Advances in Physics Theories and Applications
ISSN 2224-719X (Paper) ISSN 2225-0638 (Online)
Vol 14, 2013

Unscientific Determinism, Origin Of Universe, Free Will Theorem, Consciousness, Bell's In Equality, Standard Model Et Al.: Beatus Homo Qui Invenit Sapientiam-Tempus Edax Rerum-Tempus Fugit Models

[1]Dr. K.N.P. Kumar

Post Doctoral fellow, Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[2]Prof. B.S.Kiranagi

Distinguished Professor Emeritus of University of Mysore, Mathematics department, Manasa Gangotri, Mysore, Karnataka, India

[3]Prof. S.K. Narasimha Murthy

Chairman Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

[4]B.J.Gireesha

Associate Professor, Department of Mathematics, Jnana sahyadri, Kuvempu University, Shankaraghatta, Shimoga District, Karnataka, India

Acknowledgements: I have to acknowledge with heavy heart all those eminent and erudite connoisseurs' and ran outers who have been responsible for the work. I have put all concerted efforts for the referential integrity thereof. Should there be any act of omission, I beg pardon with utmost humbleness and humility for such an act of omissions is mentioned thereof.

Quotes:

The Buddha first taught that what we call mind or "I" is only the five skandhas of form, feelings, perceptions, impulses, and consciousness. We never get out of the suffering world because we believe that these things are real, and that they are the real "I."

Form does not differ from emptiness; emptiness does not differ from form.

Form is emptiness, emptiness is form.

No appearing, no disappearing. No taint, no purity. No increase, no decrease.

When you attain true emptiness, there is no speech or words. The Heart Sutra de-scribes our true nature by completely describing what our true nature is not. In our original nature, nothing ever appears or disappears. There is no such thing as taint or purity, because these are merely qualities created by the thinking-mind. And in original nature there is neither increase nor decrease. Our true nature is completely still and empty. It is the universal substance of which everything else is composed. Infinite in time and space, it has none of the characteristics that apply to things we can describe with speech and words.

All dharmas are marked with emptiness.

No cognition—no attainment.

In the true experience of emptiness, there are no words and no speech, so there is also no dharma. Some actual attain-ment of what these words point to is necessary. You must attain that there is actually nothing to at-tain. Everything is already truth, exactly as it is. You are already complete.

Nirvana.

Everything is suffering, which then leads to stopping suffering, stopping birth and death. This is nirvana. There are no opposites: no coming or going, no high or low, good or bad, birth or death. So in the true experience of emptiness, you perceive that there is already no birth or death, no coming or going.

Unexcelled perfect enlightenment—Anuttara samyak sambodhi.

Anuttara samyak sambodhi is a Sanskrit phrase meaning "unexcelled perfect enlightenment." It is simply another way of saying "truth." When you see, when you hear, when you smell, when you taste, when you touch, when you think—everything, just-like-this, is the truth. Before, just at the point of nirvana, there is no cognition, and no attainment with nothing to attain. So the bodhisattva depends on Prajna Paramita, and attains nirvana. But then these three words appear: Anuttara samyak sambodhi. Before, there is no attainment; now, all buddhas attain Anuttara samyak sambodhi.

If you just attain true emptiness, this is only nirvana. It is an experience of complete stillness and bliss: there is no subject or object, no good or bad, no coming or going, no life or death. There is nothing to attain. But Mahayana means your practice continues "beyond" this point, so that you attain no-attainment. You must find nirvana's function in the world. The name for that is unexcelled perfect enlightenment. If you attain no-attainment, then you attain truth. Your mind is empty and clear like space. This means your mind is clear like a mirror: If a mountain appears before the mirror, there is only mountain; water appears, and there is only water; red comes, red; white comes, white. The sky is blue. The tree is green. A dog is barking, "Woof! Woof!" Sugar is sweet. Everything that you see, hear, smell, taste, touch, and think is the truth, just as it is. Nirvana means attaining emptiness, which has nothing to attain. Anuttara samyak sambodhi means using the experience of emptiness to attain truth.

The Maha Prajna Paramita Heart Sutra Zen Master Seung Sahn (Speaking Tree TOI Collection)

Abide with me; fast falls the eventide;

The darkness deepens; Lord with me abide.

When other helpers fail and comforts flee,

Help of the helpless, O abide with me.

Swift to its close ebbs out life's little day;

Earth's joys grow dim; its glories pass away;

Change and decay in all around I see;

O Thou who changest not, abide with me.

Hymn quoted in Sakshi's letter

If we assume that the world has no beginning in time, then up to every given moment an eternity has elapsed, and there has passed away in that world an infinite series of successive states of things. Now the infinity of a series consists in the fact that it can never be completed through successive synthesis. It thus follows that it is impossible for an infinite world-series to have passed away, and that a beginning of the world is therefore a necessary condition of the world's existence.

—Immanuel Kant, First Antinomy, of Space and Time

Modern mathematics has no great problems working with infinity, but the concept still troubles some philosophers, such as Viney who argues that it is a mistake to conclude, because philosophers have been unable to answer the problems posed by the idea of an actual infinite, expounded by Kant and others, that one should not believe in an infinite past, asserting out that both metaphysical world views, that time is finite and infinite, incur paradoxes. He invokes Charles Hartshorne's principle of least paradox (As long as the problems in one's

own position are fewer than those in the positions of others, there is no justification for capitulating to the arguments of opponents.) and points out several problems with the idea of temporal finitism

One such problem is given by Hartshorne's argument against the existence of a first moment in time

Even a beginning is a change, and all change requires something changing that does not come to exist through that same change. The beginning of the world would have to happen to something other than the world, something which as the subject of happening would be in a time that did not begin with the world.

—Charles Hartshorne, Man's Vision of God, p. 233

Another, subtler, problem is that a first moment would never appear to be a first moment. Pointing to the similar arguments given by the defenders of Creation Science, and similar arguments made by Bertrand Russell, he argues that there is a paradox that infects the view that a first moment of time existed: Because every event appears to have been caused by some previous event, any first event cannot look like a first event, and so the universe must always appear to be older than it actually is. In Hartshorne's words

A first moment of time would be an ontological lie through and through, a joke of existence upon itself.

—Charles Hartshorne, Man's Vision of God, p. 234

The question of Being aims… at ascertaining the a priori conditions not only for the possibility of the sciences which examine beings as beings of such and such a type, and, in doing so, already operate with an understanding of Being, but also for the possibility of those ontologies themselves which are prior to the ontical sciences and which provide their foundations. Basically, all ontology, no matter how rich and firmly compacted a system of categories it has as its disposal, remains blind and perverted from its ownmost aim, if it has not first adequately clarified the meaning of Being, and conceived this clarification as its fundamental task. (Being and Time 3: 31) (References to Being and Time will be given in the form of 'section: page number', where 'page number' refers to the widely used Macquarrie and Robinson English translation

Whenever ontology takes for its theme entities whose character of Being is other than that of Dasein, it has its own foundation and motivation in Dasein's own ontical structure, in which a pre-ontological understanding of Being is comprised as a definite characteristic… Therefore fundamental ontology, from which alone all other ontologies can take their rise, must be sought in the existential analytic of Dasein. (Being and Time 3: 33–4 Heidegger)

What is decisive is not to get out of the circle but to come into it the right way… In the circle is hidden a positive possibility of the most primordial kind of knowing. To be sure, we genuinely take hold of this possibility only when, in our interpretation, we have understood that our first, last and constant task is never to allow our fore-having, fore-sight and fore-conception to be presented to us by fancies and popular conceptions, but rather to make the scientific theme secure by working out these fore-structures in terms of the things themselves. **(Being and Time 32: 195 Heidegger)**

The less we just stare at the hammer-thing, and the more we seize hold of it and use it, the more primordial does our relationship to it become, and the more unveiledly is it encountered as that which it is—as equipment. The hammering itself uncovers the specific 'manipulability' of the hammer. The kind of Being which equipment possesses—in which it manifests itself in its own right—we call 'readiness-to-hand'. (Being and Time 15: 98: Heidegger)

What we 'first' hear is never noises or complexes of sounds, but the creaking wagon, the motor-cycle. We hear the column on the march, the north wind, the woodpecker tapping, the fire crackling… It requires a very artificial and complicated frame of mind to 'hear' a 'pure noise'. The fact that motor-cycles and wagons are what we proximally hear is the phenomenal evidence that in every case Dasein, as Being-in-the-world, already dwells alongside what is ready-to-hand within-the-world; it certainly does not dwell proximally alongside 'sensations'; nor would it first have to give shape to the swirl of sensations to provide a springboard from which

the subject leaps off and finally arrives at a 'world'. Dasein, as essentially understanding, is proximally alongside what is understood. (Being and Time 34: 207 Heidegger)

"The wise who knows the Self as bodiless within the bodies, as unchanging among changing things, as great and omnipresent, does never grieve". "That self cannot be gained by the Veda, or by understanding, or by much learning. He whom the Self chooses, by him the Self can be gained. The Self chooses him (his body) as his own". But he who has not first turned away from his wickedness, who is not tranquil, and subdued, or whose mind is not at rest, he can never obtain the Self (even) by knowledge.

Distrust of Intellect: Katha Upanishad 1.2.22-24

He whom the world troubles not, and who troubles not the world, who is free from the emotions of joy, wrath, and fear, is dear to Me. The man who is guileless, pure, upright, unconcerned, free from distress of mind, who renounces every enterprise and worships Me, is dear to Me. He who has neither delight nor aversion, who neither mourns nor desires, who renounces good and evil fortune, and worships Me, is dear to Me. He who is the same to friend and foe, and also in honor and dishonour, who is the same in cold and heat, pleasure and pain, who is wholly free from attachment; To whom praise and blame are equal, who is silent, content with every fortune, home-renouncing, steadfast in mind, and worships Me, that man is dear to Me.

Meekness: Bhagavad Gita 12:15-19

Would he have been less full of borrowed knowledge! Then he would have accepted inspired knowledge from his father. When, with inspiration at hand, you seek book-learning, your heart, as if inspired, loads you with reproach. Traditional knowledge, when inspiration is available, Is like making ablutions in sand when water is near. Make yourself ignorant, be submissive, and then You will obtain release from your ignorance.

Masnavi: Book 4 Story 2

Sell your cleverness and buy bewilderment; Cleverness is mere opinion, bewilderment intuition.

Masnavi Book 4 Story 2

"This life of yours which you are living is not merely a piece of this entire existence, but in a certain sense the whole; only this whole is not so constituted that it can be surveyed in one single glance. This, as we know, is what the Brahmins [wise men or priests in the Vedic tradition] express in that sacred, mystic formula which is yet really so simple and so clear; tat tvam asi, this is you. Or, again, in such words as "I am in the east and the west, I am above and below, I am this entire world."

Manduka Upanishad

God isn't about making good things happen to you, or bad things happen to you. He's all about you making choices--exercising the gift of free will. God wants you to have good things and a good life, but He won't gift wrap them for you. You have to choose the actions that lead you to that life."

— Jim Butcher

It is certainly not the least charm of a theory that it is refutable; it is precisely thereby that it attracts the more subtle minds. It seems that the hundred-times-refuted theory of the "free will" owes its persistence to this charm alone; some one is always appearing who feels himself strong enough to refute it."

— Friedrich Nietzsche

If I know better than you know what I am up to, it is only because I spend more time with myself than you do."

— Daniel C. Dennett, Freedom Evolves

"Separation without separateness, reality without rift."

Charles Williams

That (pure consciousness) is full, perfect, and complete;

This (the manifest universe of matter; of names and forms) is full, perfect, complete.

Out of completion emerged completion.

And when completion emerged from completion.

Completion remained in completion.

(From the -Isa Upanishad)

Without and within all beings the unmoving and also the moving;

Because of Its subtlety, unknowable; and near and far away is That"

"And undivided, yet He exists as if divided in beings;

He is to be known as the supporter of beings;

He devours and He generates."

"No division in Consciousness is admissible at any time

As it is always one and the same.

Even the individuality of the Jiva must be known as false,

Like the delusion of a snake in a rope."

(From the various Upanishads)

The information of the whole is contained in each part.

Pribram

Order and unity are spread throughout the universe in a way which escapes our senses. In the same way that order and organisation are spread throughout the hologram. Each part of the universe contains enough information to reconstitute the whole. The form and structure of the entire universe is enfolded within each part.

Bohm

Every hundred years or so, a unique groundbreaking Copernican class volume arises unexpectedly. From ashes long thought cold of Einstein's static universe model, for the first time technically viable alternative interpretations to all pillars of Big Bang cosmology are presented in the context of a profound new “continuous-state” cosmological paradigm able to elucidate many contemporary problems plaguing the standard model of particle physics. The cosmology provides an alternative derivation of the string/brane tension formalism derived from large-scale additional dimensions that leads to a putative unique background-independent string vacuum without requiring the Higgs mechanism or SUSY superpartners. Breakthroughs presented, arising from the broad spectrum of explanatory power, include an empirical protocol for violation of the quantum uncertainty principle that provides a viable model for the implementation of universal bulk quantum computing. Other developments naturally arising from utility of the new regime include a design for constructing de Broglie-type matter-wave projectile defense shields with far-reaching consequences.

-The Holographic Anthropic Multiverse Formalizing the Complex Geometry of Reality By (author): Richard L Amoroso (Noetic Advanced Studies Institute, USA) By (author): Elizabeth A Rauscher (Tecnic Research Laboratories, USA)

Introduction:

CLASSIFICATION OF THE PROTAGONIST OR ANTAGONIST LAWS BASED ON THE CHARACTERSTICS AND PENCHANCE, PREDILECTION, PROCLIVITY AND PROPENSITIES OF THE SYSTEMS UNDER INVESTIGATION:

Towards the end of **classificational consummation, consolidation, corporation and concatenation** we take the characterstics of the systems, the predicational interiorities, ontological consonance and primordial exactitude, accolytish representations, functional topology, apocryphal aneurism and atrophied asseveration event at contracted points, and other parameters as the bastion and stylobate of the stratification purposes. Such totalistic entities would have easy paradigm of relational content, differentiated system of expressly oriented actions with primary focus and locus of homologues receptiveness and differentially instrumental activity, variable universalism and particularism, imperative compatibilities and structural variabilities, interactional dynamical orientation, institutionalization and internalisation of **pattern variables(Parsons**)common attitudinal orientation of constituionalisation of internalized dispositions, and a qualitative gradient of structural differentiation and ascribed particularistic solidarity abstraction or interactional dynamics, internal differentiation, structural morphology, comparative variability, normative aspect of expectational prediction, projection and prognostication. There are different superfield formalism and Curci-Ferrari type restrictions for different investigating systems. Equational realisation, consummation and concatenation and consubstantiation are of primal and cardinal and seminal importance Accolytish representation and other **parametricization** could also be taken in to consideration. There is no sacrosanct rule for this. It is to be remembered that there are many **investigatable systems** in the world and the equations these theories specify is applicable to them. Another example is about Off shell nil potency; Three Cf type conditions; 3-form Abelian theory; One Cf condition ; 1-form non abelian Theory B + B9bar) =I (CX C9bar) ; absolute anticommutivity are other properties that is found in many systems. Systemic characterstics thus play a primordial role in the development of the theory be it theoretical or empirical. So their characterstics could form the sentinel for the consubstantiation of the dis integrational purposes which is orderly In case of constants we talk about total invariants like the total **Schrõdinger function**, albeit we have reservations about Everett's syndrome, like for example Total gravity in the universe. However it is to be stated in unmistable terms that **Universal function** is not acceptable to the author, while instead one could talk of the various investigating systems with different conditionalities notwithstanding the fact that say a Non Linear Schrõdinger Equation holds good and the equality holds. The same is true in the Newtonian case of **Total gravity** extant in the world. Universal wave function calls for the conditionalities and functionalities of the investigating agencies and investigating systems so that those could also be taken in to consideration in the stratification scheme. Neutrinos or the protons are the same, but the source and the time factor and the space factor are also to be taken in to attribution and ascription and assignment. Two examples that could be cited are: Personality type refers to the psychological classification of different types of individuals. Personality types are sometimes distinguished from personality traits, **with the latter embodying** a smaller grouping of behavioral tendencies. Types are sometimes said **to involve** qualitative differences between people, whereas traits might be construed as quantitative differences According to type theories, for example, introverts and extraverts are two fundamentally different categories of people. According to trait theories, introversion and extraversion are part of a continuous dimension, with many people in the middle. Myers-Briggs Type Indicator (MBTI) assessment is a **psychometric questionnaire** designed to measure psychological preferences in how people perceive the world and make decisions. These preferences were extrapolated from the typological theories proposed by Carl Gustav Jung and first published in his 1921 book Psychological Types (English edition, 1923). Jung theorized that there are four principal psychological functions by which we experience the world: sensation, intuition, feeling, and thinking. One of these four functions is dominant most of the time. Equational satisfaction and consummation is of paramount and cardinal importance. We have studied discrete Schrodinger equations, NLSE, and Einstein Field equations on the same basis which must provide frame work for further analysis **In the case of Super symmetric Yang Mills Theory**

integrable systems could be taken in to consideration) The Coulomb branch of N = 2 **supersymmetric gauge theories** in four dimensions is described in general by an integrable Hamiltonian system in the holomorphic sense. A natural construction of such systems comes from two-dimensional gauge theory and spectral curves. Ground state at the critical value of the trapping potential two entanglement measures is another example. Notwithstanding the constant ground states the characterstics of the investigating systems form the bastion and stylobate of the situation. Starting from this point of view, we propose an integrable system relevant to the N = 2 SU (n) gauge theory with a hypermultiplet in the adjoint representation, and offer much evidence that it is correct. The model has an **S-duality group** (with the central element −1 of acting as charge conjugation); permutes the Higgs, confining, and oblique confining phases in the expected fashion. **Hegel** also argues strongly against the epistemological emphasis of modern philosophy from **Descartes through Kant**, which he describes as having to first establish the nature and criteria of knowledge prior to actually knowing anything, **because this would imply** an infinite regress, a foundationalism that Hegel maintains is self-contradictory and impossible. Rather, he maintains, we must examine actual knowing as it occurs in real knowledge processes. This is why Hegel uses the term "phenomenology". "Phenomenology" comes from the Greek word for "to appear", and the phenomenology of mind is thus the study of how consciousness or mind appears to itself. In Hegel's dynamic system, it is the study of the successive appearances of the mind to itself, because on examination each one dissolves into a later, more comprehensive and integrated form or structure of mind. We also study more exotic phases. **Supersymmetric Yang-Mills theory and integrable systems Ron Donagia, 1, Edward Wittenb, 2** Kind attention is also drawn to the Boltzmann arrow of time and Everett's many worlds different past and future. [Whitehead's philosophy] is provisionally the last great Anglo- American philosophy, just before the disciples of Wittgenstein spread their mists, their sufficiency and their terror. An event is not only 'a man is crushed': the great pyramid is an event, and its duration for an hour,30 minutes, 5 minutes ..., a passage of Nature, a view of God. What are the conditions for an event so that all is event? The event produces itself in a chaos, in a chaotic multiplicity, under the condition that a sort of sieve intervenes. A global methodology is assumed here. There is **Total energy and Total gravity**. Contextually reflexivity and context justification is most alluded coefficient here. As **Geroch puts it**: Theories consist of enormous number of ideas, arguments, hunches, vague feelings, all arranged in a configuration equivocation. It is the entire body of the material that is the "Theory'. (Written from memory: See Geroch 1978). **Dreyfus and Dreyfus** regard ceteris paribus condition as the universal restriction on rule systems. Classificational rules constitute prevent the regression of prevarication of such axiomatic predications and postulation alcovishness, and is a paradigmatic statement in the salient feature which does not disregard the cause effect definition and other scientific heuristic explanations. Such congressional enterprise is necessary for the totalistic view of the problem. Part of jurisprudence could be considered part of the law itself. Constitutive rules could be disregarded when the priority of the used rules are properly understood. Paradigms guide research even in the conspicuous absence of rules. Characterization **is at the heart of** naturalization. Stratification methodologies are much more vintage and montage than any other fait accompli mutatis mutandis desideratum. Manifest world in its entirety is a **holographic projection of information** embodied in its boundary. Here we are talking of the entire manifest world. But the information obtained is only for a piece of the world for which we have access. Nature's general Ledger is to be down loaded from Neuron DNA. Another factor that is taken in to consideration is **Lancan paradox and sense**, expression and event: proposition and the attribute of the state of affairs, individualistic denotation, institutionalized generalisation, personalized manifestation, organisational individuation, and corporate signification of the systems in general. According to Schrodinger actualization means to extend over a series of ordinary points, to be selected according to the rule of convergence, to be incarnated in a body, to become a state of body, and to be renewed locally namely particle coming back to its original position for the sake of limited new actualizations and extensions. It is in this sense that actualization is collective and individual, internal and external. Geometrical interpretation of the theory of differential equations clearly places in evidence two absolutely distinct realities: (a) there is a field of directions and (b**) topological accidents** which may suddenly crop up in it, as for example the existence of the plane of "singular points" to which no direction has been attached and there are integral curves with the form they take on in the vicinity of singularities of the field of directions…… The existence and distribution of singularities are notions relative to the field of vectors defined by the differential equation. The form of the integral curves **is relative to** the solution of this equation. The forms are assuredly complementary, since the nature of the field is defined by the form of integral curves in

its vicinity. But it is no less true that the field of vectors on one hand and integral curves on the other are two essentially distinct mathematical realities. In this connection of classificational process of the universal theory, it is necessary to recall what Kant had proved. Kant proves that sum of all possibilities excludes all but originary predicates and in this way constitute the completely determined concept of an individual being or objectification being. For only in this case, is a concept is equivalent to a thing. A thing completely determined in and through itself and known as the representation of an individual or a system under consideration. Thus the Universal Theory is **but the form of communication in thought between the cosmic consciousnesses, namely the nature's general ledger, and the finite individualities, namely the histories of** individual debits' and credits. It must therefore be entangled with the play of internationalities and pure retentions. Thus the universal theory has a totality. For example total gravity. Individual systems have characterstics for which say a Non Linear Schrodinger's equation holds, and such a consummation and consolidation of the equation **is built up on the corporation and actualization of the equations in simplistic and individual cases**. Compose is to make up; the parts that compose the whole. The notion of compossibility is thus defined as a continuum of the individual systems that are actualized in accord with the reductionist view which is held dearly in all subjects in this measurement world. In the Saussurean system, potential returns in language as the nonlocalizable excess of signifier over signified that accompanies the local realization of a conventional signifier-signified coupling. This excess is ensconced in the "vertical" dimension of the process (understood very differently than in the present context: as a "synchrony" rather than a temporal disparity, the disparity that is the empty form of time). It constitutes the immanent generative surplus from which the "horizontal" relay to the next linguistic realization is drawn ("diachrony": also different from the "horizontal" lines of encounter at issue here, in that diachrony is not essentially differentiating; the system remains the same across its realizations). An analogous surplus figures in Lévi-Strauss's anthropology as "mana" and its structural equivalents, which constitute the specifically cultural presentation of the staying power of processual remainder (the reserve of potential) in many societies. It is in excess-as-remainder that a reconciling of Deleuze and Guattari's thought and **structuralisms of signification might be found ("mana" converts into Deleuze's "dark precursor" or "object = x"; 1979, 315-324; 1990, 113-115; 1994, 119-123). The deleuzo-guattarian conversion of structuralism entails the signifier and signified generating an excess by escaping down a "line of flight" away from the "despotism" (self-sameness across its variations) of the signifying regime--rather than the excess regenerating signifier-signified following the chain of conventional linguistic realizations**. This amounts to an autonomization of signified-signifier coupling as a pure, "postsignifying" form in which the matter of content melds with the manner of expression. It is the melding that takes the lead, as an "abstract machine" of autonomized, depersonalized expression (Deleuze/Guattari 1986, 3-8; Deleuze/Guattari 1987, 129-135, 189-191; Haghighi. The triad thesis, antithesis, synthesis is often used to describe the thought of German philosopher Georg Wilhelm Friedrich Hegel. Hegel never used the term himself, and almost all of his biographers have been eager to discredit it. The triad is usually described in the following way: The **thesis** is an intellectual proposition. The **antithesis is simply** the negation of the thesis, a reaction to the proposition. The synthesis solves the conflict between the thesis and antithesis by reconciling their common truths and forming a new thesis, starting the process over. According to Walter Kaufmann, **although the triad is often thought to form part of an analysis of historical and philosophical progress called the Hegelian dialectic**, the assumption is erroneous. Hegel used this classification only once, and he attributed the terminology **to Immanuel Kant**. The terminology was largely developed earlier by the neo-Kantian Johann Gottlieb Fichte, also an advocate of the philosophy identified as German idealism. The triad thesis, antithesis, synthesis is often used to describe the thought of German philosopher Georg Wilhelm Friedrich Hegel. Hegel stresses the paradoxical nature of consciousness; he knows that the mind **wants to know** the whole truth, but that it cannot think **without drawing a** distinction. Unfortunately, every distinction has two terms, every argument has a counter-argument, and consciousness **can only focus on one of these at a time**. So it fixes first on the one, then under pressure fixes second on the other, until it finally comes to rest on the distinction itself. Hegel refers to this **process of alternation and rest as dialectic**. In other words, the dialectical method **involves the** notion that the form of historical movement, process or progress, **is the result of** conflicting opposites. Thus this area of Hegel's thought **has been broken down in terms** of the categories of thesis, antithesis, and synthesis. Hegel's philosophy of history embraces the concept that a conflict of opposites is a struggle between actual and potential worlds. Kant's first Critique concerns itself with determining the structure of the objects of experience. His transcendental logic allows **an**

empirical manifold, or multiplicity, to be grasped as a unified object, but only insofar as its unity is provided by the subject (via apperception). The transcendental ego synthesizes manifolds that remain distinct in kind from it, and thus its logic can only provide the a priori conditions for possible experiences, not the generative conditions for actual ones. "Kant's main preoccupation is therefore with the validity of propositions given in advance of our enquiry", for it merely seeks the conditions for subjective knowledge of an object, which are presumed to be identical with the conditions of objects themselves. Limited to schematic overviews, brief asides or even dismissive caricatures these philosophies muster enough strength stride by stride with present day theories(Henry Somers-Hall, Hegel, Deleuze, and the Critique of Representation: **Dialectics of Negation and Difference, State University of New York Press, 2012, 289pp., (hbk), ISBN 9781438440095. Reviewed by Jim Vernon, York University)** Hegel shows how the various categories of logic sublate each other, turning contradiction into the motor that drives, rather than the problematic limit that haunts, representational thinking. The extended physical and biological theory is not just a multidimensional consciousness theory, but it is, after and above all, a philosophical theory. But their connection is not hierarchical. The self-created structural and structure less system of the multi-theoretical components is necessarily (in reality and symbolically also, a multi-centred, in- and out- infinitely expanded, multi-layered spherical) logical and logical spacetime theory. But to fully unfold this concept, neither physics, nor any other partial theory is capable in itself. So, the different unified theories together are cannot be other than the wholeness theory, the unified top theory or the philosophical metatheory, but is this not just to give a name, to explain something, would this be the Creation/Non-creation itself? **On Quantum's Universality: Posted on 5/19/2004 by Dave Bacon (The Quantum Pontiff):** Often when I am thinking about the foundations of quantum theory, I am struck by the universality of the theory. Quantum theory (or its related cousin, quantum field theory) applies generically to all physical systems (disregarding the transition to some "classical" theory and of course, difficulties with both QCD and gravity.) Thus we apply quantum theory to our basic theories of physics, electromagnetism, the weak force, the strong force, but we also apply quantum theory to simple atoms and complex molecules, to single electrons and electron gases in metals, etc. Quantum theory is the universal language we use to describe any physical process. If we are thinking about ways to explain quantum theory, then this universality is a bit mysterious: the explanation had better apply to all of these different physical systems and that seems like a lot of work! Of course, this reasoning is flawed: it seems the universality is an illusion. The reason we can describe a complex molecule by quantum theory is that the fundamental constituents of that molecule obey quantum theory. Separation of different energy scales (and other scales, like localibility) allow us to ignore some of the constituents details, and the complex system behaves like a quantum system. So really any explanation of quantum theory need only apply to some basic level of physics (where this level is I refuse to speculate.) While quantum theory appears mysteriously universal, this is an illusion for those persuing understanding the mystery of the quantum. Whitehead was a mathematician as well as philosopher. Metaphysics was his main interest. He said it is the task of philosophy "to frame a coherent, logical, necessary system of general ideas in terms of which every element of our experience can be interpreted

The triad is often said to have been extended and adopted by Karl Marx and Friedrich Engels, however, Marx referred to them in The Poverty of Philosophy as speaking Greek and "Wooden trichotomies" .The classic empiricist presentation of this approach is William James's Essays in Radical Empiricism: "There is no general stuff of which appearance is made. It is made of that, or just what appears (26-27) ... what I call **'pure experience' [is] only virtually or potentially either object or subject as yet.** For the time being, it is plain, unqualified actuality, a simple that (23) ... experience as a whole is self-containing and leans on nothing (193) ... **object and subject fuse in the fact of 'presentation' or sense-perception (197) ... knower and object exist as so many ultimate thats or facts of being**" (196). As the last phrase suggests, James's "that" (or "this-that": to signal a differential, or incipient difference) is Francis Bacon's "fact," which can in turn be understood in terms of Whitehead's definition of fact as the "undifferentiated terminus of sense-awareness," the inexhaustible unthought from which thought "diversifies" and to which it "demonstratively" returns, as to its "ideal limit" (6-10, 13-15. **On the "univocity" of the coming to expression of the multiple, see Deleuze 1990, 177-180; 1994, 35-42, 303-304.** Can the nature of thinking be uncovered by thinking alone? (Cf consciousness) "My dialectical method is, in its foundations, not only different from the Hegelian, but exactly opposite to it. For Hegel, the process of thinking, which he even transforms into an independent subject, under the name of 'the idea ', is the creator of the real world, and the real world is only the external appearance of the idea. With me,

the reverse is true: the ideal is nothing but the material world reflected in the mind of man, and translated into forms of thought. "Karl Marx, Capital, vol. 1, Preface to the Second Edition. (P.102) (See Ideal / Real) For Aristotle, Thinking is the one specific activity of the Human Soul Which is Capable of Independent Existence and separate from any connection to the Body. A Strenuous counterargument has been made, precisely for body thinking. Deleuze and Guattari for, "Becoming Animal" and "Becoming intense" are Modes of resistance to abstract Thinking. Alain Badiou FOLLOWS Aristotle in using thinking **to define** specifically the Human, as Opposed to Interest. For Badiou, "the capacity which is specifically human is that of thought, and thought is nothing other than that by which the path of a truth seizes and traverses the human animal." **(Infinite Thought, p. 71)** In HIS Essay "Does Consciousness Exist?" **William James** Pointed out That Consciousness is a Process, and not a substance. (Unlike Descartes, who claimed the existence of a res cogitans?) In The Principles of Psychology, **James described five properties of what he called "thought**." Every thought, he wrote, tends to be **part of personal consciousness**. We write this despite the definition that has been held for the consciousness as a storage room and mind anagrammatically acting upon it. Concomitant quantum channeling like synapses and neuron firing might take place the in the brain. And that falls under the subterranean realm and ceratoid dualism of Neurology. Thought is always changing, is sensibly continuous, and appears to deal with objects independent of it. In addition, thought focuses on some objects to the exclusion of others. In other words, it involves attention. Alfred North Whitehead claimed that, with this Inquiry, James was to the Twentieth century What Descartes was to the Seventeenth. Louis Sullivan describes "Real Thinking," Thinking in the present tense, as Organic Thought. "It is in the present, only, that you really live, therefore it is in the present, only that you can really think. And in this sense you think organically. Pseudo-thinking is inorganic. The one is living, the other dead. The present is the organic moment, the living moment. "(Kindergarten Chats, "Thought.") **Thinking is a form of Computation? Emergence ---------- Emergence Refers to the appearance of Patterns of Organization and is one of the Key concepts of Complexity and a -life. It is sometimes referred to as a situation where the whole is greater than** the sum of the parts, because it cannot be analyzed by taking the parts apart and examining them separately. One reason for this is that in a complex phenomenon **showing** emergent properties, the parts become a determining context for each other, and these patterns of feedback **contribute to the** appearance of the emergent phenomenon. For Michael Polanyi, "**Evolution can be understood only as a feat of emergence**." For Goethe, the Whole Living was Also More than the Sum of ITS parts. "Das ist zwar Lebendige in Elemente zerlegt, aber MAN Kann ES aus Diesen nicht wieder zusammenstellen und beleben." Goethe quoted in d'Arcy Thompson, **On Growth and Form. (P.41)** But for most contemporary scientists, Goethe's meaning of emergence was a mystical one. For Francis Crick, the scientific meaning of emergent assumes that while the **whole cannot be the simple sum of the separate parts, its behavior can, at least in principle, be understood from the nature and behavior of its parts plus the knowledge of how all these parts interact. (The Astonishing Hypothesis, P.11)** (Assuming IT's Possible to have **That Laplacian Knowledge**) Stuart Kauffman Suggests That many Properties of Organisms May be Probable **emergent Collective Properties** of Their constituents. (This is precisely Kant's definition of the organic - as the result of internal interactions instead of an assembly of preexisting parts. (See Mechanism / Vitalism) "The problem of origins requires an understanding of how new levels of order emerge from complex patterns of interactions and what the properties of these emergent structures are in terms of their robustness to perturbation and their capacities for self-maintenance **"(Brian Goodwin, How the Leopard Changed its Spots, p. 181.)** Luc Steels Offers Useful definitions of emergence in Artificial Life half. He distinguishes between first order emergence, defined as a property not explicitly programmed in, and second-order emergence, defined as an emergent behavior that adds additional functionality in the system. In general a-life researchers try to create second -order emergence, then the system for use CANS ITS own emergent Properties to create an upward Spiral of Continuing evolution and emergent behaviors. For a Critique of emergence, See Peter Cariani, "Emergence and adaptivity in Organisms and Devices," World Fudtures 32 (1991):111-32. C. Lloyd Morgan's Descriptions of emergence as a Qualitative Change in direction closely resemble the definitions of bifurcations. HIS In Emergent Evolution, of 1923, he writes, "**The emergent step, although it may seem more or less saltatory, is best Regarded as a Qualitative Change of direction, or Critical turning-Point, in the Course of events**. " Daniel Stern describes the Infant's sense of Self between Birth and two months as the "emergent Self." (See subject). It is a sense of the Coming-into-Being of Organization and Remains active for the rest of life. But IT is not an overarching schema About Integrating the Self, but rather an Experience of

Process. (cf Molar / Molecular) In Gestalt Perception terms, emergence refers to our ability to perceive forms that cannot be reduced to the addition of the "atomic" parts of a pattern. Ego superimposing two squares diagonally displaced **allows for** readings of an emergent square at their overlap, two emergent L-shapes in the non-Intersecting area, in addition to the two Squares. (cf. Figure / Ground) A **Fundamental property of Open Systems is that they stabilize any improbability which Serves to elicit them**. Being / Becoming --------- ---------- According to Ernst Cassirer, Plato's Philosophy recognizes two Contrary Forms of representation, one of Which is Valid for the Realm of Being and the Other for the Realm of Becoming. (Individual and Cosmos, p.125) For Cassirer, "**Form Thinking**" Belongs to Being, while "causal Thinking" Belongs to Becoming. But strict Knowledge is Only Possible of the Always-Being. Becoming that which is CAN only be described, if at All, in the language of myth. Or rather, myth is already familiar with both the question of the "what" and the question of the "whence." "It sees everything that it grasps (the world as well as the gods) under this double aspect."(The Problem of Form and the Problem of Cause, in The Logic of the Cultural Sciences, p. 87) Socrates first Discovered the Concept, or Eidos as the relation Between the Particular and the General and as a germ of a new meaning of the general question concerning being. This meaning emerged in its full purity when the Socratic eidos went on to unfold into the (transcendental) Platonic "Idea." (See transcendence / immanence) For Plato, the soul is an intermediary between being and Becoming. The Aristotelian system seemed to Promise a Different reconciliation of the Opposition between Being and Becoming. Unlike Plato and the thinkers of pure form, Aristotle wanted to restore to Becoming ITS rightful Place, Because He was convinced **That Only in this way Could Philosophy be Transformed from a Mere Theory of concepts into a Theory of Reality. Form and Matter, Being and Becoming, must become correlative. Peculiarly Aristotelian the Concept of the formal Cause originates from this fusion**. (Cassirer) "In Nature We May Search for successfully something that endures, and in nature we may regard becoming as if any phase of it were the "reason" of a later phase and the "consequence" of an earlier one. We may even go so far as to say: Nature is the one mediate object that obeys the postulates of the rational theory of becoming. "Hans Driesch, History and Theory of Vitalism, p. 193. We are grateful to the Russian Blog that was available on Google search and Wikipedia in the consolidation and consummation of the above facts. Gilles Deluze is another person whose Pure Immanence and The Logic of sense has been helpful. Part of it expatiated and enucleated from blog Интернет версия данной статьи находится по адресу: http://www.situation.ru/app/j_art_1041.htm Copyright (c) Альманах "Восток"

One might compare Hegel's point here to that expressed by Kant in his well known claim that without concepts, those singular and immediate mental representations he calls “intuitions” are “blind.” In more recent terminology one might talk of the “concept-” or “theory-ladenness” of all experience, and the lessons of this chapter have been likened to that of Wilfred Sellars's famous criticism of the “myth of the given” (Sellars 1997). Hegel has insisted on quarantining his defensible ideas from it (e.g., Wood 1990). However, rather than be understood as a treatise in formal (or “general”) logic, it is perhaps best understood as a version of what Kant had called “transcendental logic,” and in this sense thought of as a successor to Kant's “transcendental deduction of the categories” in the Critique of Pure Reason in which Kant attempted to “deduce” a list of those non-empirical concepts, the “categories,” which he believed to be presupposed by all empirical judgments made by finite, discursive thinkers. However, whereas the latter’s is largely etymologically oriented and given to exploiting semantic ambiguities, traces and aporias, Deleuze’s strategy is more geared towards conceptual and functional differentiation, exploring the horizons of Ideas (in the Kantian sense) and bringingforth the machinic and operative features of the philosophies with which he engages. hile, for Lacan, Truth is this shattering experience of the Void - a sudden insight into the abyss of Being, "not a process so much as a brief traumatic encounter, or illuminating shock, in the midst of common reality" -, for Badiou, Truth is what comes afterward: the long arduous work of fidelity, of enforcing a new law onto the situation. 5 The choice is thus: "whether a vanishing apparition of the real as absent cause (for Lacan) or a forceful transformation of the real into a consistent truth. doctrine is precisely that, while never ceasing to be dialectical in pinpointing the absent cause and its divisive effects on the whole, it nevertheless remains tied to this whole itself and is thus unable to account for the latter's possible transformation. /.../ Surely anchored in the real as a lack of being, a truth procedure is that which gives being to this very lack. Pinpointing the absent cause or constitutive outside of a situation, in other words, remains a dialectical yet idealist tactic, unless this evanescent point of the real is forced, distorted, and extended, in order to give consistency to the real as a new generic truth. Bosteels describes the modality of the truth-procedure: Setting out from the void which prior to the event remains indiscernible in

the language of established knowledge, a subjective intervention **names** the event which disappears no sooner than it appears; /it/ faithfully connects as many elements of the situation as possible to this name which is the only trace of the vanished event, and **subsequently forces** the extended situation from the bias of the new truth as if the latter were indeed already generally applicable. Bosteels basic reproach, according to which, psychoanalysis collapses into an instantaneous act what is in reality an ongoing and impure procedure, which from a singular event leads to a generic truth by way of a forced return upon the initial situation. Whereas for Zizek, the empty place of the real that is impossible to symbolize is somehow already the act of truth itself, for Badiou a truth comes about only by forcing the real and by displacing the empty place, so as to make the impossible possible. 'Every truth is post-evental,' Badiou writes. An example is Descartes' celebrated phrase at the beginning of the Discourse on the Method: Good sense is the most evenly shared thing in the world. . the capacity to judge correctly and to distinguish the true from the false, which is properly what one calls common sense or reason, is naturally equal in all men . . For Deleuze, is an image of thought? Although images of thought take the common form of an 'Everybody knows . . .' (DR 130), they are not essentially conscious. Rather, they operate on the level of the social and the unconscious, and function, "all the more effectively in silence." (DR 167). Relationship of philosophy to thought must have two correlative aspects, Deleuze argues: an attack on the traditional moral image of thought, but also a movement towards understanding thought as self-engendering, an act of creation, not just of what is thought, but of thought itself, within thought (DR 147). he thought which is born in thought, the act of thinking which is neither given by innateness nor presupposed by reminiscence but engendered in its genitality, is a thought without image. But what is such a thought, and how does it operate in the world? (DR 167; cf. 132) Durkheim's concepts of the collective representation and the process of universalization, then, **correspond to** Kantian concepts, and Durkheim acknowledges the relevance of Kantian philosophy to his epistemological and moral considerations (1965: 494). It is to be noted and stated in unmistakable and unequivocal words that the singularities and the events constitute a life coexist with the accidents of life that corresponds to it, but they are neither grouped nor divided in the same way. They connect in a manner entirely different from how individuals connect by virtue of which classification takes place. A singular life might do without any individuality, without any other concomitant that individualizes it. Individual life or that of insentient life remains inseparable from empirical determinations. On the other hand, the indefinite as such is the mark of not an empirical indetermination but of a determination by immanence or a transcendental determinability. (**Pure Immanence Gilles Deluze: Page number 30).** It is to be realized that a phenomenon which we think out is engaged in process of actualization following the plane that gives it its particular reality. Plane of immanence therefore is actualization, as long as the events that populate them are localised by a law or other theorem tic corollary.

Deleuze uses the introduction to clarify the term "repetition." Deleuze's repetition can be understood by contrasting it to generality. Both words describe events that have some underlying connections. Generality refers to events that are connected through cycles, equalities, and laws. Most phenomena that can be directly described by science are generalities. Seemingly isolated events will occur in the same way over and over again because they are governed by the same laws. Water will flow downhill and sunlight will create warmth because of principles that apply broadly. In the human realm, behavior that accords with norms and laws counts as generality for similar reasons. Science deals mostly with generalities because it seeks to predict reality using reduction and equivalence. Repetition, for Deleuze, can only describe a unique series of things or events. The Borges story in which Pierre Menard reproduces the exact text of Don Quixote is a quintessential repetition: the repetition of Cervantes in Menard takes on a magical quality by virtue of its translation into a different time and place. Art is often a source of repetition because no artistic use of an element is ever truly equivalent to other uses. (Pop Art pushes this quality to a certain limit by bringing production near the level of capitalism.) For humans, repetition is inherently Transgressive. As in Coldness and Cruelty, Deleuze identifies humor and irony as lines of escape from the generalities of society. Humor and irony are in league with repetition because they create distance from laws and norms even while re-enacting them. Deleuze describes repetition as a shared value of an otherwise rather disparate trio: Kierkegaard, Nietzsche, and Péguy. He also connects the idea to Freud's death drive. He goes on to define repetition as "difference without a concept" . Repetition is thus reliant on difference more deeply than it is opposed. Further, profound repetition will be characterized by profound difference.

Difference in Itself

Deleuze paints a picture of philosophical history in which difference has long been subordinated to four pillars of reason: identity, opposition, analogy, and resemblance. He argues that difference has been treated as a secondary characteristic which emerges when one compares pre-existing things; these things can then be said to have differences. This network of direct relations between identities roughly overlays a much more subtle and involuted network of real differences: gradients, intensities, overlaps, and so forth .The chapter contains a discussion of how various philosophers have treated the emergence of difference within Being. This section uses Duns Scotus, Spinoza, and others to make the case that "there has only ever been one ontological proposition: Being is univocal. ... A single voice raises the clamor of being" . One then tries to understand the nature of differences that arise within Being. Deleuze describes how Hegel took contradiction—pure opposition—to be the principle underlying all difference and consequently to be the explanatory principle of the entire world's texture. He accuses this conception of having a theological and metaphysical slant.

Deleuze proposes (citing Leibniz) that difference is better understood through the use of dx, the differential. A derivative, dy/dx, determines the structure of a curve while nonetheless existing just outside the curve itself; that is, by describing a virtual tangent. Deleuze argues that difference should fundamentally be the object of affirmation and not negation. As per Nietzsche, negation becomes secondary and epiphenomenal in relation to this primary force.. Repetition for Itself: The chapter describes three different levels of time within which repetition occurs. Deleuze takes as axiomatic the notion that there is no time but the present, which contains past and future. These layers describe different ways in which past and future can be inscribed in a present. As this inscription grows more complicated, the status of the present itself becomes more abstract. Basic processes of the universe have a momentum that they carry into each present moment. A 'contraction' of reality refers to the collection of a diffuse ongoing force into the present. Prior thought and behavior, all substance performs contraction. "We are made of contracted water, earth, light, and air...Every organism, in its receptive and perceptual elements, but also in its viscera, is a sum of contractions, of retentions and expectations. Passive synthesis is exemplified by habit. Habit incarnates the past (and gestures to the future) in the present by transforming the weight of experience into an urgency. Habit creates a multitude of "larval selves," each of which functions like a small ego with desires and satisfactions. In Freudian discourse, this is the domain of bound excitations associated with the pleasure principle. Deleuze cites Hume and Bergson as relevant to his understanding of the passive synthesis. Active synthesis: The second level of time is organized by the active force of memory, which introduces discontinuity into the passage of time by sustaining relationships between more distant events. A discussion of destiny makes clear how memory transforms time and enacts a more profound form of repetition: Destiny never consists in step-by-step deterministic relations between presents which succeed one another according to the order of a represented time. Rather, it implies between successive presents non-localisable connections, actions at a distance, systems of replay, resonance and echoes, objective chances, signs, signals, and roles which transcend spatial locations and temporal successions. Relative to the passive synthesis of habit, memory is virtual and vertical. It deals with events in their depth and structure rather than in their contiguity in time. Where passive syntheses created a field of 'me's,' active synthesis is performed by 'I.' In the Freudian register, this synthesis describes the displaced energy of Eros, which becomes a searching and problematizing force rather than a simple stimulus to gratification.

Empty time

The third layer of time still exists in the present, but it does so in a way that breaks free from the simple repetition of time. This level refers to an ultimate event so powerful that it becomes omnipresent. It is a great symbolic event, like the murder to be committed by Oedipus or Hamlet. Upon rising to this level, an actor effaces herself as such and joins the abstract realm of eternal return. The me and the I give way to "the man without name, without family, without qualities, without self or I...the already-Overman whose scattered members gravitate around the sublime image" Empty time is associated with Thanatos, a desexualized energy that runs through all matter and supersedes the particularity of an individual psychic system. Deleuze is careful to point out that there is no reason for Thanatos to(eb) produce a specifically destructive impulse or 'death instinct' in the subject; he conceives of Thanatos as simply indifferent.

The Image of Thought

This chapter takes aim at an "image of thought" that permeates both popular and philosophical discourse. According to this image, thinking naturally gravitates towards truth. Thought is divided easily into categories of truth and error. The model for thought comes from the educational institution, in which a master sets a problem and the pupil produces a solution which is either true or false. This image of the subject supposes that there are different faculties, each of which ideally grasps the particular domain of reality to which it is most suited. In philosophy, this conception results in discourses predicated on the argument that "Everybody knows..." the truth of some basic idea. Descartes, for example, appeals to the idea that everyone can at least think and therefore exists. Deleuze points out that philosophy of this type attempts to eliminate all objective presuppositions while maintaining subjective ones. Deleuze maintains, with Artaud, that real thinking is one of the most difficult challenges there is. Thinking requires a confrontation with stupidity, the state of being formlessly human without engaging any real problems. One discovers that the real path to truth is through the production of sense: the creation of a texture for thought that relates it to its object. Sense is the membrane that relates thought to its other. Accordingly, learning is not the memorization of facts but the coordination of thought with a reality. "As a result, 'learning' always takes place in and through the unconscious, thereby establishing the bond of a profound complicity between nature and mind". Deleuze's alternate image of thought is based on difference, which creates a dynamism that traverses individual faculties and conceptions. This thought is fundamentally energetic and asignifying: if it produces propositions, these are wholly secondary to its development.

At the end of the chapter, Deleuze sums up the image of thought he critiques with eight attributes: the postulate of the principle, or the Cogitatio natural universalis (good will of the thinker and good nature of thought); the postulate of the ideal, or common sense (common sense as the concordia facultatum and good sense as the distribution which guarantees this concord); the postulate of the model, or of recognition (recognition inviting all the faculties to exercise themselves upon an object supposedly the same, and the consequent possibility of error in the distribution when one faculty confuses one of its objects with a different object of another faculty); the postulate of the element or of representation (when difference is subordinated to the complimentary dimensions of the Same and the Similar, the Analogous and the Opposed; the postulate of the negative, or of error (in which error expresses everything which can go wrong in thought, but only as the product of external mechanisms); the postulate of logical function, or the proposition (designation is taken to be the locus of truth, sense being no more than the neutralized double or the infinite doubling of the proposition); the postulate of modality, or solutions (problems being materially traced from propositions or indeed, formally defined by the possibility of their being solved); the postulate of the end, or result, the postulate of knowledge (the subordination of learning to knowledge, and of culture to method.

It is to be stated that we are here reproducing the following for the bastion and stylobatishness of the necessity of classificational procedural formalities and I donot believe in whatever that has been stated. Nor is it meant as propaganda. It is purely definitional with axiomatic predications and doctrinal manifestations. In Vedas and Upanishads the classification processual formalities of Universal action is undertaken thus with a mention that the proportion of the ones mentioned would be different in different individuals and individuals themselves are investigating systems notwithstanding a wider classification which is nothing but more than a Universal Law. Manduka Upanishad divides all knowledge into two categories. The knowledge that leads to Self Realization is called Para Vidya (Great or Divine Knowledge) and everything else is called Apara Vidya or Knowledge of Material world (wordly knowledge). Shaunaka approaches sage Angiras and asks "Revered Sir, by knowing what everything will be known?" Angiras replies that two know ledges should be known, one is **Para Vidya** and other is **Apara Vidya.** Knowledge of worldly things is Apara Vidya and that by which Eternal Truth or Akshara is obtained is Para Vidya. Though Apara Vidya enables one to earn ones bread and helps one to understand each object of universe separately, it does not show the Ultimate Reality (Akshara) or Root Cause of this universe. While Para Vidya doesn't teach objects of this universe but enables one to understand underlying fabric of it. Like by knowing gold all the gold ornaments could be known, by knowing Akshara, it's another manifestation, the universe is known. This Upanishad expounds the greatness of Para. To think whether a certain thing may be eaten is a thought-form of the mind; "It is good. It is not good. It can be eaten. It cannot be eaten", discriminating notions like these constitute the discriminative intellect. Because the mind alone

constitutes the root-principle manifesting as the individual, God and the world, its absorption or submergence and dissolution in the Self as pure Consciousness is the final emancipation known as Kaivalya and in the Supreme Spirit, the Brahman.(Ramana maharishi and Wikipedia). **Tamas (Darkness):** Complete delusion, ignorance, illiberality, indecision in respect of action, sleep, haughtiness, fear, cupidity, grief, censure of good acts, loss of memory, unripeness of judgment, absence of faith, violation of all rules of conduct, want of discrimination, blindness, vileness of behaviour, boastful assertions of performance when there has been no performance, presumption of knowledge in ignorance, unfriendliness (or hostility), evilness of disposition, absence of faith, stupid reasoning, crookedness, incapacity for association, sinful action, senselessness, stolidity, lassitude, absence of self-control, degradation, - all these qualities are known as belonging to Darkness **(Tamas).** Whatever other states of mind connected with delusion exist in the world, all appertain to Darkness. Frequent ill-speaking of other people, censuring the deities and the Brahmanas (priests), illiberality, vanity, delusion, wrath, unforgiveness, hostility towards all creatures, are regarded as the characteristics of Darkness. Whatever undertakings exist that are unmeritorious (in consequence of their being vain or useless), what gifts there are that are unmeritorious (in consequence of the unworthiness of the done, the unreasonableness of the time, the impropriety of the object, etc.), vain eating, - these also appertain to **Darknesss (Tamas).** Indulgence in calumny, unforgiveness, animosity, vanity, and absence of faith are also said to be characteristics of Darkness. Whatever men there are in this world who are characterised by these and other faults of a similar kind, and who break through the restraints provided by the scriptures, are all regarded as belonging to the quality of Darkness. Here Rishis and Munis (Seers and sages), and deities become deluded, desirous of pleasure. Darkness, delusion, the great delusion, the great obscurity called wrath, and death, that blinding obscurity (these are the five great afflictions). As regards wrath, that is the great obscurity (and not aversion or hatred as is sometimes included in the list). With respect then to its colour (nature), its characteristics, and its source, I have, ye learned Brahmanas, declared to you, accurately and in due order, everything about the quality of Darkness (Tamas). Who is there that truly understands it? Who is there that truly sees it? That, indeed, is the characteristic of Darkness, viz., the beholding of reality in what is not real. The qualities of Darkness have been declared to you in various ways.

Rajas (Passion): Injuring others, beauty, toil, pleasure and pain, cold and heat, lordship (or power), war, peace, arguments, dissatisfaction, endurance, might, valour, pride, wrath, exertion, quarrel, jealousy, desire, malice, battle, the sense of meum or mineness, protection of others, slaughter, bonds, and affliction, buying and selling, lopping off, cutting, piercing and cutting off the coat of mail that another has worn, fierceness, cruelty, vilifying, pointing out the faults of others, thoughts entirely devoted to worldly affairs, anxiety, animosity, reviling of others, false speech, false or vain gifts, hesitancy or doubts, boastfulness of speech, praise and criticisms, laudation, prowess, defiance, attendance (as on the weak and the sick), obedience (to the commands of preceptors and parents), service or ministrations, harbouring of thirst or desire, cleverness or dexterity of conduct, policy heedlessness, contumely, possessions, and diverse decorations that prevail in the world among men, women, animals, inanimate things, houses, grief, incredulousness, vows and regulations, actions with expectation (of good result), diverse acts of public charity, the rites in respect of Swaha salutations, rites of Swadha and Vashat, officiating at the sacrifices of others, imparting of instruction, performance of sacrifices, study, making of gifts, acceptance of gifts, rites of expiation, auspicious acts, the wish to have this and that, affection generated by the merits of the object for which or whom it is felt, treachery, deception, disrespect and respect, theft, killing, desire of concealment, vexation, wakefulness, ostentation, haughtiness, attachment, devotion, contentment, exultation, gambling, indulgence in scandal, all relations arising out of women, attachment to dancing, instrumental music and songs – all these qualities have been said to belong to the quality of Passion (Rajas). Those men on earth who meditate on the past, present and the future, who are devoted to the aggregate of the three viz., Religion, Wealth and Pleasure, who acting from impulse of desire, exult on attaining to affluence in respect of every desire, are said to be enveloped by Passion (Rajas). These men have downward courses. Repeatedly reborn in this world, they give themselves up to pleasure. They covet what belongs to the world as also all those fruits that belong to the world hereafter. They make gifts, accept gifts, offer oblations to the Pitris, and pour libations on the sacrificial fire.

Sattwa (Goodness): Brahma (the Grandsire Prajapati) said: Sattwa is beneficial to all creatures in the world, and unblamable, and constitutes the conduct of those that are good. Joy, satisfaction, nobility, enlightenment,

and happiness, absence of stinginess, absence of fear, contentment, disposition for faith, forgiveness, courage, abstention from injuring any creature, equability, truth, straightforwardness, absence of wrath, absence of malice, purity, cleverness, prowess- these appertain to the quality of Goodness (Sattwa).He who is devoted to the duty of Yoga, regarding knowledge to be vain, conduct to be vain, service to be vain, and mode of life to be vain, attains to what is highest in the world hereafter. Freedom from the idea of meum (mineness), freedom from egoism, freedom from expectations, looking upon all with an equal eye, and freedom from desire, - these constitute the eternal religion of the good. Confidence, modesty, forgiveness, renunciation, purity, absence of laziness, absence of cruelty, absence of delusion, compassion to all creatures, absence of the disposition to calumniate, exultation, satisfaction, rapture, humility, good behaviour, purity in all acts having for their object the attainment of tranquillity, righteous understanding, emancipation from attachments, indifference, Brahmacharya (celibacy), complete renunciation, freedom from the idea of meum, freedom from expectations, unbroken observance of righteousness, beliefs that gifts are vain, sacrifices are vain, study is vain, vows are vain, acceptance of gifts is vain, observance of duties is vain, and penances are vain – those Brahmanas (priests) in this world, whose conduct is marked by these virtues, who adhere to righteousness, who abide in the Vedas, are said to be wise and possessed of correctness of vision.[Note: Compare from The Mahabharata, Aswamedha Parva, and Sec.XLIV): Brahma said: The Unmanifest is the source of all the worlds as, indeed, that is the end of everything. Days end with the sun's setting and Nights with the sun's rising. The end of pleasure is always sorrow, and the end of sorrow is always pleasure. All accumulations have exhaustion for their end, and all ascent have falls for their end. All associations have dissociations for their end, and life has death for its end. All action ends in destruction and all that is born is certain to meet with death. Every mobile and immobile thing in this world is transient. Sacrifice, gift, penances, study, vows, observances, - all these have destruction for their end. Of Knowledge, there is no end. Hence, one that is possessed of a tranquil soul, that has subjugated his senses, that is freed from the sense of meum (mineness), that is devoid of egoism, is released from all sins by pure knowledge.] Casting off all sins and freed from grief, those men possessed of wisdom attain to Heaven and create diverse bodies for themselves. Attaining the power of governing everything, self-restraint, minuteness, and these high-souled ones make operations of their own mind, like the gods themselves dwelling in Heaven. Such men are said to have their courses directed upwards. They are veritable gods capable of modifying all things. Attaining to Heaven they modify all things by their very nature. They get whatever objects they desire and enjoy them. Thus have I, ye foremost of regenerate ones, described to you what that conduct is which appertains to the quality of Goodness (Sattwa). Understanding this duly, one acquires whatever objects one desire. The qualities that appertain to Goodness have been declared particularly. The conduct which those qualities constitute has also been properly set forth. That man, who always understands these qualities, succeeds in enjoying the qualities without being attached to them. **Sattwa, Rajas & Tamas** Whatever object exists in this world, everything in it is fraught with the three qualities.

It is to be noted that the primary focus and locus of homologous repetitiveness and differentially instrumental activity we are talking about is the application of some certain Universal laws should not be mistaken for the Universalistic laws and particularistic laws themselves. All we are talking about is the application of paradigmatic relational content to specific investigating systems under consideration**.** On the other hand there certainly exist **Universalistic** Laws and Particularistic Laws. We are not in from any point of view talking about the imperative compatibilities and structural variabilities of the two different forms of laws but the application of say Einstein's Law of mass energy equivalence to some certain galaxy which is under investigation. Following essay allays and ameliorates the vestiges of doubt in the eventuality of confoundment between the two schools of thought we have essayed. It is simple application systems that are also investigatory and conforms and congruenential with the given Einsteinian Theory. **This universalism which is not one Linda M. G. Zerilli From: Diacritics Volume 28, Number 2, summer 1998 pp. 3-20 | 10.1353/dia.1998.0013 In lieu of an abstract, here is a brief excerpt of the content: Diacritics 28.2 (1998) 3-20 Review Essay Ernesto Laclau. Emancipation(s). London: Verso, 1996.** Judging from the recent spate of publications devoted to the question of the universal, it appears that, in the view of some critics, we are witnessing a reevaluation of its dismantling in twentieth-century thought. One of the many oddities about this "return of the universal" is the idea that contemporary engagements with it are more or less of a piece, and that they reflect a growing consensus that poststructuralist political theories are incapable of generating a viable alternative to the collective fragmentation that characterizes late modernity. The putative return to the universal marks, on this view, both a

homecoming to Enlightenment ideals -- purified of their more poisonous elements, of course -- and a reconciliation of sorts between those who refuted these ideals and those who sought to realize them. Now that "we" all know and agree that post structuralism is critically valuable but politically bankrupt; now that we all know and agree that the "old universal" was indeed a "pseudo-universal," so the homecoming narrative goes; we can get on with the project of constructing a "new universal." This authentic universal would really be inclusive of all people, regardless of race, class, gender, sexuality, ethnicity, nationality, and whatever else attaches to the "embarrassing etcetera" that, as Judith Butler reminds us, inevitably accompanies such gestures of acknowledging human diversity.

Before signing on to this felicitous agreement about "the necessity of universalism," we may wish to know whether we have anything like a minimal agreement in language, that is, whether we who speak of this universal are even speaking about the same thing. Apart from the not insignificant problem of translating from a philosophical to a political idiom, the whole question of this agreement is virtually occluded by the rush to rescue politics from the virulent particularisms that admit no common ground or sense of collective belonging. Presented in terms of the familiar binary couple, the choice between universalism and particularism seems settled by merely pointing to global and domestic political realities. Universalism is the only alternative to social fragmentation, wild child of the collapse of communism, the rise of deadly nationalisms, and the multiculturalist romance with particularism. To invoke the name of the universal in any affirmative sense is already to sign on to the political diagnosis and its solution. One of the many virtues of Ernesto Laclau's Emancipation(s) is that it offers both an alternative to the binarisms spawned by the "return" to the universal (for example, false universalism/true universalism) and a trenchant critique of the original binary couple itself (universalism/ particularism). Demonstrating the imbrications of the universal and the particular, Laclau shows why it is a matter not of choosing one over the other but of articulating, in a scrupulously political sense, the relation between the two. He thus explicitly rejects the notion that this relation is one of mutual exclusion, and shows that the tendency to see it as just that has led to the impasse of the contemporary debate, an impasse that is glossed over in some highly visible academic cases by proclaiming the necessary return to the universal. Although the language of universalism as spoken by Laclau searches for some common ground between particularists and Universalists, it is more by way of articulating their mutual contamination, that is, how each is rendered impure by the irreducible presence of the other. Aristotle assumes that there is a plurality of individual things (substances) that are characterized by intrinsic properties (forms) each. David Lewis, an analytical philosopher, provided the thesis of Humean supervenience: at the basic level of the world, there are only local qualities in the sense of intrinsic properties instantiated by space-time points or point-sized particles or field sources at space-time points. Space-time points can qualify as individual things in the above mentioned sense. Everything there is in a world like ours supervenes on the distribution of basic intrinsic properties over all space–time points. Whether really everything supervenes on that distribution is not relevant to the present paper. What is important here is the claim that, except for spatio-temporal relations, all the relations between the things at the basic level supervene on their intrinsic properties. We can apply the fundamental principles of mathematical proofs to locate to the True form of nature natural things: Through the systematic observation and analysis (breaking down and classification) of the natural world, in combination with rigorous logic we can make True statements about the natural world and understand: 1) The nature of essences (what something is) 2) The nature of causes (why things occur) Unlike Plato, Aristotle also believes that the other arts are very useful for helping us understand things. Relevant to our course is that he believes argument -- or dialectic -- is a key ingredient for people reaching understandings: by arguing over issues the truth and falsity of the claims becomes increasingly apparent, where Plato might have believed that honing such rhetorical arts only confuses the matter.

Scheme of constitution of the paper is as follows:

SECTION ONE

Observer, Act Of Observation, And What Is Observed, The Conway-Kochen ‘Free Will Theorem’: And Unscientific Determinism Et Al: A Capite Ad Calcem-A Caelo Usque Ad Centrum-Ad Consequentiam Model

SECTION TWO

Conway-Kochen 'Free Will Theorem, Consciousness Et Al: Alis Grave Nil-Amicus Plato, Sed Magis Amica Veritas- Caesar Non Supra Grammaticos- Capax Infiniti Model

SECTION THREE

Relaxed Bell Inequalities, Determinism, And Experimental Free Will, Kochen-Specker Theorems, Theorem Of Mermin Et Al : Abusus Non Tollit Usum-Ad Augusta Per Augusta- Bellum Omnium Contra Omnes Model

SECTION FOUR

Bell's Theorem, Concept Of Space-Time, Deterministic Hidden-Variables Model, Factorizable, Stochastic Model, Et Al: Aurea Mediocritas -Aurora Musis Amica- Aut Inveniam Viam Aut Faciam Model

SECTION FIVE

Local Realism, Bell Inequalities,_Correlation Functions, Information-Theoretic Bell Inequalities, Hidden Quantum Nonlocality, Spin States Et Al: Boni Pastoris Est Tondere Pecus Non Deglubere- Adde Parvum Parvo Magnus Acervus Erit (Ovid) Model

SECTION SIX

Standard Approach To Quantum Systems, Existence Of Hidden States, Markov Context, Hidden Quantum Nonlocality, Et Al.: De Gustibus Non Est Disputandum -Legum Servi Sumus Ut Liberi Esse Possimus

SECTION SEVEN

Superposition Of Four-Dimensional Spacetimes, Geometry Of Space-Time Is A Construct, Quantum Gravity From Quantum Computation Et Al.: A Caelo Usque Ad Centrum-A Bene Placito-Gesamtkunstwerk –Gestalt Model

SECTION EIGHT

Bell States And Maximally Entangled States For N Qubits, Pitowsky Inequalities, Bohm's Example Of Two Entangled Spin 1/2 Quantum Particles, Orthogonal Basis Consisting Of Maximally Entangled States Et Al. : Abundans Cautela Non Nocet -Abyssus Abyssum Invocate- Ad Majorem Dei Gloriam-Ad Lucem Model

SECTION NINE

Algorithm Design And Analysis And Approximation Algorithms, Approximation Methods And Cloud Databases , Polynomials And Resource Management Et Al: Forsan Et Haec Olim Meminisse Iuvabit- Frustra Laborant Quotquot Se Calculationibus Fatigant Pro Inventione Quadraturae Circuli Model

SECTION TEN

Delayed-Choice Entanglement Swapping, Peres's Gedanken Experiment, Greenberger–Horne–Zeilinger Entanglement, Physical Interactions And Processes, Quantum Correlations Et Al. : Altissima Quaeque Flumina Minimo Sono Labi-Zukunftsmusik Leitfaden Model

SECTION ELEVEN

Loophole-Free Bell Inequality, Local Realism, Universal Wave Function, Quantum And Classical Bell-Type Inequalities, Universal Wave Function And Approaches To Quantum Description Of Macroscopic World, With Emphasis On Dissipation, Etal. : Abundans Cautela Non Nocet -Abyssus Abyssum Invocate-Gemutlichkeit-Ansatz Model

SECTION TWELEVE

Spin-Correlation, Quantum Mechanical Probabilities, Mass And Spin-Parity Of The Higgs Boson Candidate, Decay Of The SM Higgs Boson To A Pair Of Z Bosons, Et Al: Age Quod Agis- Agere Sequitur Credere- Agere Sequitur (Esse)- Ad Pondus Omnium (Ad Pond Om)- Ad Referendum Model

SECTION THIRTEEN

Electromagnetic, Weak, And Strong Nuclear Interactions, Particle Masses And Coupling Constants, W Bosons: Ad Pondus Omnium (Ad Pond Om)- Ad Referendum- Ab Ovo Usque Ad Mala

SECTION FOURTEEN

Bekenstein–Hawking Entropy, A Muon (Red Lines On The Right) And Neutrinos, Charged Hadron And A Neutrino, B-Quarks And An LSP, Etal.: Felix Qui Potuit Rerum Cognoscere Causas - Forsan Et Haec Olim Meminisse Iuvabit Model

SECTION FIFTEEN

Proton-Collision Event With A High Di-Muon Mass (1697 Gev), As Seen From The Direction Of The Proton Beams: Age Quod Agis- Agere Sequitur Credere- Agere Sequitur (Esse)- Credula Vitam Spes Fovet Et Melius Cras Fore Semper Dicit Model

SECTION SIXTEEN

Electroweak-Symmetry Breaking Mechanism (Popularly Known As The Brout-Englert-Higgs Mechanism), Boson's Spin-Parity, ZZ And Γγ (Two-Photon), Et Al. : Deus Et Natua Non Faciunt Frusta -Domino Optimo Maximo Model

SECTION ONE

Observer, Act Of Observation, And What Is Observed, The Conway-Kochen 'Free Will Theorem': And Origin Of The Universe Et Al : A Capite Ad Calcem-A Caelo Usque Ad Centrum-Ad Consequentiam Model

INTRODUCTION—VARIABLES USED

(1) Observer, act of observation, and what is observed (two modules)
(2) Attribution of any association in space and **time changes(e&eb)** reality
(3) Cosmic consciousness= Individual consciousness(natures general ledger and individual; general ledger) =0
(4) Space=Time
(5) Subject=Object
(6) Matter= Energy
(7) Being (Existence: by William James:"How comes the world to be here at all instead of the nonentity which might be imagined in its place? ... from nothing to being there is no logical bridge) = Nothingness
(8) Gratification Deprivation = Assets and Liabilities (in a Bank Example-like opening a bank or a branch)
(9) There was darkness hidden inside darkness (Rig-Veda. We interpret this as dark energy was hidden by lack of light)

NOTATION

Module One

Observer, act of observation, and what is observed

G_{13} : Category one of Observer,

G_{14} : Category two of Observer,

G_{15} : Category three of Observer,

T_{13} : Category one of act of observation

T_{14} : Category two of act of observation

T_{15} : Category three of act of observation

Module Two

G_{16} : Category one of act of observation

G_{17} : Category two of act of observation

G_{18} : Category three of act of observation

T_{16} : Category one of observed

T_{17} : Category two of observed

T_{18} : Category three of observed

Module three

G_{20} : Category one of observed

G_{21} : Category two of observed

G_{22} : Category three of observed

T_{20} : Category one of Observer,

T_{21} : Category two of Observer,

T_{22} : Category three of Observer,

Module four

Attribution of any association in space and time changes(e&eb)_reality

G_{24} : Category one of Attribution of any association in space and time however infinitesimal it was

G_{25} : Category two of Attribution of any association in space and time however infinitesimal it was

G_{26} : Category three of Attribution of any association in space and time however infinitesimal it was

T_{24} : Category one of reality of Big Bang

T_{25} : Category two of reality of Big Bang

T_{26} : Category three of reality of Big Bang

Module five

Cosmic consciousness= Individual consciousness(natures general ledger and individual general ledger) =0

G_{28} : Category one of Cosmic consciousness (Nature's general ledger)

G_{29} : Category two of Cosmic consciousness (Nature's general ledger)

G_{30} : Category three of Cosmic consciousness (Nature's general ledger)

T_{28} : Category one of Individual consciousness (individual general ledger)

T_{29} : Category two of Individual consciousness (individual general ledger)

T_{30} : Category three of Individual consciousness (individual general ledger)

Module six

Matter= Energy

G_{32} : Category one of Matter (any feasible and viable scale at say Planck's format could be used)

G_{33} : Category two of Matter (any feasible and viable scale at say Planck's format could be used)

G_{34} : Category three of Matter (any feasible and viable scale at say Planck's format could be used)

T_{32} : Category one of Energy

T_{33} : Category two of Energy

T_{34} : Category three of Energy

Space=Time

Module seven

G_{36} : Category one of Space

G_{37} : Category two of Space

G_{38} : Category three of Space

T_{36} : Category one of Time

T_{37} : Category two of Time

T_{38} : Category three of Time

Module eight

Being (Existence: by William James:"How comes the world to be here at all instead of the nonentity which might be imagined in its place? ... from nothing to being there is no logical bridge) = Nothingness

G_{40} : Category one of Being (Existence: by William James:"How comes the world to be here at all instead of the nonentity which might be imagined in its place? ... from nothing to being there is no logical bridge)

G_{41} : Category two of Being (Existence: by William James:"How comes the world to be here at all instead of the nonentity which might be imagined in its place? ... from nothing to being there is no logical bridge)

G_{42} : Category three of Being (Existence: by William James:"How comes the world to be here at all instead of the nonentity which might be imagined in its place? ... from nothing to being there is no logical bridge

T_{40} : Category one of Nothingness

T_{41} : Category two of Nothingness

T_{42} : Category three of Nothingness

Module Nine

There was darkness hidden inside darkness (Rig-Veda. We interpret this as dark energy was hidden by lack of light or light lessness)

G_{44} : Category one of dark energy and (Form)

G_{45} : Category two of dark energy and (Form)

G_{46} : Category three of dark energy and (Form)

T_{44} : Category one of light lessness and or emptiness

T_{45} : Category two of light lessness and or emptiness

T_{46} : Category three of light lessness and or emptiness

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \qquad 1$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}),t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}),t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}),t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}),t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

59

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{ccc} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) \\ -(b''_{21})^{(3,3,)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) \\ -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{14}$$

60

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) \\ -(b''_{22})^{(3,3,)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) \\ -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{15}$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

61

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) \\ +(a''_{20})^{(3,3,3)}(T_{21},t) & +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) \\ +(a''_{44})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{16}$$

62

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) \\ +(a''_{21})^{(3,3,3)}(T_{21},t) & +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) \\ +(a''_{45})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{17}$$

63

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) \\ +(a''_{22})^{(3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) \\ +(a''_{46})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{18}$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\,-(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\,-(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\,-(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & \boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & \boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & \boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & \boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\,-(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\,-(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\,-(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\,+(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) \quad +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) \quad +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) \quad +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$

Eighth augmentation coefficients $\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \begin{bmatrix} (a'_{42})^{(8)} \boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \begin{bmatrix} (b'_{40})^{(8)} \boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \begin{bmatrix} (b'_{41})^{(8)} \boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \begin{bmatrix} (b'_{42})^{(8)} \boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$

$$= (a_{45})^{(9)}G_{44}$$

$$-\begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$

$$= (a_{46})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\,-(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\,-(b''_{45})^{(9)}(G_{47},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\,-(b''_{46})^{(9)}(G_{47},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad (118)$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \quad (122)$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}),t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \quad (123)$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: (124)

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}((G_{31}),t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: (125)

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: (126)

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \quad (127)$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t) = (p_i)^{(6)}$ 128

$\lim_{G\to\infty}(b_i'')^{(6)}\,((G_{35}),t) = \ (r_i)^{(6)}$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$

$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}((G_{35}),t)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37},t) = (p_i)^{(7)}$

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36{,}37{,}38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36{,}37{,}38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40{,}41{,}42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)}$ 140

$\lim_{G\to\infty}(b_i'')^{(8)}\left((G_{43}),t\right) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t}$ 142

$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\left((G_{43}),t\right)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46$ 146 A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty} (a_i'')^{(9)}(T_{45},t) = (p_i)^{(9)}$

$\lim_{G\to\infty} (b_i'')^{(9)}(G_{47},t) = (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t) - (a_i'')^{(9)}(T_{45},t)| \le (\hat{k}_{44})^{(9)}|T_{45} - T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t) - (b_i'')^{(9)}((G_{47}),t)| < (\hat{k}_{44})^{(9)}||(G_{47}) - (G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^{0} + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^{0} + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^{0} + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^{0} + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^{0} + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^{0} + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^{0} + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^{0} + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^{0} + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^{0} + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^{0} + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^{0}) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^{0}\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^{0}}{G_{17}^{0}}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^{0} + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^{0} + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^{0}) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^{0}\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^{0}}{G_{21}^{0}}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^{0} + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^{0} + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_0^t\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\bar{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left(e^{(\bar{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\bar{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left(e^{(\bar{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\bar{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left(e^{(\bar{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\bar{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_{0}^{t}\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_{+}$.

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a''_{i})^{(1)}$ and $(b''_{i})^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15}. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205
unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)}|G_{21}^{(1)} - G_{21}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}|(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a_{20}'')^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)|\; e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad 214$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}), (\widetilde{T_{27}})\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{24})^{(4)}\left|G_{25}^{(1)} - G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)} +$$

$$\int_0^t\{(a_{24}')^{(4)}\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a_{24}'')^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t} \leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a_{24}')^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)}|G_{29}^{(1)} - G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)})|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ and $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29}\ \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\,(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq$$ 259
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 275

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\ t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations

$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)} v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)} u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}$ $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}$ $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}$ $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le\right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously 331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t} \right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)} t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t} \right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}$,, $(\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}$ $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}$ $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,

and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)},\ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{\left((S_1)^{(7)} - (p_{36})^{(7)}\right)t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\right] + G_{38}^0 e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t}\right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,

and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}$ $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\leq\frac{dv^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

$$\text{it follows } (\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\nu_1)^{(1)}$$

In the same manner , we get 384

$$v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq v^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\leq(\bar{\nu}_1)^{(1)}\leq\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\leq v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\leq v^{(1)}(t)\leq(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\leq u^{(1)}(t)\leq(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)\le\frac{dv^{(9)}}{dt}\le-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)},(\nu_0)^{(9)}$:-

For $0<\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}<(\nu_1)^{(9)}<(\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t)\ge\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}\quad,\quad\boxed{(C)^{(9)}=\frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(9)}=\frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\bar{\nu}_1)^{(9)}$

If $0<(\nu_1)^{(9)}<(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}<(\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)}\le\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}\le\nu^{(9)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\bar{\nu}_1)^{(9)}$$

If $0<(\nu_1)^{(9)}\le(\bar{\nu}_1)^{(9)}\le\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)}\le\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)}\le\nu^{(9)}(t)\le(m_1)^{(9)},\quad\boxed{\nu^{(9)}(t)=\frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)}\le u^{(9)}(t)\le(\mu_1)^{(9)},\quad\boxed{u^{(9)}(t)=\frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(a) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)}G_{25}^{*}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)}G_{25}^{*}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}\quad,\quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}\quad,\quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}\quad,\quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}\quad,\quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof:_Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i \qquad , T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*})=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G^*_{13}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G^*_{14}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G^*_{15}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a_{24}')^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a_{25}')^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a_{26}')^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b_{24}')^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^*\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b_{25}')^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^*\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b_{26}')^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^*\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a_{28}')^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a_{29}')^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a_{30}')^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b_{28}')^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T_{28}^*\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b_{29}')^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T_{29}^*\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b_{30}')^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T_{30}^*\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \ , \ \frac{\partial (b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \ , \frac{\partial (b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \ , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b''_i)^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left((\lambda)^{(1)}\right)^2+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\Big)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\Big]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)})$$

$$\left[\left(((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$+\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})\,((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32})$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)})$$

$$\left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$+\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})\,((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36})$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T^*_{37} + (b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\} = 0$$

$$+$$

$$(\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)})$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWO

Conway-Kochen 'Free Will Theorem, Consciousness Et Al: Alis Grave Nil-Amicus Plato, Sed Magis Amica Veritas- Caesar Non Supra Grammaticos- Capax Infiniti Model

INTRODUCTION—VARIABLES USED

(1) C = h f where C= Consciousness, h= Planck's constant and f= frequency (Equation seen on a video: theory of everything)

(2) A theorem recently propounded by Princeton mathematicians John Conway (who invented the famous Game of Life) and Simon Kochen (one of the originators of the Kochen-Specker paradox of quantum mechanics) supports a powerful challenge to the scientific credentials of determinism, by showing that two cornerstones of contemporary science, namely acceptance of the scientific method as a reliable way of finding out about the world, and relativity theory's exclusion of faster-than-light transmission of information, **together (e)conflict with** determinism, in both its versions. Belief in determinism may thus come to be seen as notably unscientific. This conclusion has previously had some support from a theorem devised by John Bell and experiments undertaken by Alain Aspect, but Conway-Kochen theorem supports it more strongly. **(THE CONWAY-KOCHEN 'FREE WILL THEOREM': AND UNSCIENTIFIC DETERMINISM: David Hodgson**)

The proof of the theorem relies on three axioms, which Conway and Kochen call "fin", "spin", and "twin". The spin and twin axioms can be verified experimentally.

(3) Fin: There is a maximum speed for propagation of information (not necessarily the speed of light). This assumption rests upon causality. Maximum speed for propagation of information **(=)** finite

(4) Spin: The squared spin component of certain elementary particles of spin one, taken in three orthogonal directions, **will (=) be** a permutation of (1, 1, and 0).

(5) Twin: It is possible to "entangle" two elementary particles, and separate them by a significant distance, **so that (=) they** have the same squared spin results if measured in parallel directions a consequence of (but more limited than) quantum entanglement.

(6) In their later paper, "The Strong Free Will Theorem," Conway and Kochen weaken the Fin axiom (thereby strengthening the theorem) to a new axiom called Min, which asserts only that two experimenters separated in a space-like way **can (eb)make choices** of measurements independently of each other. In particular, they are not asserting that all information must travel finitely fast; only the particular information about choices of measurements.

The theorem

(7) The theorem states that, given the axioms, if the two experimenters in question are free to make choices about what measurements to take, **then the (eb) results** of the measurements cannot be determined by anything previous to the experiments.

(8) Since the theorem applies to any arbitrary physical theory consistent with the axioms, it would not even be possible to place the information into the universe's past in an ad hoc way. The argument proceeds from the Kochen-Specker theorem, **which (eb) shows that the** result of any individual measurement of spin was not fixed independently of the choice of measurements.

Limitations

Conway and Kochen do not prove that free will does exist. The definition of "free will" used in the proof of this theorem **is simply that** an outcome is "not determined" by prior conditions, and some philosophers strongly dispute the equivalence of "not determined" with free will. Some critics argue that the theorem only applies to deterministic models. Others have argued that the indeterminism that Conway and Kochen claim to have established was already assumed in the premises of their proof.

NOTATION

Module One

C = h f where C= Consciousness, h= Planck's constant and f= frequency (Equation seen on a video: theory of everything)

G_{13} : Category one of r.h.s of the equation in the foregoing

G_{14} : category two of rhs of the equation in the foregoing

G_{15} : category three of rhs of the equation in the foregoing

T_{13} : Category one of LHS OF THE EQUATION IN THE FOREGOING

T_{14} : Category two of LHS OF THE EQUATION IN THE FOREGOING

T_{15} : Category three of LHS OF THE EQUATION IN THE FOREGOING

Module Two

A theorem recently propounded by Princeton mathematicians John Conway (who invented the famous Game of Life) and Simon Kochen (one of the originators of the Kochen-Specker paradox of quantum mechanics) supports a powerful challenge to the scientific credentials of determinism, by showing that two cornerstones of contemporary science, namely acceptance of the scientific method as a reliable way of finding out about the world, and relativity theory's exclusion of faster-than-light transmission of information, **together (e)conflict with** determinism, in both its versions. Belief in determinism may thus come to be seen as notably unscientific. This conclusion has previously had some support from a theorem devised by John Bell and experiments undertaken by Alain Aspect, but Conway-Kochen theorem supports it more strongly. **(the conway-kochen 'free will theorem': and unscientific determinism: David Hodgson**)

G_{16} : Category one of determinism, in both its versions

G_{17} : Category two of determinism, in both its versions

G_{18} : Category three of determinism, in both its versions

T_{16} : Category one of acceptance of the scientific method as a reliable way of finding out about the world, and relativity theory's exclusion of faster-than-light transmission of information

T_{17} : Category two of acceptance of the scientific method as a reliable way of finding out about the world, and relativity theory's exclusion of faster-than-light transmission of information

T_{18} : Category three of acceptance of the scientific method as a reliable way of finding out about the world, and relativity theory's exclusion of faster-than-light transmission of information

Fin: There is a maximum speed for propagation of information (not necessarily the speed of light). This assumption rests upon causality. Maximum speed for propagation of information finite

Module three

G_{20} : Category one of finiteness

G_{21} : Category two of finiteness

G_{22} : Category three of finiteness

T_{20} : Category one of Maximum speed for propagation of information

T_{21} : Category two of Maximum speed for propagation of information

T_{22} : Category three of Maximum speed for propagation of information

Module four

Spin: The squared spin component of certain elementary particles of spin one, taken in three orthogonal directions, **will be** a permutation of (1, 1, and 0).

G_{24} : Category one of permutation of (1, 1, and 0).

G_{25} : Category two of permutation of (1, 1, and 0).

G_{26} : Category three of permutation of (1, 1, and 0).

T_{24} : Category one of squared spin component of certain elementary particles of spin one, taken in three orthogonal directions,

T_{25} : Category two of squared spin component of certain elementary particles of spin one, taken in three orthogonal directions,

T_{26} : Category three of squared spin component of certain elementary particles of spin one, taken in three orthogonal directions,

Module five

Twin: It is possible to "entangle" two elementary particles, and separate them by a significant distance, **so that they** have the same squared spin results if measured in parallel directions a consequence of (but more limited than) quantum entanglement

G_{28} : Category one of corresponding squared spin results if measured in parallel directions a consequence of (but more limited than) quantum entanglement

G_{29} : Category two of corresponding squared spin results if measured in parallel directions a consequence of (but more limited than) quantum entanglement

G_{30} : Category three of corresponding squared spin results if measured in parallel directions a consequence of (but more limited than) quantum entanglement

T_{28} : Category one of possibilities to "entangle" two elementary particles, and separate them by a significant distance

T_{29} : Category two of possibilities to "entangle" two elementary particles, and separate them by a significant distance

T_{30} : Category three of possibilities to "entangle" two elementary particles, and separate them by a significant distance

Module six

In their later paper, "The Strong Free Will Theorem," Conway and Kochen weaken the Fin axiom (thereby strengthening the theorem) to a new axiom called Min, which asserts only that two experimenters separated in a space-like way **can (eb)make choices** of each other. In particular, they are not asserting that all information must travel measurements independently finitely fast; only the particular information about choices of measurements

G_{32} : Category one of two experimenters separated in a space-like way

G_{33} : Category two of two experimenters separated in a space-like way

G_{34} : Category three of two experimenters separated in a space-like way

T_{32} : Category one of each other. In particular, they are not asserting that all information must travel measurements independently finitely fast; only the particular information about choices of measurements

T_{33} : Category two of each other. In particular, they are not asserting that all information must travel measurements independently finitely fast; only the particular information about choices of measurements

T_{34} : Category three of each other. In particular, they are not asserting that all information must travel measurements independently finitely fast; only the particular information about choices of measurements

Module seven

The theorem states that, given the axioms, if the two experimenters in question are free to make choices about what measurements to take, **then the (eb) results** of the measurements cannot be determined by anything previous to the experiments

G_{36} : Category one of, given the axioms and if the two experimenters in question are free to make choices about what measurements to be made

G_{37} : Category two of , given the axioms, if the two experimenters in question are free to make choices about what measurements to be made

G_{38} : Category three of , given the axioms, if the two experimenters in question are free to make choices about what measurements to be made

T_{36} : Category one of **results** of the measurements cannot be determined by anything previous to the experiments

T_{37} : Category two of **results** of the measurements cannot be determined by anything previous to the experiments

T_{38} : Category three of **results** of the measurements cannot be determined by anything previous to the experiments

Module eight

Kochen-Specker theorem (Since the theorem applies to any arbitrary physical theory consistent with the axioms, it would not even be possible to place the information into the universe's past in an ad hoc way) **which (eb) shows that the** result of any individual measurement of spin was not fixed independently of the choice of measurements

G_{40} : Category one of Kochen-Specker theorem (Since the theorem applies to any arbitrary physical theory consistent with the axioms, it would not even be possible to place the information into the universe's past in an ad hoc way)

G_{41} : Category two of Kochen-Specker theorem (Since the theorem applies to any arbitrary physical theory consistent with the axioms, it would not even be possible to place the information into the universe's past in an ad hoc way)

G_{42} : Category three of Kochen-Specker theorem (Since the theorem applies to any arbitrary physical theory consistent with the axioms, it would not even be possible to place the information into the universe's past in an ad hoc way)

T_{40} : Category one of result of any individual measurement of spin was not fixed independently of the choice of measurements

T_{41} : Category two of result of any individual measurement of spin was not fixed independently of the choice of measurements

T_{42} : Category three of result of any individual measurement of spin was not fixed independently of the choice of measurements

Module Nine

G_{44} : Category one of outcome is "not determined" by prior conditions and some philosophers strongly dispute the equivalence of "not determined" with free will. (Some critics argue that the theorem only applies to deterministic models. Others have argued that the indeterminism that Conway and Kochen claim to have established was already assumed in the premises of their proof).

G_{45} : Category two of outcome is "not determined" by prior conditions, and some philosophers strongly dispute the equivalence of "not determined" with free will. (Some critics argue that the theorem only applies to deterministic models. Others have argued that the indeterminism that Conway and Kochen claim to have established was already assumed in the premises of their proof).

G_{46} : Category three of outcome is "not determined" by prior conditions, and some philosophers strongly dispute the equivalence of "not determined" with free will. (Some critics argue that the theorem only applies to deterministic models. Others have argued that the indeterminism that Conway and Kochen claim to have established was already assumed in the premises of their proof).

T_{44} : Category one of definition of "free will" used in the proof of the theorem

T_{45} : Category two of definition of "free will" used in the proof of the theorem

T_{46} : Category three of definition of "free will" used in the proof of the theorem

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$: $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$

$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad (1)$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad (2)$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad (3)$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad (4)$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad (5)$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad (7)$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad (8)$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad (9)$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad (10)$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad (11)$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad (13)$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad (14)$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad (15)$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad (16)$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad (17)$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}),t)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}),t)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3
$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3
$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3
$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}\ \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,)}(G,t)}\ \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,)}(G,t)}\ \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,)}(G,t)}\ \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \tag{69}$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b_{20}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b_{36}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{37}'')^{(7,7,7,7,7)}(G_{39},t)$, $-(b_{38}'')^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b_{40}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b_{46}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a_{28}')^{(5)} & +(a_{28}'')^{(5)}(T_{29},t) & +(a_{24}'')^{(4,4,)}(T_{25},t) & +(a_{32}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{16}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{44}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a_{29}')^{(5)} & +(a_{29}'')^{(5)}(T_{29},t) & +(a_{25}'')^{(4,4,)}(T_{25},t) & +(a_{33}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{14}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{17}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{45}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a_{30}')^{(5)} & +(a_{30}'')^{(5)}(T_{29},t) & +(a_{26}'')^{(4,4,)}(T_{25},t) & +(a_{34}'')^{(6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1)}(T_{14},t) & +(a_{18}'')^{(2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3)}(T_{21},t) \\ & +(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t) & +(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t) & +(a_{46}'')^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a_{28}'')^{(5)}(T_{29},t)$, $+(a_{29}'')^{(5)}(T_{29},t)$, $+(a_{30}'')^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $+(a_{24}'')^{(4,4,)}(T_{25},t)$, $+(a_{25}'')^{(4,4,)}(T_{25},t)$, $+(a_{26}'')^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3
$+(a_{32}'')^{(6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3
$+(a_{13}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{14}'')^{(1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3
$+(a_{16}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a_{20}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a_{36}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{37}'')^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a_{38}'')^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a_{40}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a_{42}'')^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a_{46}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b_{28}')^{(5)} & -(b_{28}'')^{(5)}(G_{31},t) & -(b_{24}'')^{(4,4,)}(G_{27},t) & -(b_{32}'')^{(6,6,6)}(G_{35},t) \\ & -(b_{13}'')^{(1,1,1,1,1)}(G,t) & -(b_{16}'')^{(2,2,2,2,2)}(G_{19},t) & -(b_{20}'')^{(3,3,3,3,3)}(G_{23},t) \\ & -(b_{36}'')^{(7,7,7,7,7,7)}(G_{39},t) & -(b_{40}'')^{(8,8,8,8,8,8)}(G_{43},t) & -(b_{44}'')^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$
$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$
$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$
$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[\begin{matrix}(a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[\begin{matrix}(a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{14} \quad (92)$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[\begin{matrix}(a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{15} \quad (93)$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{37}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right]T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$ *96*
$$= (a_{44})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \tag{97}$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \tag{98}$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: (99)

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: (100)

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\widehat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\widehat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\widehat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants

$(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}\ (T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\ \big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - \ T_{41}'| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\widehat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ , \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (a_i)^{(8)} + (a_i')^{(8)} + \ (\hat{A}_{40})^{(8)} + \ (\hat{P}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ \ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\widehat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\widehat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\overline{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\overline{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\overline{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(1)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{16})^{(2)}\left|G_{17}^{(1)}-G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)\left|G_{16}^{(1)}-G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)}-(G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\ (G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\widehat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{20})^{(3)}\right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})} \right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)} - G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\tilde{M}_{24})^{(4)}s_{(24)}}e^{(\tilde{M}_{24})^{(4)}s_{(24)}}ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 228

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\leq\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25}\leq\left((\widehat{M}_{24})^{(4)}\right)_2=G_{25}^0+2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26}\leq\left((\widehat{M}_{24})^{(4)}\right)_3=G_{26}^0+2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t),t))=(b'_{25})^{(4)}$ then $T_{25}\ \to\infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a_{28}')^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{-(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a_{28}'')^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$G_{28}^{(2)}|(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a_{28}'')^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a_{28}')^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a_{28}'')^{(5)}$ and $(b_{28}'')^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\geq 0$

$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(5)}t\right)}>0\ \ \text{for t}>0$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\leq\left((\widehat{M}_{28})^{(5)}\right)_1-(a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29}\leq\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30}\leq\left((\widehat{M}_{28})^{(5)}\right)_3=G_{30}^0+2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t))=(b_{29}')^{(5)}$ then $T_{29}\ \to\infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ >t_5$

$(b_{29})^{(5)}-(b_i'')^{(5)}((G_{31})(t),t)<\varepsilon_5, T_{28}\ (t)>(m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{-(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)}\left|(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a_{32}'')^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t), t(t), t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\tilde{M}_{36})^{(7)}} < 1$ and to choose $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{-(\tilde{M}_{36})^{(7)}s_{(36)}} +$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}t} \leq \quad 259$$
$$\frac{1}{(\tilde{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered 260

as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is

unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right)=(b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)}+\left((\widehat{P}_{40})^{(8)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{40})^{(8)}\right]\leq(\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ \ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}\,((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t\ > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279

A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and

respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\ (t) = 0\ and\ T_i\ (t) = 0$

From 99 to 44 it results

$$G_i\ (t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\ (t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\ (t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\ \frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)},\ \boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)},$ 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\Big(\frac{(a_{22})^{(3)}G^0_{20}}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\Big[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\Big] + G^0_{22}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \quad 324$$

$$\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}\right]+G_{22}^{0}e^{-(a'_{22})^{(3)}t})$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$$

$$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$$

$$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)},(v_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},v^{(4)},u^{(4)}$: 329

By $(v_1)^{(4)}>0$, $(v_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_1)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)}>0$, $(\bar{v}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$$(m_2)^{(4)}=(v_0)^{(4)},(m_1)^{(4)}=(v_1)^{(4)},\ \boldsymbol{if}\ (v_0)^{(4)}<(v_1)^{(4)}$$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}$ $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}$ $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}$ $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}$ $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}$ $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \leq G_{26}(t) \leq \right.$$

334

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \leq T_{26}(t) \leq$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,
and $\boxed{(v_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^{0}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G_{28}^{0}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t}-e^{-(S_2)^{(5)}t}\right]+G_{30}^{0}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left.\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+G_{30}^{0}e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$ and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$ and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)},(v_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},v^{(7)},u^{(7)}$: 361

By $(v_1)^{(7)}>0,(v_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0,(u_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2+(\sigma_1)^{(7)}v^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{v}_1)^{(7)},,(\bar{v}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)}>0,(\bar{v}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0,(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2+(\sigma_2)^{(7)}v^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(v_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(v_0)^{(7)},(m_1)^{(7)}=(v_1)^{(7)},\ \boldsymbol{if}\ (v_0)^{(7)}<(v_1)^{(7)}$

$(m_2)^{(7)}=(v_1)^{(7)},(m_1)^{(7)}=(\bar{v}_1)^{(7)},\boldsymbol{if}\ (v_1)^{(7)}<(v_0)^{(7)}<(\bar{v}_1)^{(7)},$
and $\boxed{(v_0)^{(7)}=\frac{G_{36}^{0}}{G_{37}^{0}}}$

$(m_2)^{(7)}=(v_1)^{(7)},(m_1)^{(7)}=(v_0)^{(7)},\ \boldsymbol{if}\ (\bar{v}_1)^{(7)}<(v_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, , (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t}\le G_{42}(t)\le\right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}\right]+G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\le T_{42}(t)\le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)}\le-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\le-(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)}\le-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\le-(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}, \ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$$

and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \le \frac{d\nu^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})v^{(2)} - (a_{17})^{(2)}v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)}v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}_{16}^0}{\mathrm{G}_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\leq\frac{dv^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\geq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\leq v^{(3)}(t)\leq(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\leq v^{(3)}(t)\leq(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\leq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\leq v^{(3)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\leq(\bar{v}_1)^{(3)}\leq(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\leq v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\leq v^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$
In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows
$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also**

defines $(\nu_0)^{(8)}$ for the special case .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $v^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\ ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\ , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(b) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows

that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$$T^*_{24} = \frac{(b_{24})^{(4)}T^*_{25}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T^*_{26} = \frac{(b_{26})^{(4)}T^*_{25}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a_{13}')^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a_{14}')^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a_{15}')^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b_{13}')^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^*\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b_{14}')^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^*\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b_{15}')^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^*\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}(\,(G_{19})^*\,) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(16)(j)}T_{16}^*\mathbb{G}_j\big) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}} (T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j} ((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)} G_{40}^* \mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)} G_{41}^* \mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)} G_{42}^* \mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)} T_{40}^* \mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)} T_{41}^* \mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)} T_{42}^* \mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is

$$(\lambda)^{(1)}+(b'_{15})^{(1)}-(r_{15})^{(1)})\{((\lambda)^{(1)}+(a'_{15})^{(1)}+(p_{15})^{(1)})$$

$$\left[\left(((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)})(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\right]$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(14)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\right)$$

$$+\left(((\lambda)^{(1)}+(a'_{14})^{(1)}+(p_{14})^{(1)})(q_{13})^{(1)}G^{*}_{13}+(a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(13)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$+\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)})\{((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$+\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)})\{((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)})$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left(\left((\lambda)^{(9)}\right)^2 + \left((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2 + \left((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45} + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45} + (b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\} = 0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION THREE

Relaxed Bell Inequalities, Determinism, And Experimental Free Will, Kochen-Specker Theorems, Theorem Of Mermin Et Al : Abusus Non Tollit Usum-Ad Augusta Per Augusta- Bellum Omnium Contra Omnes Model

INTRODUCTION—VARIABLES USED

Relaxed Bell inequalities and Kochen-Specker theorems: Michael J.W. Hall

(1) The combination of various physically plausible properties, such as no signaling, determinism, and experimental free will, is known **to be (ac = dc =0: STUDY ALL THE THREE CASES) incompatible** with quantum correlations.

(2) No signaling, determinism, and experimental free will, properties must be individually or jointly **relaxed (e)** in any model of such correlations.

(3) The necessary degrees of relaxation are quantified here**, (e&eb) via** natural distance and information-theoretic measures. Such quantification of necessary degrees of relaxation through (e&eb) natural distance and information theoretic measures **allows (eb)** quantitative comparisons between different models in terms of the resources, such as the number of bits, of randomness, communication, and/or correlation that they require. For example, measurement dependence is a relatively strong resource for modeling singlet state correlations, with only 1/15 of one bit of correlation **required (e)** between measurement settings and the underlying variable.

(4) It is shown how various 'relaxed' Bell inequalities may be obtained, **which (eb) precisely specify** the complementary degrees of relaxation required to model any given violation of a standard Bell inequality.

(5) The robustness of a class of Kochen-Specker theorems, **to** relaxation of measurement independence, is also investigated.

(6) It is shown that a theorem of Mermin remains **valid (eb)** unless measurement independence is relaxed by 1/3.

(7) The Conway-Kochen 'free will' theorem and a result of Hardy are less robust, **failing (e)** if measurement independence is relaxed by only 6.5% and 4.5%, respectively.

(8) An appendix shows the existence of an outcome independent model thereof is **equivalent to** the existence of a deterministic model. Journal reference: Phys. Rev. A 84, 022102 (2011) DOI: 10.1103/PhysRevA.84.022102 Cite as: arXiv:1102.4467 [quant-ph] (or

arXiv:1102.4467v3 [quant-ph] for this version)

NOTATION

Module One

G_{13} : Category one of combination of various physically plausible properties, such as no signaling, determinism, and experimental free will

G_{14} : Category two of combination of various physically plausible properties, such as no signaling, determinism, and experimental free will

G_{15} : Category three of combination of various physically plausible properties, such as no signaling, determinism, and experimental free will

T_{13} : Category one of quantum correlations

T_{14} : Category two of quantum correlations

T_{15} : Category three of quantum correlations

Module Two

G_{16} : Category one of models of such correlations for feasible results

G_{17} : Category two of models of such correlations for feasible results

G_{18} : Category three of models of such correlations for feasible results

T_{16} : Category one of Relaxation of the properties such as no signaling, determinism, and experimental free will

T_{17} : Category two of Relaxation of the properties such as no signaling, determinism, and experimental free will

T_{18} : Category three of Relaxation of the properties such as no signaling, determinism, and experimental free will

Module three

G_{20} : Category one of necessary degrees of relaxation are quantified **(e&eb) via** natural distance and information-theoretic measures which allows for necessary degrees of relaxation through (e&eb) natural distance and information theoretic measures

G_{21} : Category two of necessary degrees of relaxation are quantified **(e&eb) via** natural distance and information-theoretic measures which allows for necessary degrees of relaxation through (e&eb) natural distance and information theoretic measures

G_{22} : Category three of necessary degrees of relaxation are quantified **(e&eb) via** natural distance and information-theoretic measures which allows for necessary degrees of relaxation through (e&eb) natural distance and information theoretic measures

T_{20} : Category one of quantitative comparisons between different models in terms of the resources, such as the number of bits, of randomness, communication, and/or correlation that they require. For example, measurement dependence is a relatively strong resource for modeling singlet state correlations, with only 1/15 of one bit of correlation **required (e)** between measurement settings and the underlying variable.

T_{21} : Category two of quantitative comparisons between different models in terms of the resources, such as the number of bits, of randomness, communication, and/or correlation that they require. For example, measurement dependence is a relatively strong resource for modeling singlet state correlations,

with only 1/15 of one bit of correlation **required (e)** between measurement settings and the underlying variable.

T_{22} : Category three of quantitative comparisons between different models in terms of the resources, such as the number of bits, of randomness, communication, and/or correlation that they require. For example, measurement dependence is a relatively strong resource for modeling singlet state correlations, with only 1/15 of one bit of correlation **required (e)** between measurement settings and the underlying variable.

Module four

G_{24} : Category one of obtention of 'relaxed' Bell inequalities

G_{25} : Category two of obtention of 'relaxed' Bell inequalities

G_{26} : Category three of obtention of 'relaxed' Bell inequalities

T_{24} : Category one of **precise specification of** the complementary degrees of relaxation required to model any given violation of a standard Bell inequality.

T_{25} : Category two of **precise specification of** the complementary degrees of relaxation required to model any given violation of a standard Bell inequality.

T_{26} : Category three of **precise specification of** the complementary degrees of relaxation required to model any given violation of a standard Bell inequality.

Module five

G_{28} : Category one of robustness of a class of Kochen-Specker theorems

G_{29} : Category two of robustness of a class of Kochen-Specker theorems

G_{30} : Category three of robustness of a class of Kochen-Specker theorems

T_{28} : Category one of relaxation of measurement independence

T_{29} : Category two of relaxation of measurement independence

T_{30} : Category three of relaxation of measurement independence

Module six

G_{32} : Category one of theorem of Mermin (Systems that hold for the theorem and the concomitant characterstics)

G_{33} : Category two of theorem of Mermin (Systems that hold for the theorem and the concomitant characterstics. Please see the explanation in the introduction)

G_{34} : Category three of Mermin (Systems that hold for the theorem and the concomitant characterstics. Please see the explanation in the introduction)

T_{32} : Category one of measurement independence relaxation by 1/3.

T_{33} : Category two of measurement independence relaxation by 1/3.

T_{34} : Category three of measurement independence relaxation by 1/3.

Module seven

G_{36} : Category one of measurement independence relaxation by only 6.5% and 4.5%, respectively

G_{37} : Category two of measurement independence relaxation by only 6.5% and 4.5%, respectively

G_{38} : Category three of measurement independence relaxation by only 6.5% and 4.5%, respectively

T_{36} : Category one of Conway-Kochen 'free will' theorem and a result of Hardy

T_{37} : Category two of Conway-Kochen 'free will' theorem and a result of Hardy

T_{38} : Category three of Conway-Kochen 'free will' theorem and a result of Hardy

Module eight

G_{40} : Category one of existence of a deterministic model

G_{41} : Category two of existence of a deterministic model

G_{42} : Category three of existence of a deterministic model

T_{40} : Category one of existence of an outcome independent model thereof

T_{41} : Category two of existence of an outcome independent model thereof

T_{42} : Category three of existence of an outcome independent model thereof

Module Nine

Physics Research International Volume 2012 (2012), Article ID 352543, 5 pages doi:10.1155/2012/352543 Research Article Reality or Locality? Proposed Test to Decide How Nature Breaks Bell's Inequality Johan Hansson Department of Physics, Luleå University of Technology, 971 87 Luleå, Sweden Academic Editors: Ali Hussain Reshak. Copyright © 2012 Johan Hansson. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bell's theorem, and its experimental tests, has shown that the two premises for Bell's inequality—locality and objective reality—**cannot (e) both hold in nature**, as Bell's inequality is broken. A simple test is proposed, which for the first time may decide which alternative nature actually prefers on the fundamental, quantum level. If each microscopic event is truly random (e.g., as assumed in orthodox quantum mechanics) objective reality is not valid whereas if each event is described by an unknown but deterministic mechanism (“hidden variables”) locality is not valid. This may be analyzed and decided by the well-known reconstruction method of Ruelle and Takens; in the former case no structure should be discerned, in the latter a reconstructed structure should be visible. This could in principle be tested by comparing individual “hits” in a double-slit experiment, but in practice a single fluorescent atom, and its (seemingly random) temporal switching between active/inactive states would possibly be better/more practical, easier to set up, observe, and analyze. However, only imagination limits the list of possible experimental setups.

G_{44} : Category one of actualization of the principle of locality (again classification is based on systemic characterstics for which the locality holds. For further detailed explanation please see the introduction)

G_{45} : Category two of actualization of the principle of locality (again classification is based on systemic

characterstics for which the locality holds. For further detailed explanation please see the introduction)

G_{46} : Category three of actualization of the principle of locality (again classification is based on systemic characterstics for which the locality holds. For further detailed explanation please see the introduction)

T_{44} : Category one of spatio temporal realisation of the principle of objective reality (again classification is based on systemic characterstics for which the locality holds. For further detailed explanation please see the introduction)

T_{45} : Category two of spatio temporal realisation of the principle of objective reality (again classification is based on systemic characterstics for which the locality holds. For further detailed explanation please see the introduction)

T_{46} : Category three of spatio temporal realisation of the principle of objective reality (again classification is based on systemic characterstics for which the locality holds. For further detailed explanation please see the introduction)

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients
Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27}),t\big)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\big((G_{27}),t\big) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}), t\right)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}), t\right)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}), t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3
$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3
$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3
$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3
$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3
$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3
$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3
$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients*

for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients

for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \qquad 82$$

$$\frac{dT_{29}}{dt}=(b_{29})^{(5)}T_{28}-\left[\begin{array}{ccc}(b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{29} \qquad 83$$

$$\frac{dT_{30}}{dt}=(b_{30})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{30} \qquad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
$are\ first\ detrition\ coefficients\ \ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ \ third\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} & -(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} & -(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} & -(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second

detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \left[\begin{matrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)}\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{matrix}\right]G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{matrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)}\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{matrix}\right]G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{matrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)}\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{matrix}\right]G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)} & \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\begin{bmatrix} (b'_{45})^{(9)} & \boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad (97)$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$lim_{T_2\to\infty}(a_i'')^{(1)}\,(T_{14},t)=(p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}\,(G,t)=\ \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)},(p_i)^{(1)},\ \ (r_i)^{(1)}}$ are positive constants and $\boxed{i=13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},$ are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\ ,\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)}+(a_i')^{(1)}+\ (\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)}+(b_i')^{(1)}+\ (\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le\left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le\ \ (r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}\,(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=\ \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26$$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$ 118

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$

and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}<1$$

Definition of $(\hat{P}_{28})^{(5)},(\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)},(\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)},(a_i')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(p_i)^{(5)},\ (r_i)^{(5)},i=28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)}+(a_i')^{(5)}+\ (\hat{A}_{28})^{(5)}+\ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}]<1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)}+(b_i')^{(5)}+\ (\hat{B}_{28})^{(5)}+\ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}]<1$$

Where we suppose

$$(a_i)^{(6)},(a_i')^{(6)},(a_i'')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(b_i'')^{(6)}>0,\quad i,j=32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)},(b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t)\le(p_i)^{(6)}\le(\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t)\le\ (r_i)^{(6)}\le(b_i')^{(6)}\le(\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}),t\right)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\left((G_{35}),t\right)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\widehat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)}$$ 132

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It

is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+(\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)}+(b_i')^{(8)}+(\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,,T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{24})^{(4)}$, $T_i^0 \leq (\hat{Q}_{24})^{(4)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\overline{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that (172)

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that (174)

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)}+G_{29}^{0}\right)e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{P}_{32})^{(6)}+G_{33}^{0}\right)e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{P}_{36})^{(7)}+G_{37}^{0}\right)e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{P}_{40})^{(8)}+G_{41}^{0}\right)e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$G_{13}^{(2)}|(a_{13}'')^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a_{13}'')^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a_{13}')^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a_{13}'')^{(1)}$ and $(b_{13}'')^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{(-(b_i')^{(1)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a_{14}')^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\ (G(t),t))=(b_{14}')^{(1)}$ then $T_{14}\ \to\infty$. (189)

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1, T_{13}\ (t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$

$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205

unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b''_{18})^{(2)}\left((G_{19})(t),t\right)=(b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}<1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 215

$(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself (224)

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric (225)

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\leq \tag{226}$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2 \; and \; \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25} \; and \; G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25} \; or \; G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24} \; is$ bounded, from below, the same property holds for $G_{25} \; and \; G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$ 239

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad (245)$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 250

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 261

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{40})^{(8)}|G_{41}^{(1)}-G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}|(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})-(a''_{40})^{(8)}(T_{41}^{(2)},s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$|(G_{43})^{(1)}-(G_{43})^{(2)}|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)})d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(8)}t)}>0 \quad \text{for t} > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1,((\widehat{M}_{40})^{(8)})_2$ *and* $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40}<(\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt}\le((\widehat{M}_{40})^{(8)})_1-(a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})} \right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{44})^{(9)} \left|G_{45}^{(1)} - G_{45}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\, ds_{(44)} +$$
$$\int_0^t \{(a'_{44})^{(9)} \left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$G_{44}^{(2)} \left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\} ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)}\right) d\left(((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ *and* $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)} \left((\widehat{M}_{44})^{(9)}\right)_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, $\boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, $\boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16} e^{((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)})\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16} e^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$

$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

332

$$G^0_{24} e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24} e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)}-(a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)},$ $\boldsymbol{if}$ $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}$ $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)},$ $\boldsymbol{if}$ $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)},$ $\boldsymbol{if}$ $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)},$ $\boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},, (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^{0}}{G_{33}^{0}}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0$, $(v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0$, $(\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G^0_{36} e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t}\leq G_{38}(t)\leq\right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\leq T_{38}(t)\leq \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$$

$$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$$

$$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)}\leq-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\leq-(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)}\leq-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\leq-(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)},(\nu_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},\nu^{(8)},u^{(8)}$: 372

By $(\nu_1)^{(8)}>0$, $(\nu_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0$, $(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2+(\sigma_1)^{(8)}\nu^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{\nu}_1)^{(8)},,(\bar{\nu}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)}>0$, $(\bar{\nu}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0$, $(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2+(\sigma_2)^{(8)}\nu^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, $\boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}\left((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)}\right)}\left[e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}\left((S_1)^{(8)} - (a'_{42})^{(8)}\right)}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}\left((R_1)^{(8)} - (b'_{42})^{(8)}\right)}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}\left((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)}\right)}\left[e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, $\boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, $\boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also

defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $v^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq v^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)\le\frac{dv^{(4)}}{dt}\le-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_4)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)},(v_0)^{(4)}$:-

For $0<\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}<(v_1)^{(4)}<(\bar{v}_1)^{(4)}$

$$v^{(4)}(t)\ge\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}\quad,\quad\boxed{(C)^{(4)}=\frac{(v_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(v_2)^{(4)}}}$$

it follows $(v_0)^{(4)}\le v^{(4)}(t)\le(v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(4)}=\frac{(\bar{v}_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)}\le v^{(4)}(t)\le(\bar{v}_1)^{(4)}$

If $0<(v_1)^{(4)}<(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}<(\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)}\le\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}\le v^{(4)}(t)\le$$

$$\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(\bar{v}_1)^{(4)}$$

If $0<(v_1)^{(4)}\le(\bar{v}_1)^{(4)}\le\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)}\le v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)}\le v^{(4)}(t)\le(m_1)^{(4)},\quad\boxed{v^{(4)}(t)=\frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)}\le u^{(4)}(t)\le(\mu_1)^{(4)},\quad\boxed{u^{(4)}(t)=\frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \le \frac{dv^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$v^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le v^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le v^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le v^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le v^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \le v^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad (477)$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad (478)$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad (479)$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad (480)$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad (481)$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad (482)$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad (483)$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad (484)$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad (484A)$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: (485)

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: (486)

(c) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: (487)

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502
$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{\left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})\right]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{\left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})\right]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{\left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)\right]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{\left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)\right]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \tag{513}$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b''_i)^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \ , \ \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}(\,(G_{35})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\right)\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left(\,(a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left(\,(b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^{2}+\left(\,(a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left(\,(a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^{2}+\left(\,(b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left((\lambda)^{(9)}\right)^{2}+\left(\,(a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FOUR

Bell's Theorem, Concept Of Space-Time, Deterministic Hidden-Variables Model, Factorizable, Stochastic Model, Et Al: Aurea Mediocritas -Aurora Musis Amica- Aut Inveniam Viam Aut Faciam Model

INTRODUCTION—VARIABLES USED

Bell's theorem. Experimental tests and implications J F Clauser and A Shimony

(1) Bell's theorem represents a significant advance in understanding the conceptual foundations of quantum mechanics. The theorem shows that essentially all local theories of natural phenomena that are formulated within the framework of realism may be tested using a single experimental arrangement. Moreover, the predictions by those theories must significantly differ from those by quantum mechanics. Experimental results evidently refute the theorem's predictions for these theories and favour those of quantum mechanics. The **conclusions (eb) are philosophically startling**:

Either one must totally abandon the realistic philosophy of most working scientists or dramatically revision of concept of space-time. Rep. Prog. Phys. 41 1881 doi:10.1088/0034-4885/41/12/002

Hidden Variables, Joint Probability, and the Bell Inequalities: Arthur Fine © 1982 The American Physical Society URL: http://link.aps.org/doi/10.1103/PhysRevLett.48.291 DOI: 10.1103/PhysRevLett.48.291 PACS: 03.50.Bz, 02.50. +s

It is shown that the following statements about a quantum correlation experiment are mutually equivalent.

(1) There is a deterministic hidden-variables model for the experiment. (2) There is a factorizable, stochastic model. (3) There is one joint distribution for all observables of the experiment, returning the experimental probabilities. (4) There are well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables. (5) The Bell inequalities hold.

NOTATION

Module One

G_{13} : Category one of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{14} : Category two of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{15} : Category three of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

T_{13} : Category one of abandon the realistic philosophy of most working scientists,

T_{14} : Category two of abandon the realistic philosophy of most working scientists,

T_{15} : Category three of abandon the realistic philosophy of most working scientists,

Module Two

G_{16} : Category one of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{17} : Category two of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{18} : Category three of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

T_{16} : Category one of revision of concept of space-time

T_{17} : Category two of revision of concept of space-time

T_{18} : Category three of revision of concept of space-time

Module three

G_{20} : Category one of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{21} : Category two of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

G_{22} : Category three of Bell's theorem (We are talking of the systems to which bell's Theorem holds and the concomitant and corporate characterstics thereof)

T_{20} : Category one of Either one must totally abandon the realistic philosophy of most working scientists or dramatically revision of concept of space-time

T_{21} : Category two of Either one must totally abandon the realistic philosophy of most working scientists or dramatically revision of concept of space-time

T_{22} : Category three of Either one must totally abandon the realistic philosophy of most working scientists or dramatically revision of concept of space-time

Module four

G_{24} : Category one of factorizable, stochastic model

G_{25} : Category two of factorizable, stochastic model

G_{26} : Category three of factorizable, stochastic model

T_{24} : Category one of deterministic hidden-variables model for the experiment

T_{25} : Category two of deterministic hidden-variables model for the experiment

T_{26} : Category three of deterministic hidden-variables model for the experiment

Module five

G_{28} : Category one of joint distributions for all observables of the experiment, returning the experimental probabilities.

G_{29} : Category two of joint distributions for all observables of the experiment, returning the experimental probabilities.

G_{30} : Category three of joint distributions for all observables of the experiment, returning the

experimental probabilities.

T_{28} : Category one of factorizable, stochastic model

T_{29} : Category two of factorizable, stochastic model

T_{30} : Category three of factorizable, stochastic model

Module six

G_{32} : Category one of factorizable, stochastic model

G_{33} : Category two of factorizable, stochastic model

G_{34} : Category three of factorizable, stochastic model

T_{32} : Category one of joint distributions for all observables of the experiment, returning the experimental probabilities.

T_{33} : Category two of joint distributions for all observables of the experiment, returning the experimental probabilities.

T_{34} : Category three of joint distributions for all observables of the experiment, returning the experimental probabilities.

Module seven

G_{36} : Category one of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

G_{37} : Category two of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

G_{38} : Category three of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

T_{36} : Category one of factorizable, stochastic model

T_{37} : Category two of factorizable, stochastic model

T_{38} : Category three of factorizable, stochastic model

Module eight

G_{40} : Category one of Spatio temporal actualization of Bell's inequalities

G_{41} : Category two of Spatio temporal actualization of Bell's inequalities

G_{42} : Category three of Spatio temporal actualization of Bell's inequalities

T_{40} : Category one of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

T_{41} : Category two of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

T_{42} : Category three of well-defined, compatible joint distributions for all pairs and triples of commuting and noncommuting observables

Module Nine

G_{44} : Category one of factorizable, stochastic model

G_{45} : Category two of factorizable, stochastic model

G_{46} : Category three of factorizable, stochastic model

T_{44} : Category one of Spatio temporal actualization of Bell's inequalities

T_{45} : Category two of Spatio temporal actualization of Bell's inequalities

T_{46} : Category three of Spatio temporal actualization of Bell's inequalities

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15}$ 6

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16}$ 7

$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17}$ 8

$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18}$ 9

$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16}$ 10

$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17}$ 11

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30}$ 27

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45}$ 53

$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46}$ 54

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{matrix}(a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}\\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}\\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{matrix}(a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}\\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}\\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{matrix}(a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}\\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}\\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}\end{matrix}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[\begin{matrix}(b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{matrix}(b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}\\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}\\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}\end{matrix}\right]T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) \end{bmatrix} G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) \end{bmatrix} G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) \end{bmatrix} G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3
$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3
$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3
$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3
$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3
$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3
$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3
$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients*

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) \quad +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) \quad +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) \quad +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients

for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \qquad 82$$

$$\frac{dT_{29}}{dt}=(b_{29})^{(5)}T_{28}-\left[\begin{array}{ccc}(b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{29} \qquad 83$$

$$\frac{dT_{30}}{dt}=(b_{30})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{30} \qquad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
$are\ first\ detrition\ coefficients\ \ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ \ third\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} & -(b''_{36})^{(7)}(G_{39},t) & -(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} & -(b''_{37})^{(7)}(G_{39},t) & -(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} & -(b''_{38})^{(7)}(G_{39},t) & -(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1)}(G,t) & -(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)$ are second

detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)} & \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\begin{bmatrix} (b'_{45})^{(9)} & \boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$lim_{T_2\to\infty}(a_i'')^{(1)}\,(T_{14},t)=(p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}\,(G,t)=\ \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)},(p_i)^{(1)},\ \ (r_i)^{(1)}}$ are positive constants and $\boxed{i=13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},$ are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\ ,\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)}+(a_i')^{(1)}+\ (\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)}+(b_i')^{(1)}+\ (\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le\left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le\ \ (r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}\,(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=\ \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26$$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$ 118

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)}=1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}},\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$$

Definition of $(\hat{P}_{24})^{(4)},(\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)},(\hat{A}_{24})^{(4)}and\,(\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)},(a_i')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(p_i)^{(4)},\ (r_i)^{(4)},i=24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)}+(a_i')^{(4)}+\ (\hat{A}_{24})^{(4)}+\ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}]<1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)}+(b_i')^{(4)}+\ (\hat{B}_{24})^{(4)}+\ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}]<1$$

Where we suppose

$$(a_i)^{(5)},(a_i')^{(5)},(a_i'')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(b_i'')^{(5)}>0,\quad i,j=28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)},(b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)},\ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t)\le(p_i)^{(5)}\le(\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big)\le\ (r_i)^{(5)}\le(b_i')^{(5)}\le(\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}\,(T_{29},t)=(p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}\,(G_{31},t)=\ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)},(p_i)^{(5)},\ (r_i)^{(5)}}$ are positive constants and $\boxed{i=28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t)-(a_i'')^{(5)}(T_{29},t)|\le(\hat{k}_{28})^{(5)}|T_{29}-T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t)-(b_i'')^{(5)}\big((G_{31}),t\big)|<(\hat{k}_{28})^{(5)}||(G_{31})-(G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$

and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}<1$$

Definition of $(\hat{P}_{28})^{(5)},(\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)},(\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)},(a_i')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(p_i)^{(5)},\ (r_i)^{(5)},i=28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)}+(a_i')^{(5)}+\ (\hat{A}_{28})^{(5)}+\ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}]<1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)}+(b_i')^{(5)}+\ (\hat{B}_{28})^{(5)}+\ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}]<1$$

Where we suppose

$$(a_i)^{(6)},(a_i')^{(6)},(a_i'')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(b_i'')^{(6)}>0,\quad i,j=32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)},(b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t)\le(p_i)^{(6)}\le(\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t)\le\ (r_i)^{(6)}\le(b_i')^{(6)}\le(\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}),t\right)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\left((G_{35}),t\right)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\widehat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$ 132

$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It

is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+\ (\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[\ (b_i)^{(8)}+(b_i')^{(8)}+\ (\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-\ T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{20})^{(3)}$, $T_i^0 \leq (\hat{Q}_{20})^{(3)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{24})^{(4)}$, $T_i^0 \leq (\hat{Q}_{24})^{(4)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)}t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)}t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\overline{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167 that

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that (172)

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that (174)

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T}$: $\left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$G_{13}^{(2)}|(a_{13}'')^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a_{13}'')^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a_{13}')^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a_{13}'')^{(1)}$ and $(b_{13}'')^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a_{14}')^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\ (G(t),t))=(b_{14}')^{(1)}$ then $T_{14}\ \to\infty$. (189)

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\ (t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$

$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205

unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b''_{18})^{(2)}\left((G_{19})(t),t\right)=(b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\hat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 215

$(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself (224)

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric (225)

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\leq \tag{226}$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. (227)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2 \; and \; \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25} \; and \; G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25} \; or \; G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24} \; is$ bounded, from below, the same property holds for $G_{25} \; and \; G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}((G_{31}),(T_{31}))$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 239

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(5)} t\right)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad (245)$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad (246)$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (250)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{40})^{(8)}|G_{41}^{(1)}-G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}|(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})-(a''_{40})^{(8)}(T_{41}^{(2)},s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$|(G_{43})^{(1)}-(G_{43})^{(2)}|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)})d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(8)}t)}>0 \quad \text{for t} > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1,((\widehat{M}_{40})^{(8)})_2$ *and* $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40}<(\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt}\le((\widehat{M}_{40})^{(8)})_1-(a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})} \right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{44})^{(9)} \left|G_{45}^{(1)} - G_{45}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\, ds_{(44)} +$$
$$\int_0^t \{(a'_{44})^{(9)} \left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$G_{44}^{(2)} \left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\} ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)}\right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)}; (G_{47})^{(2)}, (T_{47})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ *and* $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(9)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, $\boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, $\boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16} e^{((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)})\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16} e^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$

$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

332

$$G^0_{24} e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24} e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)}-(a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)},$ $\boldsymbol{if}$ $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}$ $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)},$ $\boldsymbol{if}$ $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)},$ $\boldsymbol{if}$ $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)},$ $\boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$

$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$

Definition of $(\nu_1)^{(6)}, (\nu_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, \nu^{(6)}, u^{(6)}$: 350

By $(\nu_1)^{(6)} > 0$, $(\nu_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)}, , (\bar{\nu}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)}$, ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)}$,
and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^{0}}{G_{33}^{0}}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)}$, ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0$, $(v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0$, $(\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (v_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (v_0)^{(7)}, (m_1)^{(7)} = (v_1)^{(7)}, \ \boldsymbol{if}\ (v_0)^{(7)} < (v_1)^{(7)}$$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (\bar{v}_1)^{(7)}, \boldsymbol{if}\ (v_1)^{(7)} < (v_0)^{(7)} < (\bar{v}_1)^{(7)},$$

and $\boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (v_1)^{(7)}, (m_1)^{(7)} = (v_0)^{(7)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(7)} < (v_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G^0_{36} e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t}\leq G_{38}(t)\leq\right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\leq T_{38}(t)\leq \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)}\leq-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\leq-(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)}\leq-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\leq-(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0$, $(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0$, $(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0$, $(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0$, $(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, $\boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}\left((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)}\right)}\left[e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}\left((S_1)^{(8)} - (a'_{42})^{(8)}\right)}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}\left((R_1)^{(8)} - (b'_{42})^{(8)}\right)}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}\left((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)}\right)}\left[e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, $\boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, $\boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)}\right) \leq \frac{dv^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also

defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)\le\frac{dv^{(4)}}{dt}\le-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_4)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)},(v_0)^{(4)}$:-

For $0<\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}<(v_1)^{(4)}<(\bar{v}_1)^{(4)}$

$$v^{(4)}(t)\ge\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}\quad,\quad\boxed{(C)^{(4)}=\frac{(v_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(v_2)^{(4)}}}$$

it follows $(v_0)^{(4)}\le v^{(4)}(t)\le(v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(4)}=\frac{(\bar{v}_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)}\le v^{(4)}(t)\le(\bar{v}_1)^{(4)}$

If $0<(v_1)^{(4)}<(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}<(\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)}\le\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}\le v^{(4)}(t)\le$$

$$\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(\bar{v}_1)^{(4)}$$

If $0<(v_1)^{(4)}\le(\bar{v}_1)^{(4)}\le\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)}\le v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)}\le v^{(4)}(t)\le(m_1)^{(4)},\quad\boxed{v^{(4)}(t)=\frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)}\le u^{(4)}(t)\le(\mu_1)^{(4)},\quad\boxed{u^{(4)}(t)=\frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)}=\frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)}=\frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)}=\frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t)=\frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t)=\frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \le \frac{dv^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \ge \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \le v^{(7)}(t) \le (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \le \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \le v^{(7)}(t) \le (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \le \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \le v^{(7)}(t) \le$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \le (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \le (\bar{v}_1)^{(7)} \le \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \le v^{(7)}(t) \le \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \le (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \le v^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain

421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get

422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)}T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(d) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}\mathrm{G}_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}\mathrm{G}_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502
$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{\left[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})\right]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{\left[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})\right]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{\left[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)\right]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{\left[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)\right]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \tag{513}$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b''_i)^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \ , \ \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(6)}$ *and* $(b''_i)^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b''_i)^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(7)}$ *and* $(b''_i)^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\right)\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\Big(\big((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\big)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\Big)$$

$$\Big(\big((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\big)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\Big)$$

$$\Big(\big((\lambda)^{(8)}\big)^{2}+\big((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\big)(\lambda)^{(8)}\Big)$$

$$\Big(\big((\lambda)^{(8)}\big)^{2}+\big((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\big)(\lambda)^{(8)}\Big)$$

$$+\Big(\big((\lambda)^{(8)}\big)^{2}+\big((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\big)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\big((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\big)\big((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\big)$$

$$\Big(\big((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\big)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\big((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\big)$$

$$\Big[\Big(\big((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\big)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\Big)\Big]$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\Big(\big((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\big)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\Big(\big((\lambda)^{(9)}\big)^{2}+\big((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\big)(\lambda)^{(9)}\Big)$$

$$\Big(\big((\lambda)^{(9)}\big)^{2}+\big((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\big)(\lambda)^{(9)}\Big)$$

$$+\Big(\big((\lambda)^{(9)}\big)^{2}+\big((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\big)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\big((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\big)\big((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\big)$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIVE

Local Realism, Bell Inequalities, Correlation Functions, Information-Theoretic Bell Inequalities, Hidden Quantum Nonlocality, Spin States Et Al: Boni Pastoris

Est Tondere Pecus Non Deglubere- Adde Parvum Parvo Magnus Acervus Erit (Ovid) Model

INTRODUCTION—VARIABLES USED

Wringing out better Bell inequalities Samuel L Braunstein and Carlton M Cavesb

(1) Local realism implies (eb) constraints on the statistics of two physically separated systems.
(2) These constraints, known collectively as Bell inequalities, **can be (eb) violated** by quantum mechanics. In other words quantum mechanics **violates (e)** Bell's inequality.
(3) The standard Bell inequalities apply to a pair of two-state systems and **constrain (e)** the value of some linear combination of correlation functions between the two systems.
(4) Authors generalize these standard Bell inequalities in two ways. First, we "chain" the Clauser-Horne-Shimony-Holt Bell inequality **to (eb)obtain** chained correlation Bell inequalities for two-state systems
(5) Real experiment shows that these chained Bell inequalities **lead (eb) to** stronger quantum violations. Second, we formulate information-theoretic Bell inequalities, which are written in terms of the average information obtained in several measurements on a pair of physically separated systems (not just two-state systems); these information Bell inequalities have an appealing interpretation: if local realism holds, **then (eb)** two systems must carry an amount of information consistent with the inequality.

Hidden quantum nonlocality revealed by local filters N. Gisin Group of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland http://dx.doi.org/10.1016/S0375-9601(96)80001-6, PHYSICS Letters A Volume 210, Issue 3, 8 January 1996, Pages 151–156

(6) Family of two spin states with the following properties is presented, these states are "local" in the sense that they do not violate any Bell—CHSH inequality which, however, after each spins interacts with two independent local environments, and the resulting states of the two spin systems **violate (e) a** Bell inequality.
(7) In particular, for mixed states the (e)violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open) and the concept of nonlocality for mixed states differ
(8) The fact that some dissipative environments can increase quantum correlations **might (e) be useful for** quantum computation.
(9) Careless application of generalized quantum measurements **can (e) violates** Bell's inequality by more than $2\sqrt{2}$, even for mixtures of product states.

NOTATION

Module One

G_{13} : Category one of Local realism

G_{14} : Category two of Local realism

G_{15} : Category three of Local realism

T_{13} : Category one of constraints on the statistics of two physically separated systems

T_{14} : Category two of constraints on the statistics of two physically separated systems

T_{15} : Category three of constraints on the statistics of two physically separated systems

Module Two

G_{16} : Category one of Bell's inequality

G_{17} : Category two of Bell's inequality

G_{18} : Category three of Bell's inequality

T_{16} : Category one of quantum mechanics

T_{17} : Category two of quantum mechanics

T_{18} : Category three of quantum mechanics

Module three

G_{20} : Category one of value of some linear combination of correlation functions between the two systems.

G_{21} : Category two of value of some linear combination of correlation functions between the two systems.

G_{22} : Category three of value of some linear combination of correlation functions between the two systems.

T_{20} : Category one of standard Bell inequalities apply to a pair of two-state systems

T_{21} : Category two of standard Bell inequalities apply to a pair of two-state systems

T_{22} : Category three of standard Bell inequalities apply to a pair of two-state systems

Module four

G_{24} : Category one of Clauser-Horne-Shimony-Holt Bell inequality (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{25} : Category two of Clauser-Horne-Shimony-Holt Bell inequality (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{26} : Category three of Clauser-Horne-Shimony-Holt Bell inequality (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{24} : Category one of Bell inequalities (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{25} : Category two of Bell inequalities

T_{26} : Category three of Bell inequalities

Module five

G_{28} : Category one of Result of module four

G_{29} : Category two of Result of module four

G_{30} : Category three of Result of module four

T_{28} : Category one of chained correlation Bell inequalities for two-state systems

T_{29} : Category two of chained correlation Bell inequalities for two-state systems

T_{30} : Category three of chained correlation Bell inequalities for two-state systems

Module six

G_{32} : Category one of chained Bell inequalities

G_{33} : Category two of chained Bell inequalities

G_{34} : Category three of chained Bell inequalities

T_{32} : Category one of stronger quantum violations

T_{33} : Category two of stronger quantum violations

T_{34} : Category three of stronger quantum violations

Module seven

G_{36} : Category one of local realism holds

G_{37} : Category two of local realism holds

G_{38} : Category three of local realism holds

T_{36} : Category one of two systems must carry an amount of information consistent with the inequality

T_{37} : Category two of two systems must carry an amount of information consistent with the inequality

T_{38} : Category three of two systems must carry an amount of information consistent with the inequality

Module eight

Family of two spin states with the following properties is presented, these states are "local" in the sense that they do not violate any Bell—CHSH inequality which, however, after each spins interacts with two independent local environments, and the resulting states of the two spin systems **violate (e) a** Bell inequality.

G_{40} : Category one of Bell inequality (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{41} : Category two of Bell inequality

G_{42} : Category three of Bell inequality

T_{40} : Category one of Family of two spin states with the following properties is presented, these states are "local" in the sense that they do not violate any Bell—CHSH inequality which, however, after each spins interacts with two independent local environments, and the resulting states of the two spin systems

T_{41} : Category two of Family of two spin states with the following properties is presented, these states are "local" in the sense that they do not violate any Bell—CHSH inequality which, however, after each spins interacts with two independent local environments, and the resulting states of the two spin systems

T_{42} : Category three of Family of two spin states with the following properties is presented, these states are "local" in the sense that they do not violate any Bell—CHSH inequality which, however, after each spins interacts with two independent local environments, and the resulting states of the two spin systems

Module Nine

G_{44} : Category one of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

G_{45} : Category two of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

G_{46} : Category three of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

T_{44} : Category one of concept of nonlocality for mixed states

T_{45} : Category two of concept of nonlocality for mixed states

T_{46} : Category three of concept of nonlocality for mixed states

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad (1)$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad (2)$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad (3)$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad (4)$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad (5)$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad (7)$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad (8)$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad (9)$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad (10)$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad (11)$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad (13)$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad (14)$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad (15)$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad (16)$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad (17)$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad (19)$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad (20)$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad (21)$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad (22)$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad (23)$$

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32}$ 31

$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33}$ 32

$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34}$ 33

$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32}$ 34

$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33}$ 35

$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34}$ 36

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36}$ 37

$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37}$ *38*

$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38}$ *39*

$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36}$ *40*

$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37}$ *41*

$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38}$ *42*

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40}$ *43*

$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41}$ *44*

$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42}$ *45*

$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40}$ *46*

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\left[-(b''_{24})^{(4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\left[-(b''_{25})^{(4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\left[-(b''_{26})^{(4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\left[+(a''_{28})^{(5)}(T_{29},t)\right] & \left[+(a''_{24})^{(4,4,)}(T_{25},t)\right] & \left[+(a''_{32})^{(6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)\right] & \left[+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)\right] & \left[+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)\right] & \left[+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)\right] & \left[+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)\right] \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) \quad +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) \quad +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) \quad +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$

Eighth augmentation coefficients $\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) \\ +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & & \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + \ (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + \ (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + \ (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad (118)$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \quad (122)$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}),t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \quad (123)$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: (124)

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}((G_{31}),t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: (125)

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: (126)

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \quad (127)$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t) = (p_i)^{(6)}$ 128

$\lim_{G\to\infty}(b_i'')^{(6)}\,((G_{35}),t) = \ (r_i)^{(6)}$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$

$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}((G_{35}),t)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37},t) = (p_i)^{(7)}$

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}',t) - (a_i'')^{(7)}(T_{37},t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})',t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}',t)$ and$(a_i'')^{(7)}(T_{37},t)$. (T_{37}',t) and (T_{37},t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ \ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41},t) \le (p_i)^{(8)} \le (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}),t) \le \ (r_i)^{(8)} \le (b_i')^{(8)} \le (\hat{B}_{40})^{(8)} \qquad 139$$

$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)}$ 140

$\lim_{G\to\infty}(b_i'')^{(8)}\big((G_{43}),t\big) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t}$ 142

$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\big((G_{43}),t\big)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46$ 146 A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$

$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)}and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}\left(s_{(13)}\right) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}\left(s_{(13)}\right) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}\left(s_{(16)}\right) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}\left(T_{17}\left(s_{(16)}\right), s_{(16)}\right)\right) G_{16}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^{0} + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^{0} + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^{0}) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^{0}\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^{0}}{G_{17}^{0}}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^{0} + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^{0} + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^{0}) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^{0}\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^{0}}{G_{21}^{0}}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^{0} + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^{0} + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_0^t\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\bar{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left(e^{(\bar{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\bar{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left(e^{(\bar{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\bar{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left(e^{(\bar{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\bar{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(P_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_{0}^{t}\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_{+}$.

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a''_{i})^{(1)}$ and $(b''_{i})^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15}. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205
unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}}$:$\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}((G_{23}), (T_{23}))$

It results 213

$$|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)}|G_{21}^{(1)} - G_{21}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)})|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}|(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)}) - (a_{20}'')^{(3)}(T_{21}^{(2)}, s_{(20)})|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|{G_{23}}^{(1)} - {G_{23}}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad (214)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(3)}$ and $(b''_i)^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ (216)

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(3)} - (a''_i)^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b'_i)^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: (217)

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. (218)

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. (219)

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b''_i)^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to (220)

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}), (\widetilde{T_{27}})\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{24})^{(4)}\left|G_{25}^{(1)} - G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)} +$$

$$\int_0^t\{(a_{24}')^{(4)}\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a_{24}'')^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t} \leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a_{24}')^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)}|G_{29}^{(1)} - G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)})|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)}G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29}\ \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{32})^{(6)}|G_{33}^{(1)}-G_{33}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)|\;e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$|(G_{35})^{(1)}-(G_{35})^{(2)}|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq$$ 259
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 275

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\ t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations

$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)} v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)} u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}$ $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}$ $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}$ $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le\right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t} \right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)} t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t} \right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$, $(v_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,
and $\boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$
$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}$ $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}$ $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,

and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)},\ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{\left((S_1)^{(7)} - (p_{36})^{(7)}\right)t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\right] + G_{38}^0 e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t}\right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, ***if*** $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, ***if*** $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,

and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, ***if*** $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}$ $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\le\frac{dv^{(1)}}{dt}\le-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\ge\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

$$\text{it follows } (\nu_0)^{(1)}\le v^{(1)}(t)\le(\nu_1)^{(1)}$$

In the same manner , we get 384

$$v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\le v^{(1)}(t)\le(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\le\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\le v^{(1)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\le(\bar{\nu}_1)^{(1)}\le\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\le v^{(1)}(t)\le\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\le(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\le v^{(1)}(t)\le(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\le u^{(1)}(t)\le(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)\le\frac{dv^{(9)}}{dt}\le-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(9)},(v_0)^{(9)}$:-

For $0<\boxed{(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}<(v_1)^{(9)}<(\bar{v}_1)^{(9)}$

$$v^{(9)}(t)\ge\frac{(v_1)^{(9)}+(C)^{(9)}(v_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)}-(v_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)}-(v_0)^{(9)}\right)t\right]}}\quad,\quad\boxed{(C)^{(9)}=\frac{(v_1)^{(9)}-(v_0)^{(9)}}{(v_0)^{(9)}-(v_2)^{(9)}}}$$

it follows $(v_0)^{(9)}\le v^{(9)}(t)\le(v_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t)\le\frac{(\bar{v}_1)^{(9)}+(\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(9)}=\frac{(\bar{v}_1)^{(9)}-(v_0)^{(9)}}{(v_0)^{(9)}-(\bar{v}_2)^{(9)}}}$$

From which we deduce $(v_0)^{(9)}\le v^{(9)}(t)\le(\bar{v}_1)^{(9)}$

If $0<(v_1)^{(9)}<(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}<(\bar{v}_1)^{(9)}$ we find like in the previous case,

$$(v_1)^{(9)}\le\frac{(v_1)^{(9)}+(C)^{(9)}(v_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)}-(v_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((v_1)^{(9)}-(v_2)^{(9)}\right)t\right]}}\le v^{(9)}(t)\le$$

$$\frac{(\bar{v}_1)^{(9)}+(\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}\le(\bar{v}_1)^{(9)}$$

If $0<(v_1)^{(9)}\le(\bar{v}_1)^{(9)}\le\boxed{(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(v_1)^{(9)}\le v^{(9)}(t)\le\frac{(\bar{v}_1)^{(9)}+(\bar{C})^{(9)}(\bar{v}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{v}_1)^{(9)}-(\bar{v}_2)^{(9)}\right)t\right]}}\le(v_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)}\le v^{(9)}(t)\le(m_1)^{(9)},\quad\boxed{v^{(9)}(t)=\frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)}\le u^{(9)}(t)\le(\mu_1)^{(9)},\quad\boxed{u^{(9)}(t)=\frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(e) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T_{14}^* :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)}G_{25}^{*}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)}G_{25}^{*}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}\quad,\quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}\quad,\quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}\quad,\quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}\quad,\quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof:_Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i \qquad , T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*})=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G^*_{13}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G^*_{14}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G^*_{15}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a_{24}')^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a_{25}')^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a_{26}')^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b_{24}')^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^*\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b_{25}')^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^*\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b_{26}')^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^*\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a_{28}')^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a_{29}')^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a_{30}')^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b_{28}')^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T_{28}^*\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b_{29}')^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T_{29}^*\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b_{30}')^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T_{30}^*\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \ , \ \frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \ , \ \frac{\partial (b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \ , \ \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b''_i)^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left((\lambda)^{(1)}\right)^2+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\Big)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\Big]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)})$$

$$\left[\left(((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$+\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)})$$

$$\left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$+\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T^*_{37} + (b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\} = 0$$

$$+$$

$$(\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)})$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIX

Standard Approach To Quantum Systems, Existence Of Hidden States, Markov Context, Hidden Quantum Nonlocality, Et Al.: De Gustibus Non Est Disputandum -Legum Servi Sumus Ut Liberi Esse Possimus

INTRODUCTION—VARIABLES USED

A Markovian Model for Joint Observations, Bell's Inequality and Hidden States Ulrich Faigle, Alexander Schönhuth

(1) standard approach to quantum systems studies length preserving linear transformations of wave functions, the Markov picture focuses on trace preserving operators on the space of Hermitian (self-adjoint) matrices thus the Markov approach extends the standard one and provides(eb) **a** refined analysis of measurements and quantum Markov chains, in particular, Bell's inequality becomes structurally clear.
(2) Hidden state models are natural in the Markov context and in particular, a violation of Bell's inequality **is seen to (eb) be compatible** with the existence of hidden states.
(3) The Markov model moreover clarifies the role of the "negative probabilities" in Feynman's analysis of the EPR paradox. 60J99, 81P45, 81P68 Cite as: arXiv: 1011.1295 [cs.IT] (or arXiv: 1011.1295v2 [cs.IT] for this version)

Hidden quantum nonlocality revealed by local filters N. Gisin Group of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland http://dx.doi.org/10.1016/S0375-9601(96)80001-6, PHYSICS Letters A Volume 210, Issue 3, 8 January 1996, Pages 151–156.

(4) The fact that some dissipative environments can increase quantum correlations might (e) be useful for quantum computation.
(5) Careless application of generalized quantum measurements **can (e) violates** Bell's inequality by more than $2\sqrt{2}$, even for mixtures of product states.

Bell's inequality holds for all non-product states: N. Gisin

(6) Authors prove that any non-product state of two-particle systems **violates(e)** a Bell inequality

A theory of quantum gravity based on quantum computation Seth Lloyd

(7) This paper proposes a method of unifying quantum mechanics and gravity **based (e) on** quantum computation.
(8) In this theory, fundamental processes are described in terms of pair wise interactions between quantum degrees of freedom.
(9) The geometry of space-time is a construct, derived (e) from the underlying quantum information processing.
(10) The computation **gives (eb) rise to** a superposition of four-dimensional spacetimes, each of which obeys the Einstein-Regge equations. The theory makes explicit predictions for the back-reaction of the metric to computational `matter,' black-hole evaporation, holography, and quantum cosmology. Comments: Plain TeX; 43 pages; 5 figures (pdf); this version with more extensive discussion of quantum cosmology Subjects: Quantum Physics (quant-ph); General Relativity and Quantum Cosmology (gr-qc) Cite as: arXiv: quant-ph/0501135 (or arXiv: quant-ph/0501135v8 for this version)

NOTATION

Module One

G_{13} : Category one of concept of nonlocality for mixed states

G_{14} : Category two of concept of nonlocality for mixed states

G_{15} : Category three of concept of nonlocality for mixed states

T_{13} : Category one of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

T_{14} : Category two of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

T_{15} : Category three of violation of Bell's inequality for mixed states (problem of classifying the nonlocal states of two spin systems is still open)

Module Two

G_{16} : Category one of standard approach to quantum systems studies length preserving linear transformations of wave functions, the Markov picture focuses on trace preserving operators on the space of Hermitian (self-adjoint) matrices thus the Markov approach extends the standard one

G_{17} : Category two of standard approach to quantum systems studies length preserving linear transformations of wave functions, the Markov picture focuses on trace preserving operators on the space of Hermitian (self-adjoint) matrices thus the Markov approach extends the standard one

G_{18} : Category three of standard approach to quantum systems studies length preserving linear transformations of wave functions, the Markov picture focuses on trace preserving operators on the space of Hermitian (self-adjoint) matrices thus the Markov approach extends the standard one

T_{16} : Category one of refined analysis of measurements and quantum Markov chains, in particular, Bell's inequality becomes structurally clear.

T_{17} : Category two of refined analysis of measurements and quantum Markov chains, in particular, Bell's inequality becomes structurally clear.

T_{18} : Category three of refined analysis of measurements and quantum Markov chains, in particular, Bell's inequality becomes structurally clear.

Module three

G_{20} : Category one of hidden state models are natural in the Markov context with a violation of Bell's inequality

G_{21} : Category two of hidden state models are natural in the Markov context with a violation of Bell's inequality

G_{22} : Category three of hidden state models are natural in the Markov context with a violation of Bell's inequality

T_{20} : Category one of existence of hidden states

T_{21} : Category two of existence of hidden states

T_{22} : Category three of existence of hidden states

Module four

Markov model moreover clarifies the role of the "negative probabilities" in Feynman's analysis of the EPR paradox

G_{24} : Category one of Markov model (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{25} : Category two of Markov model

G_{26} : Category three of Markov model

T_{24} : Category one of role of the "negative probabilities" in Feynman's analysis of the EPR paradox

T_{25} : Category two of role of the "negative probabilities" in Feynman's analysis of the EPR paradox

T_{26} : Category three of role of the "negative probabilities" in Feynman's analysis of the EPR paradox

Module five

G_{28} : Category one of quantum computation

G_{29} : Category two of quantum computation

G_{30} : Category three of quantum computation

T_{28} : Category one of some dissipative environments can increase quantum correlations

T_{29} : Category two of some dissipative environments can increase quantum correlations

T_{30} : Category three of some dissipative environments can increase quantum correlations

Module six

G_{32} : Category one of Bell's inequality by more than $2\sqrt{2}$, even for mixtures of product states.

G_{33} : Category two of Bell's inequality by more than $2\sqrt{2}$, even for mixtures of product states.

G_{34} : Category three of Bell's inequality by more than $2\sqrt{2}$, even for mixtures of product states.

T_{32} : Category one of Careless application of generalized quantum measurements

T_{33} : Category two of Careless application of generalized quantum measurements

T_{34} : Category three of Careless application of generalized quantum measurements

Module seven

G_{36} : Category one of Bell inequality (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{37} : Category two of Bell inequality

G_{38} : Category three of Bell inequality

T_{36} : Category one of any non-product state of two-particle systems

T_{37} : Category two of any non-product state of two-particle systems

T_{38} : Category three of any non-product state of two-particle systems

Module eight

G_{40} : Category one of gravity based on quantum computation.

G_{41} : Category two of gravity based on quantum computation.

G_{42} : Category three of gravity based on quantum computation.

T_{40} : Category one of quantum mechanics

T_{41} : Category two of quantum mechanics

T_{42} : Category three of quantum mechanics

Module Nine

G_{44} : Category one of fundamental processes

G_{45} : Category two of fundamental processes

G_{46} : Category three of fundamental processes

T_{44} : Category one of pair wise interactions between quantum degrees of freedom

T_{45} : Category two of pair wise interactions between quantum degrees of freedom

T_{46} : Category three of pair wise interactions between quantum degrees of freedom

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$

$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$
$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$
$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for

1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $\left[-(b''_{13})^{(1)}(G,t)\right]$, $\left[-(b''_{14})^{(1)}(G,t)\right]$, $\left[-(b''_{15})^{(1)}(G,t)\right]$ are first detrition coefficients for category 1, 2 and 3

$\left[-(b''_{16})^{(2,2,)}(G_{19},t)\right]$, $\left[-(b''_{17})^{(2,2,)}(G_{19},t)\right]$, $\left[-(b''_{18})^{(2,2,)}(G_{19},t)\right]$ are second detrition coefficients for category 1, 2 and 3

$\left[-(b''_{20})^{(3,3,)}(G_{23},t)\right]$, $\left[-(b''_{21})^{(3,3,)}(G_{23},t)\right]$, $\left[-(b''_{22})^{(3,3,)}(G_{23},t)\right]$ are third detrition coefficients for category 1, 2 and 3

$\left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right]$, $\left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right]$, $\left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right]$ are fourth detrition coefficients for category 1, 2 and 3

$\left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right]$, $\left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right]$, $\left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right]$ are fifth detrition coefficients for category 1, 2 and 3

$\left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right]$, $\left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right]$, $\left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right]$ are sixth detrition coefficients for category 1, 2 and 3

$\left[-(b''_{37})^{(7,7,)}(G_{39},t)\right]$, $\left[-(b''_{36})^{(7,7,)}(G_{39},t)\right]$, $\left[-(b''_{38})^{(7,7,)}(G_{39},t)\right]$ are seventh detrition coefficients for category 1, 2 and 3

$\left[-(b''_{40})^{(8,8)}(G_{43},t)\right]$ $\left[-(b''_{41})^{(8,8)}(G_{43},t)\right]$ $\left[-(b''_{42})^{(8,8)}(G_{43},t)\right]$ are eight detrition coefficients for category 1, 2 and 3

$\left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right]$, $\left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right]$, $\left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right]$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\left[+(a''_{16})^{(2)}(T_{17},t)\right] & \left[+(a''_{13})^{(1,1,)}(T_{14},t)\right] & \left[+(a''_{20})^{(3,3,3)}(T_{21},t)\right] \\ \left[+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)\right] & \left[+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)\right] & \left[+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{36})^{(7,7,7)}(T_{37},t)\right] & \left[+(a''_{40})^{(8,8,8)}(T_{41},t)\right] & \left[+(a''_{44})^{(9,9)}(T_{45},t)\right] \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\left[+(a''_{17})^{(2)}(T_{17},t)\right] & \left[+(a''_{14})^{(1,1,)}(T_{14},t)\right] & \left[+(a''_{21})^{(3,3,3)}(T_{21},t)\right] \\ \left[+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)\right] & \left[+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)\right] & \left[+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{37})^{(7,7,7)}(T_{37},t)\right] & \left[+(a''_{41})^{(8,8,8)}(T_{41},t)\right] & \left[+(a''_{45})^{(9,9)}(T_{45},t)\right] \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \tag{71}$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \tag{72}$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \tag{73}$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \tag{74}$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \tag{75}$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{matrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{matrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{matrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3
And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3
$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3
$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3
$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3
$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3
$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3
$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3
$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2, and 3
$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 91$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 92$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 93$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$
are eighth augmentation coefficient for 1,2,3
$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$
$$= (a_{44})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)}G_{45}$$
$$-\left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right] G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\widehat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\widehat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\widehat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants

$(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T_{25}'| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and $(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}\ (T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\ \big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - \ T_{41}'| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\widehat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ , \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (a_i)^{(8)} + (a_i')^{(8)} + \ (\hat{A}_{40})^{(8)} + \ (\hat{P}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ \ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\widehat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\widehat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\overline{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\overline{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\overline{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(1)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{16})^{(2)}|G_{17}^{(1)}-G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)|\;e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)}-(G_{19})^{(2)}|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\;(G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\widehat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{20})^{(3)}\right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})} \right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)} - G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\tilde{M}_{24})^{(4)}s_{(24)}}e^{(\tilde{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+\ (a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 228

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\leq\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25}\leq\left((\widehat{M}_{24})^{(4)}\right)_2=G_{25}^0+2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26}\leq\left((\widehat{M}_{24})^{(4)}\right)_3=G_{26}^0+2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t),t))=(b'_{25})^{(4)}$ then $T_{25}\ \to\infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a_{28}')^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{-(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a_{28}'')^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$G_{28}^{(2)}|(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a_{28}'')^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a_{28}')^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a_{28}'')^{(5)}$ and $(b_{28}'')^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\geq 0$

$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(5)}t\right)}>0\ \ \text{for t}>0$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\leq\left((\widehat{M}_{28})^{(5)}\right)_1-(a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29}\leq\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30}\leq\left((\widehat{M}_{28})^{(5)}\right)_3=G_{30}^0+2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t))=(b_{29}')^{(5)}$ then $T_{29}\ \to\infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ >t_5$

$(b_{29})^{(5)}-(b_i'')^{(5)}((G_{31})(t),t)<\varepsilon_5, T_{28}\ (t)>(m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)}, (T_{35})^{(1)}), ((G_{35})^{(2)}, (T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}), (\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}), (\widetilde{T_{35}})\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{-(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)}\left|(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a_{32}'')^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t), t(t), t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}), (\widetilde{T_{39}}) : \left((\widetilde{G_{39}}), (\widetilde{T_{39}})\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered 260

as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ and $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(7)} - (a_i'')^{(7)}(T_{37}(s_{(36)}), s_{(36)})\} ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t), t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is

unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right)=(b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)}+\left((\widehat{P}_{40})^{(8)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{40})^{(8)}\right]\leq(\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\;e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\,(G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279

A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself

The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$

It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and

respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\ (t) = 0\ and\ T_i\ (t) = 0$

From 99 to 44 it results

$$G_i\ (t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\ (t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\ (t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\ \frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)},\ \boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)},$ 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}\ (v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}\ (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\Big(\frac{(a_{22})^{(3)}G^0_{20}}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\Big[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\Big] + G^0_{22}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \quad 324$$

$$\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+\ G_{22}^{0}e^{-(a'_{22})^{(3)}t})$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$$

$$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$$

$$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)},(v_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},v^{(4)},u^{(4)}$: 329

By $(v_1)^{(4)}>0$, $(v_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_1)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)}>0$, $(\bar{v}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$(m_2)^{(4)}=(v_0)^{(4)},(m_1)^{(4)}=(v_1)^{(4)},$ ***if*** $(v_0)^{(4)}<(v_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}$ $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}$ $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}$ $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}$ $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}$ $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \leq G_{26}(t) \leq \right.$$

334

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \leq T_{26}(t) \leq$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,
and $\boxed{(v_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^{0}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G_{28}^{0}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t}-e^{-(S_2)^{(5)}t}\right]+G_{30}^{0}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left.\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+G_{30}^{0}e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$ and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$ and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0,(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0,(u_2)^{(7)}<0$ the roots of the equations
$(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0,(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0,(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)},\ \boldsymbol{if}\ (\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\bar{\nu}_1)^{(7)},\boldsymbol{if}\ (\nu_1)^{(7)}<(\nu_0)^{(7)}<(\bar{\nu}_1)^{(7)}$,
and $\boxed{(\nu_0)^{(7)}=\frac{G_{36}^{0}}{G_{37}^{0}}}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)}<(\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, , (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, ***if*** $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, ***if*** $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, ***if*** $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t}\le G_{42}(t)\le\right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\le T_{42}(t)\le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)}\le-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\le-(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)}\le-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\le-(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, ***if*** $(\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, ***if*** $(\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,

and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, ***if*** $(\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, ***if*** $(u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, ***if*** $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}$, ***if*** $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \leq \frac{d\nu^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\leq\frac{dv^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\geq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\leq v^{(3)}(t)\leq(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\leq v^{(3)}(t)\leq(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\leq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\leq v^{(3)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\leq(\bar{v}_1)^{(3)}\leq(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\leq v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\leq v^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$
In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows
$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also**

defines $(\nu_0)^{(8)}$ for the special case .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $v^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\ ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\ , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(f) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows

that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$$T^*_{24} = \frac{(b_{24})^{(4)}T^*_{25}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T^*_{26} = \frac{(b_{26})^{(4)}T^*_{25}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \ , \ \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\ G^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a_{13}')^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a_{14}')^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a_{15}')^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b_{13}')^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^*\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b_{14}')^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^*\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b_{15}')^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^*\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \ , \ \frac{\partial(b_i'')^{(2)}}{\partial G_j}(\ (G_{19})^*\ ) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(16)(j)}T_{16}^*\mathbb{G}_j\big) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}} (T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j} ((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)} G_{40}^* \mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)} G_{41}^* \mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)} G_{42}^* \mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)} T_{40}^* \mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)} T_{41}^* \mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)} T_{42}^* \mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is

$$(\lambda)^{(1)}+(b'_{15})^{(1)}-(r_{15})^{(1)})\{((\lambda)^{(1)}+(a'_{15})^{(1)}+(p_{15})^{(1)})$$

$$\left[\left(((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)})(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\right]$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(14)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\right)$$

$$+\left(((\lambda)^{(1)}+(a'_{14})^{(1)}+(p_{14})^{(1)})(q_{13})^{(1)}G^{*}_{13}+(a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(13)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\right)$$

$$\left((\lambda)^{(1)}\right)^{2}+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$\left((\lambda)^{(1)}\right)^{2}+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^{2}+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)})\{((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)})\{((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)})$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left((\lambda)^{(9)}\right)^2 + \left((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left((\lambda)^{(9)}\right)^2 + \left((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45} + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45} + (b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\} = 0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SEVEN

Superposition Of Four-Dimensional Spacetimes, Geometry Of Space-Time Is A Construct, Quantum Gravity From Quantum Computation Et Al. : A Caelo Usque Ad Centrum-A Bene Placito-Gesamtkunstwerk –Gestalt Model

INTRODUCTION—VARIABLES USED

One Response to Quantum Gravity from Quantum Computation: Jeff Hodges says: 1/25/2005 at 9:04 pm (I have deleted paragraphs due to spatial constraints. However the point author makes is clearly conspicuous)

From A3.4, pg 27:

(1) A fluctuating cosmological term then arises naturally…This picture of a fluctuating cosmological 'constant' with a scale given by the age of the universe **is (e&eb) consistent with** the causal set account of a fluctuating cosmological term.

From A3.5 (the necessary discussion about "scaling up" to the classical world):

(2) In other words, the computational universe naturally **gives(eb) rise to the** a semiclassical spacetime via the decoherent histories approach to quantum mechanics…Coarse-graining about stable histories yields hydrodynamic variables that approximately decohere and obey classical probabilites.

Bell inequality, Bell states and maximally entangled states for n qubits: N Gisin, H Bechmann-Pasquinucci Group of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland http://dx.doi.org/10.1016/S0375-9601(98)00516-7

(3) First, authors present a Bell-type inequality for n qubits, assuming that m out of the n qubits is independent. Quantum mechanics ***violates (e) this*** inequality by a ratio that increases exponentially with m.
(4) Hence an experiment on n qubits violating this inequality sets (eb) a lower bound on the number m of

entangled qubits.

(5) Next, they propose a definition of maximally entangled states of n qubits. For this purpose authors study five different criteria. Four of these criteria are found compatible. For any number n of qubits, **they (eb) determine** an orthogonal basis consisting of maximally entangled states generalizing the Bell states.

The Violation of Bell Inequalities in the Macro world Diederik Aerts, Sven Aerts, Jan Broekaert, Liane Gabora Foundations of Physics September 2000, Volume 30, Issue 9, pp 1387-1414 Springer Link

(6) Authors show that Bell inequalities can be violated in the macroscopic world. The macro world violation is illustrated using an example involving connected vessels of water. Authors show that whether the violation of inequalities occurs in the micro world or the macro world, it is the identification of nonidentical events that plays a crucial role.

(7) Specifically, they prove that if nonidentical events are consistently differentiated, Bell-type Pitowsky inequalities **are (eb) = e(e) no longer violated**, even for Bohm's example of two entangled spin 1/2 quantum particles.

NOTATION

Module One

G_{13} : Category one of underlying quantum information processing.

G_{14} : Category two of underlying quantum information processing.

G_{15} : Category three of underlying quantum information processing.

T_{13} : Category one of geometry of space-time is a construct

T_{14} : Category two of geometry of space-time is a construct

T_{15} : Category three of geometry of space-time is a construct

Module Two

G_{16} : Category one of computation

G_{17} : Category two of computation

G_{18} : Category three of computation

T_{16} : Category one of superposition of four-dimensional spacetimes, each of which obeys the Einstein-Regge equations. The theory makes explicit predictions for the back-reaction of the metric to computational `matter,' black-hole evaporation, holography, and quantum cosmology

T_{17} : Category two of superposition of four-dimensional spacetimes, each of which obeys the Einstein-Regge equations. The theory makes explicit predictions for the back-reaction of the metric to computational `matter,' black-hole evaporation, holography, and quantum cosmology

T_{18} : Category three of superposition of four-dimensional spacetimes, each of which obeys the Einstein-Regge equations. The theory makes explicit predictions for the back-reaction of the metric to computational `matter,' black-hole evaporation, holography, and quantum cosmology

Module three

Fluctuating cosmological ‘constant’ with a scale given by the age of the universe **is (e&eb) consistent with** the causal set account of a fluctuating cosmological term.

G_{20} : Category one of Fluctuating cosmological ‘constant’ with a scale given by the age of the universe

G_{21} : Category two of Fluctuating cosmological 'constant' with a scale given by the age of the universe

G_{22} : Category three of Fluctuating cosmological 'constant' with a scale given by the age of the universe

T_{20} : Category one of causal set account of a fluctuating cosmological term

T_{21} : Category two of causal set account of a fluctuating cosmological term

T_{22} : Category three of causal set account of a fluctuating cosmological term

Module four

G_{24} : Category one of causal set account of a fluctuating cosmological term

G_{25} : Category two of causal set account of a fluctuating cosmological term

G_{26} : Category three of causal set account of a fluctuating cosmological term

T_{24} : Category one of Fluctuating cosmological 'constant' with a scale given by the age of the universe

T_{25} : Category two of Fluctuating cosmological 'constant' with a scale given by the age of the universe

T_{26} : Category three of Fluctuating cosmological 'constant' with a scale given by the age of the universe

Module five

G_{28} : Category one of computational universe (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{29} : Category two of computational universe

G_{30} : Category three of computational universe

T_{28} : Category one of semiclassical spacetime via the decoherent histories approaches to quantum mechanics…

T_{29} : Category two of semiclassical spacetime via the decoherent histories approaches to quantum mechanics…

T_{30} : Category three of semiclassical spacetime via the decoherent histories approaches to quantum mechanics…

Module six

G_{32} : Category one of Coarse-graining about stable histories

G_{33} : Category two of Coarse-graining about stable histories

G_{34} : Category three of Coarse-graining about stable histories

T_{32} : Category one of hydrodynamic variables that approximately decohere and obey classical probabilites.

T_{33} : Category two of hydrodynamic variables that approximately decohere and obey classical probabilites.

T_{34} : Category three of hydrodynamic variables that approximately decohere and obey classical probabilites.

Module seven

G_{36} : Category one of Bell-type inequality for n qubits, assuming that m out of the n qubits is independent.

G_{37} : Category two of Bell-type inequality for n qubits, assuming that m out of the n qubits is independent.

G_{38} : Category three of Bell-type inequality for n qubits, assuming that m out of the n qubits is independent.

T_{36} : Category one of Quantum mechanics

T_{37} : Category two of Quantum mechanics

T_{38} : Category three of Quantum mechanics

Module eight

G_{40} : Category one of inequality by a ratio that increases exponentially with m.

G_{41} : Category two of inequality by a ratio that increases exponentially with m.

G_{42} : Category three of inequality by a ratio that increases exponentially with m.

T_{40} : Category one of Quantum mechanics

T_{41} : Category two of Quantum mechanics

T_{42} : Category three of Quantum mechanics

Module Nine

G_{44} : Category one of experiment on n qubits violating this inequality

G_{45} : Category two of experiment on n qubits violating this inequality

G_{46} : Category three of experiment on n qubits violating this inequality

T_{44} : Category one of lower bound on the number m of entangled qubits

T_{45} : Category two of lower bound on the number m of entangled qubits

T_{46} : Category three of lower bound on the number m of entangled qubits

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$

$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}), t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}), t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}), t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}), t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}),t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}),t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}),t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for

1,2,3
$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3
$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt}=(b_{13})^{(1)}T_{14}-\begin{bmatrix}(b'_{13})^{(1)}\,-(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t)\\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt}=(b_{14})^{(1)}T_{13}-\begin{bmatrix}(b'_{14})^{(1)}\,-(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) & -(b''_{21})^{(3,3,)}(G_{23},t)\\ -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,)}(G_{35},t)\\ -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt}=(b_{15})^{(1)}T_{14}-\begin{bmatrix}(b'_{15})^{(1)}\,-(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) & -(b''_{22})^{(3,3,)}(G_{23},t)\\ -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,)}(G_{35},t)\\ -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\end{bmatrix}T_{15} \quad (60)$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3
$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3
$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3
$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3
$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3
$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3
$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}\,+(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)}\,+(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad (63)$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9)}(T_{45},t)}$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad (66)$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}+(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{22} \tag{69}$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \tag{70}$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \qquad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[\begin{matrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[\begin{matrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[\begin{matrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$

are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $+(a''_{24})^{(4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,)}(T_{25},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2, and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)} & +(a''_{36})^{(7)}(T_{37},t) & +(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 91$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)} & +(a''_{37})^{(7)}(T_{37},t) & +(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 92$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)} & +(a''_{38})^{(7)}(T_{37},t) & +(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 93$$

Where $(a''_{36})^{(7)}(T_{37},t)$, $(a''_{37})^{(7)}(T_{37},t)$, $(a''_{38})^{(7)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt}$$

$$= (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$ *96*
$$= (a_{44})^{(9)}G_{45}$$
$$- \begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$- \begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)}G_{45}$$
$$- \begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\widehat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\widehat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\widehat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants

$(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \le (p_i)^{(4)} \le (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \le (r_i)^{(4)} \le (b_i')^{(4)} \le (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \le (\hat{k}_{24})^{(4)} |T_{25} - T_{25}'| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}\ (T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\ \big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - \ T_{41}'| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\widehat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ , \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (a_i)^{(8)} + (a_i')^{(8)} + \ (\hat{A}_{40})^{(8)} + \ (\hat{P}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ \ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}\left(s_{(13)}\right) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}\left(s_{(13)}\right) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\overline{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\overline{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\overline{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\widehat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(1)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)} G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{16})^{(2)}|G_{17}^{(1)}-G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)|\;e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)}-(G_{19})^{(2)}|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\;(G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\widehat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{20})^{(3)}\right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})} \right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)} - G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\tilde{M}_{24})^{(4)}s_{(24)}}e^{(\tilde{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 228

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\leq\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25}\leq\left((\widehat{M}_{24})^{(4)}\right)_2=G_{25}^0+2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26}\leq\left((\widehat{M}_{24})^{(4)}\right)_3=G_{26}^0+2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t),t))=(b'_{25})^{(4)}$ then $T_{25}\ \to\infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a_{28}')^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{-(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a_{28}'')^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$G_{28}^{(2)}|(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a_{28}'')^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a_{28}')^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a_{28}'')^{(5)}$ and $(b_{28}'')^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\geq 0$

$T_i\ (t)\geq T_i^0 e^{(-(b_i')^{(5)}t)}>0\ \ \text{for t}>0$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\leq\left((\widehat{M}_{28})^{(5)}\right)_1-(a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29}\leq\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30}\leq\left((\widehat{M}_{28})^{(5)}\right)_3=G_{30}^0+2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t))=(b_{29}')^{(5)}$ then $T_{29}\ \to\infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ >t_5$

$(b_{29})^{(5)}-(b_i'')^{(5)}((G_{31})(t),t)<\varepsilon_5, T_{28}\ (t)>(m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)} + T_j^0}{T_j^0}\right)} + (\widehat{Q}_{32})^{(6)}\right] \leq (\widehat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a_{32}')^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$$(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)}\left|(a_{32}'')^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a_{32}'')^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t), t(t), t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}), (\widetilde{T_{39}}) : \left((\widetilde{G_{39}}), (\widetilde{T_{39}})\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t} \leq \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\widehat{A}_{36})^{(7)} + (\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered 260

as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is

unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right)=(b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)}+\left((\widehat{P}_{40})^{(8)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{40})^{(8)}\right]\leq(\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\;e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\,(G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ \ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t\ > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279

A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and

respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\ (t) = 0\ and\ T_i\ (t) = 0$

From 99 to 44 it results

$$G_i\ (t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\ (t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\ (t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\ \frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}, \ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}, \ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}, \ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)},\ \boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)},$ 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}\ (v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}\ (v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\Big(\frac{(a_{22})^{(3)}G^0_{20}}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\Big[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\Big] + G^0_{22}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \quad 324$$

$$\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+\ G_{22}^{0}e^{-(a'_{22})^{(3)}t})$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\le T_{20}(t)\le\frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\le T_{22}(t)\le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$$

$$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$$

$$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)}\le-(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\le-(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)}\le-(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\le-(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)},(v_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},v^{(4)},u^{(4)}$: 329

By $(v_1)^{(4)}>0$, $(v_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_1)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)}>0$, $(\bar{v}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$$(m_2)^{(4)}=(v_0)^{(4)},(m_1)^{(4)}=(v_1)^{(4)},\ \boldsymbol{if}\ (v_0)^{(4)}<(v_1)^{(4)}$$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}$ $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}$ $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}$ $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}$ $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}$ $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \leq G_{26}(t) \leq \right.$$

334

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \leq T_{26}(t) \leq$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,
and $\boxed{(v_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,
and $\boxed{(u_0)^{(5)} = \frac{T^0_{28}}{T^0_{29}}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^{0}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G_{28}^{0}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t}-e^{-(S_2)^{(5)}t}\right]+G_{30}^{0}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left.\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+G_{30}^{0}e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$ and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$ and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0,(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0,(u_2)^{(7)}<0$ the roots of the equations
$(a_{37})^{(7)}\left(\nu^{(7)}\right)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0,(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0,(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)},\ \boldsymbol{if}\ (\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\bar{\nu}_1)^{(7)},\boldsymbol{if}\ (\nu_1)^{(7)}<(\nu_0)^{(7)}<(\bar{\nu}_1)^{(7)}$,
and $\boxed{(\nu_0)^{(7)}=\frac{G_{36}^{0}}{G_{37}^{0}}}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)}<(\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, , (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t}\le G_{42}(t)\le\right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\le T_{42}(t)\le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)}\le-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\le-(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)}\le-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\le-(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$
$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \le \frac{d\nu^{(1)}}{dt} \le -\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \ge \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \le \nu^{(1)}(t) \le (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \le \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \le \nu^{(1)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \le (\bar{\nu}_1)^{(1)} \le \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \le \nu^{(1)}(t) \le \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \le (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21},t)\right) - (a_{21}'')^{(3)}(T_{21},t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\leq\frac{dv^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\geq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\leq v^{(3)}(t)\leq(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\leq v^{(3)}(t)\leq(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\leq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\leq v^{(3)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\leq(\bar{v}_1)^{(3)}\leq(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\leq v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\leq v^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$
In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows
$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also**

defines $(\nu_0)^{(8)}$ for the special case .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then $(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $v^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\ ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\ , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(g) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows

that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$$T^*_{24} = \frac{(b_{24})^{(4)}T^*_{25}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T^*_{26} = \frac{(b_{26})^{(4)}T^*_{25}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a_{13}')^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a_{14}')^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a_{15}')^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b_{13}')^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^*\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b_{14}')^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^*\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b_{15}')^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^*\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}(\,(G_{19})^*\,) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(16)(j)}T_{16}^*\mathbb{G}_j\big) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ and $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ and $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}} (T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j} ((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)} G_{40}^* \mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)} G_{41}^* \mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)} G_{42}^* \mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)} T_{40}^* \mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)} T_{41}^* \mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)} T_{42}^* \mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is

$$(\lambda)^{(1)}+(b'_{15})^{(1)}-(r_{15})^{(1)})\{((\lambda)^{(1)}+(a'_{15})^{(1)}+(p_{15})^{(1)})$$

$$\left[\left(((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)})(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\right]$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(14)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\right)$$

$$+\left(((\lambda)^{(1)}+(a'_{14})^{(1)}+(p_{14})^{(1)})(q_{13})^{(1)}G^{*}_{13}+(a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(13)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$+\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)})\{((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$+\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)})\{((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)})$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left(\,(a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left(\,(b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left(\,(a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left(\,(a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left(\,(b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left(\,(a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left((\lambda)^{(9)}\right)^2 + \left((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$+\left((\lambda)^{(9)}\right)^2 + \left((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45} + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45} + (b_{45})^{(9)}s_{(44),(46)}T^*_{44}\Big)\} = 0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION EIGHT

Bell States And Maximally Entangled States For N Qubits, Pitowsky Inequalities, Bohm's Example Of Two Entangled Spin 1/2 Quantum Particles, Orthogonal Basis Consisting Of Maximally Entangled States Et Al. : Abundans Cautela Non Nocet -Abyssus Abyssum Invocate- Ad Majorem Dei Gloriam-Ad Lucem Model

INTRODUCTION—VARIABLES USED

Bell inequality, Bell states and maximally entangled states for n qubits: N Gisin, H Bechmann-Pasquinucci Group of Applied Physics, University of Geneva, 1211 Geneva 4, Switzerland http://dx.doi.org/10.1016/S0375-9601(98)00516-7

(1) Next, they propose a definition of maximally entangled states of n qubits. For this purpose authors study five different criteria. Four of these criteria are found compatible. For any number n of qubits, **they (eb) determine** an orthogonal basis consisting of maximally entangled states generalizing the Bell states.

The Violation of Bell Inequalities in the Macro world Diederik Aerts, Sven Aerts, Jan Broekaert, Liane Gabora Foundations of Physics September 2000, Volume 30, Issue 9, pp 1387-1414 Springer Link

(2) Authors show that Bell inequalities can be violated in the macroscopic world. The macro world violation is illustrated using an example involving connected vessels of water. Authors show that whether the violation of inequalities occurs in the micro world or the macro world, it is the identification of nonidentical events that plays a crucial role.

(3) Specifically, they prove that if nonidentical events are consistently differentiated, Bell-type

Pitowsky inequalities **are (eb) = e(e) no longer violated**, even for Bohm's example of two entangled spin 1/2 quantum particles.

(4) Authors also_show how Bell inequalities can be violated in cognition, specifically in the relationship between abstract concepts and specific instances of these concepts which supports the hypothesis that genuine quantum structure **exists (eb) in** the mind. This goes on to prove essentially that quantum mechanics is a subjective science.

(5) Authors introduce a model where the amount of nonlocality and the degree of quantum uncertainty are parameterized, and demonstrate that increasing nonlocality (nonlocality and the degree of quantum uncertainty are parameterized) produces **(eb)** increases the degree of violation, while increasing quantum uncertainty decreases the degree of violation.

Preparation of macroscopic superpositions in many-atom systems J. I. Cirac and P. Zoller © 1994 The American Physical Society URL: http://link.aps.org/doi/10.1103/PhysRevA.50.R2799 DOI: 10.1103/PhysRevA.50.R2799PACS: 03.65.Bz, 42.50.Wm

(6) Authors propose a technique to prepare two or more atoms in certain entangled states, **based (e) on** the interaction of the atoms with a cavity mode.

(7) After the atomic state is prepared, the cavity mode is left in the vacuum state, so dissipation does not affect the generated entangled states which **could (e) be used** for improved tests that challenge local realistic theories, and to examine related phenomena, which, thus far, have been realized only with photons

NOTATION

Module One.

G_{13} : Category one of number n of qubits

G_{14} : Category two of number n of qubits

G_{15} : Category three of number n of qubits

T_{13} : Category one of orthogonal basis consisting of maximally entangled states generalizing the Bell states

T_{14} : Category two of orthogonal basis consisting of maximally entangled states generalizing the Bell states

T_{15} : Category three of orthogonal basis consisting of maximally entangled states generalizing the Bell states

Module Two

G_{16} : Category one of identification of nonidentical events

G_{17} : Category two of identification of nonidentical events

G_{18} : Category three of identification of nonidentical events

T_{16} : Category one of violation of inequalities occurs in the micro world or the macro world

T_{17} : Category two of violation of inequalities occurs in the micro world or the macro world

T_{18} : Category three of violation of inequalities occurs in the micro world or the macro world

Module three

G_{20} : Category one of nonidentical events are consistently differentiated, Bell-type Pitowsky inequalities

G_{21} : Category two of nonidentical events are consistently differentiated, Bell-type Pitowsky inequalities

G_{22} : Category three of nonidentical events are consistently differentiated, Bell-type Pitowsky inequalities

T_{20} : Category one of Bohm's example of two entangled spin 1/2 quantum particles.

T_{21} : Category two of Bohm's example of two entangled spin 1/2 quantum particles.

T_{22} : Category three of Bohm's example of two entangled spin 1/2 quantum particles.

Module four

G_{24} : Category one of Bell inequalities

G_{25} : Category two of Bell inequalities

G_{26} : Category three of Bell inequalities

T_{24} : Category one of cognition

T_{25} : Category two of cognition

T_{26} : Category three of cognition

Module five

G_{28} : Category one of quantum structure

G_{29} : Category two of quantum structure

G_{30} : Category three of quantum structure

T_{28} : Category one of mind

T_{29} : Category two of mind

T_{30} : Category three of mind

Module six

G_{32} : Category one of increasing nonlocality (nonlocality and the degree of quantum uncertainty are parameterized)

G_{33} : Category two of increasing nonlocality (nonlocality and the degree of quantum uncertainty are parameterized)

G_{34} : Category three of increasing nonlocality (nonlocality and the degree of quantum uncertainty are parameterized)

T_{32} : Category one of increases the degree of violation

T_{33} : Category two of increases the degree of violation

T_{34} : Category three of increases the degree of violation

Module seven

G_{36} : Category one of increasing the degree of violation

G_{37} : Category two of increasing the degree of violation

G_{38} : Category three of increasing the degree of violation

T_{36} : Category one of quantum uncertainty

T_{37} : Category two of quantum uncertainty

T_{38} : Category three of quantum uncertainty

Module eight

Authors propose a technique to prepare two or more atoms in certain entangled states, **based (e) on** the interaction of the atoms with a cavity mode.

G_{40} : Category one of interaction of the atoms with a cavity mode.

G_{41} : Category two of interaction of the atoms with a cavity mode.

G_{42} : Category three of interaction of the atoms with a cavity mode.

T_{40} : Category one of preparation of two or more atoms in certain entangled states

T_{41} : Category two of preparation of two or more atoms in certain entangled states

T_{42} : Category three of preparation of two or more atoms in certain entangled states

Module Nine

G_{44} : Category one of improved tests that challenge local realistic theories, and to examine related phenomena, which, thus far, have been realized only with photons

G_{45} : Category two of improved tests that challenge local realistic theories, and to examine related phenomena, which, thus far, have been realized only with photons

G_{46} : Category three of improved tests that challenge local realistic theories, and to examine related phenomena, which, thus far, have been realized only with photons

T_{44} : Category one of atomic state preparation and consequent ramification of the cavity mode is left in the vacuum state, so dissipation does not affect the generated entangled states

T_{45} : Category two of atomic state preparation and consequent ramification of the cavity mode is left in the vacuum state, so dissipation does not affect the generated entangled states

T_{46} : Category three of atomic state preparation and consequent ramification of the cavity mode is left

in the vacuum state, so dissipation does not affect the generated entangled states

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32}$ 31

$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33}$ 32

$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34}$ 33

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}),t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}),t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t) & \end{bmatrix}G_{16} \quad 61$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t) & \end{bmatrix}G_{17} \quad 62$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t) & \end{bmatrix}G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) & \end{bmatrix}T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation

coefficients for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3
$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3
$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3
$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*
$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*
$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*
$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*
$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*
$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*
$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)}&\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)}&\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)}&\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}&\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}&\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}&\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}&\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}&\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)}&\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)}&\boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)}&\boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}&\boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}&\boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}&\boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}&\boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}&\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \quad (78)$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh

augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for

category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} =$$

94

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)} & +(a''_{40})^{(8)}(T_{41},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \begin{bmatrix} (a'_{44})^{(9)} \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \begin{bmatrix} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \begin{bmatrix} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \begin{bmatrix} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\leq(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\big((G_{19}),t\big)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)}=1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}},\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}<1$$

Definition of $(\hat{P}_{13})^{(2)},(\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},(\hat{A}_{16})^{(2)}and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)},(a_i')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(p_i)^{(2)},\ (r_i)^{(2)},i=16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)}+(a_i')^{(2)}+(\hat{A}_{16})^{(2)}+(\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)}+(b_i')^{(2)}+(\hat{B}_{16})^{(2)}+(\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 111$$

Where we suppose

$$(a_i)^{(3)},(a_i')^{(3)},(a_i'')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(b_i'')^{(3)}>0,\quad i,j=20,21,22 \quad 112$$

The functions $(a_i'')^{(3)},(b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)},\ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t)\leq(p_i)^{(3)}\leq(\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t)\leq\ (r_i)^{(3)}\leq(b_i')^{(3)}\leq(\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t)=(p_i)^{(3)} \quad 113$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t)=\ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)},(p_i)^{(3)},\ (r_i)^{(3)}}$ are positive constants and $\boxed{i=20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\le(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\le(p_i)^{(4)}\le(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\le\ (r_i)^{(4)}\le(b_i')^{(4)}\le(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\le(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It

is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)}=1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}},\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}<1$$

Definition of $(\hat{P}_{24})^{(4)},(\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)},(\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)},(a_i')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}, i=24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)}+(a_i')^{(4)}+(\hat{A}_{24})^{(4)}+(\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}]<1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)}+(b_i')^{(4)}+(\hat{B}_{24})^{(4)}+(\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}]<1$$

Where we suppose

$$(a_i)^{(5)},(a_i')^{(5)},(a_i'')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(b_i'')^{(5)}>0,\quad i,j=28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)},(b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)},\ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t)\le(p_i)^{(5)}\le(\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big)\le\ (r_i)^{(5)}\le(b_i')^{(5)}\le(\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t)=(p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t)=\ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)},(p_i)^{(5)},\ (r_i)^{(5)}}$ are positive constants and $\boxed{i=28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t)-(a_i'')^{(5)}(T_{29},t)|\le(\hat{k}_{28})^{(5)}|T_{29}-T_{29}'|e^{-(\widehat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t)-(b_i'')^{(5)}\big((G_{31}),t\big)|<(\hat{k}_{28})^{(5)}||(G_{31})-(G_{31})'||e^{-(\widehat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and $(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\widehat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\widehat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\widehat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} \ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\,\big((G_{39}), t\big) = \ (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - \ T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \quad 146A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \quad , \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}(s_{(13)}), s_{(13)}\right)\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{34}(t) = T_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\overline{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \,,\; T_i(0) = T_i^0 \,,\; G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} \,, T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\,\hat{P}_{13}\,)^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\,\hat{P}_{13}\,)^{(1)}}{(\,\hat{M}_{13}\,)^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\,\hat{P}_{13}\,)^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\,\hat{P}_{13}\,)^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for G_{14} , $G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that (175)

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)} t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[\left((\hat{P}_{32})^{(6)} + G_{33}^0 \right) e^{\left(- \frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0} \right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0 \right) e^{\left(- \frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \quad 180$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0 \right) e^{\left(- \frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{-(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\hat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(1)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b''_i)^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\hat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ and $(\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and

respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\le\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)},i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$$G_i\ (t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i\ (t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$$G_{29}\le\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_{0}^{t}\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \qquad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_{+}$. 249

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 250

it results

$$G_i(t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0 e^{(-(b_i')^{(6)}t)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32}<(\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt}\leq\left((\widehat{M}_{32})^{(6)}\right)_1-(a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33}\leq\left((\widehat{M}_{32})^{(6)}\right)_2=G_{33}^0+2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34}\leq\left((\widehat{M}_{32})^{(6)}\right)_3=G_{34}^0+2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}(T_{37}^{(1)}, s_{(36)})|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a_{36}'')^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a_{36}')^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right) \quad (259)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a_{36}'')^{(7)}$ and $(b_{36}'')^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (261)

it results

$$G_i(t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0 e^{\left(-(b_i')^{(7)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: (262)

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36}<(\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt}\leq\left((\widehat{M}_{36})^{(7)}\right)_1-(a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37}\leq\left((\widehat{M}_{36})^{(7)}\right)_2=G_{37}^0+2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38}\leq\left((\widehat{M}_{36})^{(7)}\right)_3=G_{38}^0+2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36}\ is$ bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. (263)

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t))=(b_{37}')^{(7)}$ then $T_{37}\to\infty$. (264)

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{40})^{(8)}|G_{41}^{(1)} - G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)} +$$
$$\int_0^t\{(a_{40}')^{(8)}|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a_{40}'')^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right)|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}|(a_{40}'')^{(8)}\left(T_{41}^{(1)}, s_{(40)}\right) - (a_{40}'')^{(8)}\left(T_{41}^{(2)}, s_{(40)}\right)|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)}t} \leq \frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right) d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right) \quad 273$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\big((G_{43})(t), t\big) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\big((G_{43})(t), t(t), t\big) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \big((\hat{P}_{44})^{(9)} + G_j^0\big)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\big((\hat{Q}_{44})^{(9)} + T_j^0\big)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\Big(\big((G_{47})^{(1)}, (T_{47})^{(1)}\big), \big((G_{47})^{(2)}, (T_{47})^{(2)}\big)\Big) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \big(\widetilde{(G_{47})}, \widetilde{(T_{47})}\big) = \mathcal{A}^{(9)}\big((G_{47}), (T_{47})\big)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a_{44}')^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a_{44}'')^{(9)}\big(T_{45}^{(1)}, s_{(44)}\big)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a_{44}'')^{(9)}\big(T_{45}^{(1)}, s_{(44)}\big) - (a_{44}'')^{(9)}\big(T_{45}^{(2)}, s_{(44)}\big)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b'_i)^{(9)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right)) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t), t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^{0}e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}}G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}}G_{13}^{0}e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)},$ $\boldsymbol{if}$ $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, $\boldsymbol{if}$ $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)},$ 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},$ $\boldsymbol{if}$ $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},$ $\boldsymbol{if}$ $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, $\boldsymbol{if}$ $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},$ $\boldsymbol{if}$ $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$ 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^{0}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G_{20}^{0}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, , (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, ***if*** $(v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, ***if*** $(v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,
and $\boxed{(v_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, ***if*** $(\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big) - (b''_{33})^{(6)}\big((G_{35}),t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}\ (v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}\ (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T^0_{32}}{T^0_{33}}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G^0_{32}e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}}G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}}G^0_{32}e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)}G^0_{32}}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G^0_{34}e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)}G^0_{32}}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G^0_{34}e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0$, $(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0$, $(u_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0$, $(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0$, $(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)}$, ***if*** $(\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}$, $\boldsymbol{if}$ $(\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}$, $\boldsymbol{if}$ $(u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}$, $\boldsymbol{if}$ $(u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)}$,

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}$ $(\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \leq G_{38}(t) \leq \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^{0}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G_{40}^{0}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)}=(v_0)^{(9)},(m_1)^{(9)}=(v_1)^{(9)},$ $\boldsymbol{if}$ $(v_0)^{(9)}<(v_1)^{(9)}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(\bar{v}_1)^{(9)}$, $\boldsymbol{if}$ $(v_1)^{(9)}<(v_0)^{(9)}<(\bar{v}_1)^{(9)}$,

and $\boxed{(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(v_0)^{(9)},$ $\boldsymbol{if}$ $(\bar{v}_1)^{(9)}<(v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)}=(u_0)^{(9)},(\mu_1)^{(9)}=(u_1)^{(9)},$ $\boldsymbol{if}$ $(u_0)^{(9)}<(u_1)^{(9)}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)}<(u_0)^{(9)}<(\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)}=\frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(u_0)^{(9)},\boldsymbol{if}$ $(\bar{u}_1)^{(9)}<(u_0)^{(9)}$ where $(u_1)^{(9)},(\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{44}(t)\le G_{44}^0e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{45}(t)\le\frac{1}{(m_2)^{(9)}}G_{44}^0e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^0e^{-(S_2)^{(9)}t}\le G_{46}(t)\le\right.$$
$$\left.\frac{(a_{46})^{(9)}G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}]+\ G_{46}^0e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le\frac{1}{(\mu_2)^{(9)}}T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq\nu^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \le v^{(3)}(t) \le (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also**

defines $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $v^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\quad \boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq v^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424

Proof : From 99,20,44,22,23,44 we obtain A

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \qquad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\,, \qquad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \qquad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(h) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a_{32}')^{(6)} + (a_{32}'')^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a_{34}')^{(6)} + (a_{34}'')^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a_{36}')^{(7)} + (a_{36}'')^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a_{38}')^{(7)} + (a_{38}'')^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a_{40}')^{(8)} + (a_{40}'')^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a_{42}')^{(8)} + (a_{42}'')^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a_{44}')^{(9)} + (a_{44}'')^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a_{46}')^{(9)} + (a_{46}'')^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b_{13}')^{(1)}(b_{14}')^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b_{13}')^{(1)}(b_{14}'')^{(1)}(G) + (b_{14}')^{(1)}(b_{13}'')^{(1)}(G)\right] + (b_{13}'')^{(1)}(G)(b_{14}'')^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b_{16}')^{(2)}(b_{17}')^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b_{16}')^{(2)}(b_{17}'')^{(2)}(G_{19}) + (b_{17}')^{(2)}(b_{16}'')^{(2)}(G_{19})\right] + (b_{16}'')^{(2)}(G_{19})(b_{17}'')^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504

that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi({G_{39}}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}{}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b'_{13})^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b'_{14})^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b'_{15})^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^{*}) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a'_{16})^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a'_{17})^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^{*}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a'_{18})^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^{*}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ and $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ and $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right)$ 586

G 587

The characteristic equation of this system is

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\Big]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\Big)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\Big]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION NINE

Algorithm Design And Analysis And Approximation Algorithms, Approximation Methods And Cloud Databases , Polynomials And Resource Management Et Al: Forsan Et Haec Olim Meminisse Iuvabit- Frustra Laborant Quotquot Se Calculationibus Fatigant Pro Inventione Quadraturae Circuli Model

INTRODUCTION—VARIABLES USED

Making cloud computing more efficient: for database-driven applications, new software could reduce hardware requirements by 95 percent while actually improving performance. Larry Hardesty, MIT News Office ILLUSTRATION: CHRISTINE DANILOFF/MIT (Please see article for figurative representations) March 12, 2013

For many companies, moving their web-application servers to the cloud is an attractive option, since cloud-computing services can offer economies of scale, extensive technical support and easy accommodation of demand fluctuations.

But for applications that depend heavily on database queries, cloud hosting can pose as many problems as it solves. Cloud services often partition their servers into "virtual machines," each of which gets so many operations per second on a server's central processing unit, so much space in memory, and the like. That makes cloud servers easier to manage, but for database-intensive applications, it can result in the allocation of about 20 times as much hardware as should be necessary. And the cost of that over provisioning gets passed on to customers. MIT researchers are developing a new system called DBSeer that should help solve this problem and others, such as the pricing of cloud services and the diagnosis of application slowdowns. At the recent Biennial Conference on Innovative Data Systems Research, the researchers laid out their vision for DBSeer. And in June, at the annual meeting of the Association for Computing Machinery's Special Interest Group on Management of Data (SIGMOD), they will unveil the algorithms at the heart of DBSeer, which use machine-learning techniques to build accurate models of performance and resource demands of database-driven applications. DBSeer's advantages aren't restricted to cloud computing, either. Teradata, a major database company, has already assigned several of its engineers the task of importing the MIT researchers' new algorithm — which has been released under an open-source license — into its own software.

Virtual limitations

Barzan Mozafari, a postdoc in the lab of professor of electrical engineering and computer science Samuel Madden and lead author on both new papers, explains that, with virtual machines, server resources must be allocated according to an application's peak demand. "You're not going to hit your peak load all the time," Mozafari says. "So that means that these resources are going to be underutilized most of the time."

Moreover, Mozafari says, the provisioning for peak demand is largely guesswork. "It's very counterintuitive," Mozafari says, "but you might take on certain types of extra load that might help your overall performance." Increased demand means that a database server will store more of its frequently used data in its high-speed memory, which can help it process requests more quickly.

On the other hand, a slight increase in demand could cause the system to slow down precipitously — if, for instance, too many requests require modification of the same pieces of data, which need to be updated on multiple servers. "It's extremely nonlinear," Mozafari says.

Mozafari, Madden, postdoc Alekh Jindal, and Carlo Curino, a former member of Madden's group who's now at Microsoft, use two different techniques in the SIGMOD paper to predict how a database-driven application will respond to increased load. Mozafari describes the first as a "black box" approach: DBSeer simply monitors fluctuations in both the number and type of user requests and system performance and uses machine-learning techniques to correlate the two. This approach is good at predicting the consequences of fluctuations that don't fall too far outside the range of the training data.

Gray areas

Often, however, database managers — or prospective cloud-computing customers — will be interested in the consequences of a fourfold, tenfold, or even hundredfold increase in demand. For those types of predictions, Mozafari explains, DBSeer uses a "gray box" model, which takes into account the idiosyncrasies of particular database systems.

For instance, Mozafari explains, updating data stored on a hard drive is time-consuming, so most database servers will try to postpone that operation as long as they can, instead storing data modifications in the much faster — but volatile — main memory. At some point, however, the server has to commit its pending modifications to disk, and the criteria for making that decision can vary from one database system to another.

The version of DBSeer presented at SIGMOD includes a gray-box model of My SQL, one of the most widely used database systems. The researchers are currently building a new model for another popular system, Postgre SQL. Although adapting the model isn't a negligible undertaking, models tailored to just a handful of systems would cover the large majority of database-driven Web applications.

The researchers tested their prediction algorithm against both a set of benchmark data, called TPC-C, that's commonly used in database research and against real-world data on modifications to the Wikipedia database. On average, the model was about 80 percent accurate in predicting CPU use and 99 percent accurate in predicting the bandwidth consumed by disk operations.

"We're really fascinated and thrilled that someone is doing this work," says Doug Brown, a database software architect at Teradata. "We've already taken the code and are prototyping right now." Initially, Brown says, Teradata will use the MIT researchers' prediction algorithm to determine customers' resource requirements. "The really big question for our customers is, 'How are we going to scale?'" Brown says.

Brown hopes, however, that the algorithm will ultimately help allocate server resources on the fly, as database requests come in. If servers can assess the demands imposed by individual requests and budget

accordingly, they can ensure that transaction times stay within the bounds set by customers' service agreements. For instance, "if you have two big, big resource consumers, you can calculate ahead of time that we're only going to run two of these in parallel," Brown says. "There are all kinds of games you can play in workload management."

Optimal Resource Allocation in Clouds Cloud Computing (CLOUD), 2010 IEEE 3rd International Conference on Date of Conference: 5-10 July 2010Author(s): Chang, Fangzhe Ren, Jennifer; Viswanathan, Ramesh: Conference Publications

Cloud platforms enable enterprises to lease computing power in the form of virtual machines. An important problem for such enterprise users is to understand how many and what kinds of virtual machines will be needed from clouds. We formulate demand for computing power and other resources as a resource allocation problem with multiplicity, where computations that have to be performed concurrently are represented as tasks and a later task can reuse resources released by an earlier task. We show that finding a minimized allocation is NP-complete. This paper presents an approximation algorithm with a proof of its approximation bound that can yield close to optimum solutions in polynomial time. Enterprise users can exploit the solution to reduce the leasing cost and amortize the administration overhead (e.g., setting up VPNs or configuring a cluster). Cloud providers may utilize the solution to share their resources among a larger number of users.

(i) Algorithm design and analysis and Approximation algorithms
(ii) Approximation methods and Cloud Databases
(iii) Polynomials and Resource management

NOTATION

Module One

G_{13} : Category one of Algorithm design and analysis

G_{14} : Category two of Algorithm design and analysis

G_{15} : Category three of Algorithm design and analysis

T_{13} : Category one of Approximation algorithms
T_{14} : Category two of Approximation algorithms
T_{15} : Category three of Approximation algorithms

Module Two

G_{16} : Category one of Approximation algorithms
G_{17} : Category two of Approximation algorithms
G_{18} : Category three of Approximation algorithms

T_{16} : Category one of Algorithm design and analysis

T_{17} : Category two of Algorithm design and analysis

T_{18} : Category three of Algorithm design and analysis

Module three

G_{20} : Category one of Approximation methods

G_{21} : Category two of Approximation methods

G_{22} : Category three of Approximation methods

T_{20} : Category one of Cloud Databases

T_{21} : Category two of Cloud Databases

T_{22} : Category three of Cloud Databases

Module four

G_{24} : Category one of Polynomials

G_{25} : Category two of Polynomials

G_{26} : Category three of Polynomials

T_{24} : Category one of Resource management

T_{25} : Category two of Resource management

T_{26} : Category three of Resource management

Module five

G_{28} : Category one of Resource management

G_{29} : Category two of Resource management

G_{30} : Category three of Resource management

T_{28} : Category one of Cloud Databases

T_{29} : Category two of Cloud Databases

T_{30} : Category three of Cloud Databases

Module six

Dense Coding in Experimental Quantum Communication BibTeX or EndNote (RIS):Klaus Mattle1, Harald Weinfurter1, Paul G. Kwiat1,2, and Anton Zeilinger1

Classically, sending more than one bit of information **requires (e)** manipulation of more than one two-state particle. Authors demonstrate experimentally that one can transmit one of three messages, i.e., 1 "trit" ≈1.58bit, by **manipulating** only one of two entangled particles. The increased channel capacity is proven by transmitting ASCII characters in five trits instead of the usual 8 bits.© 1996 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.76.4656 DOI: 10.1103/PhysRevLett.76.4656PACS: 03.65.-w, 89.70.+c

G_{32} : Category one of manipulation of more than one two-state particle.

G_{33} : Category two of manipulation of more than one two-state particle.

G_{34} : Category three of manipulation of more than one two-state particle.

T_{32} : Category one of sending more than one bit of information

T_{33} : Category two of sending more than one bit of information

T_{34} : Category three of sending more than one bit of information

Module seven

G_{36} : Category one of transmission of one of three messages, i.e., 1 "trit" ≈1.58bit,

G_{37} : Category two of transmission of one of three messages, i.e., 1 "trit" ≈1.58bit,

G_{38} : Category three of transmission of one of three messages, i.e., 1 "trit" ≈1.58bit,

T_{36} : Category one of only one of two entangled particles (and there are many such pairs which satisfy the relation and whose characterstics form the background for the assimilation of the classificational procedure)

T_{37} : Category two of only one of two entangled particles (and there are many such pairs which satisfy the relation and whose characterstics form the background for the assimilation of the classificational procedure)

T_{38} : Category three of only one of two entangled particles (and there are many such pairs which satisfy the relation and whose characterstics form the background for the assimilation of the classificational procedure)

Module eight

Quantum Cryptography with Entangled Photons: Thomas Jennewein1, Christoph Simon1, Gregory Weihs1, Harald Weinfurter2, and Anton Zeilinger1

By realizing a quantum cryptography system based on polarization entangled photon pairs **authors establish** highly secure keys, because a single photon source is approximated and the inherent randomness of quantum measurements is exploited. They implement a novel key distribution scheme **using** Wigner's inequality to test the security of the quantum channel, and, alternatively, realize a variant of the BB84 protocol. System has two completely independent users separated by 360 m, and generates raw keys at rates of 400–800 bits/s with bit error rates around 3%.© 2000 The American Physical SocietyURL: http://link.aps.org/doi/10.1103/PhysRevLett.84.4729DOI: 10.1103/PhysRevLett.84.4729 PACS: 03.67.Dd, 42.79.Sz, 89.80.+h

G_{40} : Category one of realisation of quantum cryptography system based on polarization entangled photon pairs

G_{41} : Category two of realisation of quantum cryptography system based on polarization entangled photon pairs

G_{42} : Category three of realisation of quantum cryptography system based on polarization entangled photon pairs

T_{40} : Category one of highly secure keys, because a single photon source is approximated and the inherent randomness of quantum measurements is exploited

T_{41} : Category two of highly secure keys, because a single photon source is approximated and the inherent randomness of quantum measurements is exploited

T_{42} : Category three of highly secure keys, because a single photon source is approximated and the

inherent randomness of quantum measurements is exploited

Module Nine

G_{44} : Category one of Wigner's inequalities to test the security of the quantum channels, and, alternatively, realizes a variant of the BB84 protocol. System has two completely independent users separated by 360 m, and generates raw keys at rates of 400–800 bits/s with bit error rates around 3%.

G_{45} : Category two of Wigner's inequalities to test the security of the quantum channels, and, alternatively, realizes a variant of the BB84 protocol. System has two completely independent users separated by 360 m, and generates raw keys at rates of 400–800 bits/s with bit error rates around 3%.

G_{46} : Category three of Wigner's inequalities to test the security of the quantum channels, and, alternatively, realizes a variant of the BB84 protocol. System has two completely independent users separated by 360 m, and generates raw keys at rates of 400–800 bits/s with bit error rates around 3%.

T_{44} : Category one of key distribution scheme

T_{45} : Category two of key distribution scheme

T_{46} : Category three of key distribution scheme

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad (1)$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad (2)$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad (3)$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad (4)$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad (5)$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad (7)$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad (8)$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad (9)$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad (10)$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad (11)$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad (13)$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad (14)$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad (15)$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad (16)$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad (17)$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad (19)$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad (20)$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad (21)$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad (22)$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad (23)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \qquad (69)$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad (75)$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\,-(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\,-(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\,-(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\,+(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) \quad +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \left[\begin{array}{ccc} (a'_{33})^{(6)} & +(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) \quad +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \left[\begin{array}{ccc} (a'_{34})^{(6)} & +(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) \quad +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{array}\right] G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$

Eighth augmentation coefficients $\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt}$$

$$= (a_{45})^{(9)}G_{44}$$

$$-\begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt}$$

$$= (a_{46})^{(9)}G_{45}$$

$$-\begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad (118)$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\widehat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\widehat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \quad (122)$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}),t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \quad (123)$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: (124)

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}((G_{31}),t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and $(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: (125)

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: (126)

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \quad (127)$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}, (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t) = (p_i)^{(6)}$ 128

$\lim_{G\to\infty}(b_i'')^{(6)}\,((G_{35}),t) = \ (r_i)^{(6)}$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$|(a_i'')^{(6)}(T_{33}',t) - (a_i'')^{(6)}(T_{33},t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$

$|(b_i'')^{(6)}((G_{35})',t) - (b_i'')^{(6)}((G_{35}),t)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$

$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$

Where we suppose

$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$(a_i'')^{(7)}(T_{37},t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$

$(b_i'')^{(7)}(G_{39},t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$

132

$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37},t) = (p_i)^{(7)}$

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36{,}37{,}38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}), t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36{,}37{,}38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40{,}41{,}42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)}$ 140

$\lim_{G\to\infty}(b_i'')^{(8)}\left((G_{43}),t\right) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \leq (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t}$ 142

$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\left((G_{43}),t\right)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46$ 146 A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$

$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)}and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)$, $T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}\left(s_{(13)}\right) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}\left(s_{(13)}\right) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}\left(s_{(16)}\right) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}\left(T_{17}\left(s_{(16)}\right), s_{(16)}\right)\right) G_{16}\left(s_{(16)}\right)\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^{0} + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^{0} + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^{0}) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^{0}\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^{0}}{G_{17}^{0}}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^{0} + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^{0} + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^{0}) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^{0}\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^{0}}{G_{21}^{0}}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^{0} + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^{0} + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_0^t\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\bar{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left(e^{(\bar{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\bar{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left(e^{(\bar{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\bar{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left(e^{(\bar{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\bar{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(P_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_{0}^{t}\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_{+}$.

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a''_{i})^{(1)}$ and $(b''_{i})^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15}. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205
unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)}|G_{21}^{(1)} - G_{21}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)})|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}|(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)}) - (a_{20}'')^{(3)}(T_{21}^{(2)}, s_{(20)})|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad 214$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}), (\widetilde{T_{27}})\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{24})^{(4)}\left|G_{25}^{(1)} - G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)} +$$

$$\int_0^t\{(a_{24}')^{(4)}\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a_{24}'')^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t} \leq \tag{226}$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a_{24}')^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)}|G_{29}^{(1)} - G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)})|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ and $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]} \geq 0$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29}\ \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right),\quad t = log\,\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{32})^{(6)}|G_{33}^{(1)}-G_{33}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)|\;e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$|(G_{35})^{(1)}-(G_{35})^{(2)}|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq$$ 259
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 275

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\ t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations

$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)} v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)} u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}$ $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}$ $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}$ $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le\right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously 331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t} \right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)} t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t} \right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$, $(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}$ $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}$ $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,

and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)},\ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{\left((S_1)^{(7)} - (p_{36})^{(7)}\right)t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\right] + G_{38}^0 e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t}\right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, ***if*** $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, ***if*** $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, ***if*** $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}$ $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\leq\frac{dv^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

$$\text{it follows } (\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\nu_1)^{(1)}$$

In the same manner , we get 384

$$v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq v^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\leq(\bar{\nu}_1)^{(1)}\leq\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\leq v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\leq v^{(1)}(t)\leq(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\leq u^{(1)}(t)\leq(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \ge \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \le \nu^{(5)}(t) \le (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \le \nu^{(5)}(t) \le (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)\le\frac{dv^{(9)}}{dt}\le-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)},(\nu_0)^{(9)}$:-

For $0<\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}<(\nu_1)^{(9)}<(\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t)\ge\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}\quad,\quad\boxed{(C)^{(9)}=\frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(9)}=\frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\bar{\nu}_1)^{(9)}$

If $0<(\nu_1)^{(9)}<(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}<(\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)}\le\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}\le\nu^{(9)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\bar{\nu}_1)^{(9)}$$

If $0<(\nu_1)^{(9)}\le(\bar{\nu}_1)^{(9)}\le\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)}\le\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)}\le\nu^{(9)}(t)\le(m_1)^{(9)},\quad\boxed{\nu^{(9)}(t)=\frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)}\le u^{(9)}(t)\le(\mu_1)^{(9)},\quad\boxed{u^{(9)}(t)=\frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0$ 446

$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0$ 447

$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0$ 448

has a unique positive solution , which is an equilibrium solution

$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0$ 449

$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0$ 450

$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0$ 451

$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0$ 452

$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0$ 453

$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0$ 454

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(i) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^* :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]} \quad , \quad G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$$

$$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]} \quad , \quad T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]} \quad , \quad G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]} \qquad 513$$

$$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]} \quad , \quad T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]} \qquad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]} \quad , \quad G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$$

$$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]} \quad , \quad T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$$G_{24}^{*} = \frac{(a_{24})^{(4)}G_{25}^{*}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^{*})]} \quad , \quad G_{26}^{*} = \frac{(a_{26})^{(4)}G_{25}^{*}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^{*})]}$$

$$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]} \quad , \quad T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]} \qquad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}\quad,\quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}\quad,\quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}\quad,\quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}\quad,\quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof:_Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i \qquad , T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*})=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G^*_{13}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G^*_{14}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G^*_{15}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a_{24}')^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a_{25}')^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a_{26}')^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b_{24}')^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^*\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b_{25}')^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^*\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b_{26}')^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^*\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a_{28}')^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a_{29}')^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a_{30}')^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b_{28}')^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T_{28}^*\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b_{29}')^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T_{29}^*\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b_{30}')^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T_{30}^*\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \ , \ \frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \ , \ \frac{\partial (b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \ , \ \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b''_i)^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left((\lambda)^{(1)}\right)^2+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\Big)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\Big]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)})$$

$$\left[\left(((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$+\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)})$$

$$\left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$+\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T^*_{37} + (b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\} = 0$$

$$+$$

$$(\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)})$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TEN

Delayed-Choice Entanglement Swapping, Peres's Gedanken Experiment, Greenberger–Horne–Zeilinger Entanglement, Physical Interactions And Processes, Quantum Correlations Et Al. : Altissima Quaeque Flumina Minimo Sono Labi-Zukunftsmusik Leitfaden Model

INTRODUCTION—VARIABLES USED

Experimental delayed-choice entanglement swapping Xiao-song Ma, Stefan Zotter, Johannes Kofler, Rupert Ursin, Thomas Jennewein, Časlav Brukner & Anton Zeilinger Nature Physics 8, 479–484 (2012) doi: 10.1038/nphys2294 Received 23 September 2011 Accepted 16 March 2012 Published online 22 April 2012

(1) Motivated by the question of which kind of physical interactions and processes **are (e) needed** for the production of quantum entanglement, Peres has put forward the radical idea of delayed-choice entanglement swapping for **production of** quantum entanglement.
(2) There, entanglement can be 'produced a posteriori**, such a production being done** after the entangled particles have been measured and may no longer exist'. Here, we report the realization of Peres's gedanken experiment.
(3) Using four photons, one can actively **delay (e) the** choice of measurement—implemented through a high-speed tunable bipartite-state analyser and a quantum random-number generator—on two of the photons into the time-like future of the registration of the other two photons.
(4) This effectively projects the two already registered photons **onto(e&eb)** one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'. Nature 403, 515-519 (3 February 2000) | doi:10.1038/35000514

Experimental test of quantum nonlocality in three-photon Greenberger–Horne–Zeilinger entanglement Jian-Wei Pan1, Dik Bouwmeester2, Matthew Daniell1, Harald Weinfurter3 & Anton Zeilinger1

(5) Bell's theorem states that certain statistical correlations predicted by quantum physics for measurements on two-particle systems cannot be understood within a realistic picture **based on** local properties of each individual particle—even if the two particles are separated by large distances.
(6) Einstein, Podolsky and Rosen first recognized the fundamental significance of these quantum correlations (termed 'entanglement' by Schrödinger3) and the two-particle quantum predictions have found ever-increasing experimental support4. A more striking conflict between quantum mechanical **and (e)** local realistic predictions (for perfect correlations) has been discovered,
(7) but experimental verification has been difficult, as it requires entanglement between at least three particles. Here authors report experimental confirmation of this conflict, **using their (e) recently developed** method to observe three-photon entanglement, or 'Greenberger–Horne–Zeilinger' (GHZ) states.
(8) The results of three specific experiments, involving measurements of polarization correlations between three photons, **lead to predictions** for a fourth experiment; quantum physical predictions are mutually **contradictory (e)** with expectations based on local realism. Authors find the results of the fourth experiment to be in agreement with the quantum prediction and in striking conflict with local realism.

NOTATION

Module One

G_{13} : Category one of delayed-choice entanglement swapping

G_{14} : Category two of delayed-choice entanglement swapping

G_{15} : Category three of delayed-choice entanglement swapping

T_{13} : Category one of quantum entanglement.
T_{14} : Category two of quantum entanglement.
T_{15} : Category three of quantum entanglement.

Module Two

There, entanglement can be 'produced a posteriori**, such a production being done** after the entangled particles have been measured and may no longer exist'. Here, we report the realization of Peres's gedanken experiment

G_{16} : Category one of entanglement 'production a posteriori

G_{17} : Category two of entanglement 'production a posteriori

G_{18} : Category three of entanglement 'production a posteriori

T_{16} : Category one of entangled particles that have been measured and may no longer exist'.

T_{17} : Category two of entangled particles that have been measured and may no longer exist'

T_{18} : Category three of entangled particles that have been measured and may no longer exist'

Module three

G_{20} : Category one of CHOICE OF MEASUREMENT

G_{21} : Category two of CHOICE OF MEASUREMENT

G_{22} : Category three of CHOICE OF MEASUREMENT

T_{20} : Category one of four photons

T_{21} : Category two of four photons

T_{22} : Category three of four photons

Module four

G_{24} : Category one of Projection of the two already registered photons

G_{25} : Category two of Projection of the two already registered photons

G_{26} : Category three of Projection of the two already registered photons

T_{24} : Category one of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'

T_{25} : Category two of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as

'quantum steering into the past'

T_{26} : Category three of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'

Module five(Case two)

G_{28} : Category one of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'

G_{29} : Category two of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'

G_{30} : Category three of one of two mutually exclusive quantum states in which the photons are either entangled (quantum correlations) or separable (classical correlations). This can also be viewed as 'quantum steering into the past'

T_{28} : Category one of Projection of the two already registered photons

T_{29} : Category two of Projection of the two already registered photons

T_{30} : Category three of Projection of the two already registered photons

Module six

G_{32} : Category one of local properties of each individual particle—even if the two particles are separated by large distances.

G_{33} : Category two of local properties of each individual particle—even if the two particles are separated by large distances.

G_{34} : Category three of local properties of each individual particle—even if the two particles are separated by large distances.

T_{32} : Category one of Bell's theorem states that certain statistical correlations predicted by quantum physics for measurements on two-particle systems cannot be understood within a realistic picture

T_{33} : Category two of Bell's theorem states that certain statistical correlations predicted by quantum physics for measurements on two-particle systems cannot be understood within a realistic picture

T_{34} : Category three of Bell's theorem states that certain statistical correlations predicted by quantum physics for measurements on two-particle systems cannot be understood within a realistic picture

Module seven

G_{36} : Category one of local realistic predictions (for perfect correlations) Einstein, Podolsky and Rosen first recognized the fundamental significance of these quantum correlations (termed 'entanglement' by Schrödinger) and the two-particle quantum predictions have found ever-increasing experimental support

G_{37} : Category two of local realistic predictions (for perfect correlations) Einstein, Podolsky and Rosen first recognized the fundamental significance of these quantum correlations (termed 'entanglement' by

Schrödinger) and the two-particle quantum predictions have found ever-increasing experimental support

G_{38} : Category three of local realistic predictions (for perfect correlations) Einstein, Podolsky and Rosen first recognized the fundamental significance of these quantum correlations (termed 'entanglement' by Schrödinger) and the two-particle quantum predictions have found ever-increasing experimental support

T_{36} : Category one of quantum mechanics (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{37} : Category two of quantum mechanics

T_{38} : Category three of quantum mechanics

Module eight

G_{40} : Category one of three-photon entanglement, or 'Greenberger–Horne–Zeilinger' (GHZ) states.

G_{41} : Category two of three-photon entanglement, or 'Greenberger–Horne–Zeilinger' (GHZ) states.

G_{42} : Category three of three-photon entanglement, or 'Greenberger–Horne–Zeilinger' (GHZ) states.

T_{40} : Category one of conflict between quantum mechanical **and (e)** local realistic predictions (for perfect correlations)

T_{41} : Category two of conflict between quantum mechanical **and (e)** local realistic predictions (for perfect correlations)

T_{42} : Category three of conflict between quantum mechanical **and (e)** local realistic predictions (for perfect correlations)

. Module Nine

G_{44} : Category one of expectations based on local realism

G_{45} : Category two of expectations based on local realism

G_{46} : Category three of expectations based on local realism

T_{44} : Category one of quantum physical predictions (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{45} : Category two of quantum physical predictions

T_{46} : Category three of quantum physical predictions

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}), t\right)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}), t\right)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}), t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\left((G_{19}), t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$
$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor
$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24} \quad 22$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25} \quad 23$$
$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26} \quad 24$$
$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**
$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$
$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$
$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$
$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for

category 1, 2 and 3
$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)}+(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3
$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3
$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)}-(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt}=(b_{17})^{(2)}T_{16}-\begin{bmatrix}(b'_{17})^{(2)}-(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t)\\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t)\end{bmatrix}T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right]\left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right]\left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right]\left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right]\left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right]\left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right]\left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right]\left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right]\left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right]\left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right]\left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right]\left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right]\left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right]\left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right]\left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right]\left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right]\left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right]\left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right]\left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad 78$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)} & \boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \qquad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)} & \boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \qquad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)} & \boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} \quad \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \qquad 81$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh

augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)} G_{32} - \begin{bmatrix} (a'_{33})^{(6)} \boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)} G_{33} - \begin{bmatrix} (a'_{34})^{(6)} \boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)} T_{33} - \begin{bmatrix} (b'_{32})^{(6)} \boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)} T_{32} - \begin{bmatrix} (b'_{33})^{(6)} \boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for

category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)} & +(a''_{40})^{(8)}(T_{41},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \begin{bmatrix} (a'_{44})^{(9)} \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \begin{bmatrix} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \begin{bmatrix} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \begin{bmatrix} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right]T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\leq(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\big((G_{19}),t\big)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)}=1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}},\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}<1$$

Definition of $(\hat{P}_{13})^{(2)},(\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},(\hat{A}_{16})^{(2)}and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)},(a_i')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(p_i)^{(2)},\ (r_i)^{(2)},i=16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)}+(a_i')^{(2)}+(\hat{A}_{16})^{(2)}+(\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)}+(b_i')^{(2)}+(\hat{B}_{16})^{(2)}+(\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 111$$

Where we suppose

$$(a_i)^{(3)},(a_i')^{(3)},(a_i'')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(b_i'')^{(3)}>0,\quad i,j=20,21,22 \quad 112$$

The functions $(a_i'')^{(3)},(b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)},\ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t)\leq(p_i)^{(3)}\leq(\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t)\leq\ (r_i)^{(3)}\leq(b_i')^{(3)}\leq(\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t)=(p_i)^{(3)} \quad 113$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t)=\ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)},(p_i)^{(3)},\ (r_i)^{(3)}}$ are positive constants and $\boxed{i=20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\le(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\le(p_i)^{(4)}\le(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\le\ (r_i)^{(4)}\le(b_i')^{(4)}\le(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\le(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It

is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)}=1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}},\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}<1$$

Definition of $(\hat{P}_{24})^{(4)},(\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)},(\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)},(a_i')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}, i=24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)}+(a_i')^{(4)}+(\hat{A}_{24})^{(4)}+(\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}]<1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)}+(b_i')^{(4)}+(\hat{B}_{24})^{(4)}+(\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}]<1$$

Where we suppose

$$(a_i)^{(5)},(a_i')^{(5)},(a_i'')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(b_i'')^{(5)}>0,\quad i,j=28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)},(b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)},\ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t)\le(p_i)^{(5)}\le(\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big)\le\ (r_i)^{(5)}\le(b_i')^{(5)}\le(\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t)=(p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t)=\ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)},(p_i)^{(5)},\ (r_i)^{(5)}}$ are positive constants and $\boxed{i=28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t)-(a_i'')^{(5)}(T_{29},t)|\le(\hat{k}_{28})^{(5)}|T_{29}-T_{29}'|e^{-(\widehat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t)-(b_i'')^{(5)}\big((G_{31}),t\big)|<(\hat{k}_{28})^{(5)}||(G_{31})-(G_{31})'||e^{-(\widehat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \leq (p_i)^{(6)} \leq (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \leq (r_i)^{(6)} \leq (b_i')^{(6)} \leq (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \leq (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\widehat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\widehat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\widehat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}, \ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le \ (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad 132$$

$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37}, t) = (p_i)^{(7)}$

$\lim_{G\to\infty}(b_i'')^{(7)}\,\big((G_{39}), t\big) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - \ T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\big[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\big]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, (r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \quad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \quad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^{0} + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^{0} + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^{0} + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^{0} + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^{0} + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\overline{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \,,\; T_i(0) = T_i^0 \,,\; G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} \,, T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\,\hat{P}_{13}\,)^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\,\hat{P}_{13}\,)^{(1)}}{(\,\hat{M}_{13}\,)^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\,\hat{P}_{13}\,)^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\,\hat{P}_{13}\,)^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for G_{14} , $G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that (175)

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)} t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[\left((\hat{P}_{32})^{(6)} + G_{33}^0 \right) e^{\left(- \frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0} \right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0 \right) e^{\left(- \frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \quad 180$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0 \right) e^{\left(- \frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{-(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\hat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(1)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b''_i)^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\hat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ and $(\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and

respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}),(\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}),(\widetilde{T_{27}})\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\le\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)},i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$$G_i\ (t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i\ (t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$$G_{29}\le\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right)$, $t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_{0}^{t}\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \qquad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}\ and\ (\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_{+}$. 249

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 250

it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{(-(b_i')^{(6)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32}<(\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt}\leq\left((\widehat{M}_{32})^{(6)}\right)_1-(a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33}\leq\left((\widehat{M}_{32})^{(6)}\right)_2=G_{33}^0+2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34}\leq\left((\widehat{M}_{32})^{(6)}\right)_3=G_{34}^0+2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If $G_{32}\ \ is$ bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}(T_{37}^{(1)}, s_{(36)})|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a_{36}'')^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a_{36}')^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right) \quad 259$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a_{36}'')^{(7)}$ and $(b_{36}'')^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 261

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36}<(\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt}\leq\left((\widehat{M}_{36})^{(7)}\right)_1-(a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37}\leq\left((\widehat{M}_{36})^{(7)}\right)_2=G_{37}^0+2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38}\leq\left((\widehat{M}_{36})^{(7)}\right)_3=G_{38}^0+2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36}\ \ is$ bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}\ ((G_{39})(t),t))=(b_{37}')^{(7)}$ then $T_{37}\ \to\infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{40})^{(8)}|G_{41}^{(1)} - G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)} +$$
$$\int_0^t\{(a_{40}')^{(8)}|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a_{40}'')^{(8)}(T_{41}^{(1)}, s_{(40)})|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}|(a_{40}'')^{(8)}(T_{41}^{(1)}, s_{(40)}) - (a_{40}'')^{(8)}(T_{41}^{(2)}, s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)}t} \leq \frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right) d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right) \quad 273$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\big((G_{43})(t), t\big) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\big((G_{43})(t), t(t), t\big) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \big((\hat{P}_{44})^{(9)} + G_j^0\big)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\big((\hat{Q}_{44})^{(9)} + T_j^0\big)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\Big(\big((G_{47})^{(1)}, (T_{47})^{(1)}\big), \big((G_{47})^{(2)}, (T_{47})^{(2)}\big)\Big) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\big|G_i^{(1)}(t) - G_i^{(2)}(t)\big|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\big|T_i^{(1)}(t) - T_i^{(2)}(t)\big|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \big(\widetilde{(G_{47})}, \widetilde{(T_{47})}\big) = \mathcal{A}^{(9)}\big((G_{47}), (T_{47})\big)$
It results

$$\big|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\big| \leq \int_0^t (a_{44})^{(9)}\big|G_{45}^{(1)} - G_{45}^{(2)}\big|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a_{44}')^{(9)}\big|G_{44}^{(1)} - G_{44}^{(2)}\big|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a_{44}'')^{(9)}\big(T_{45}^{(1)}, s_{(44)}\big)\big|G_{44}^{(1)} - G_{44}^{(2)}\big|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\big|(a_{44}'')^{(9)}\big(T_{45}^{(1)}, s_{(44)}\big) - (a_{44}'')^{(9)}\big(T_{45}^{(2)}, s_{(44)}\big)\big|\; e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\big|(G_{47})^{(1)} - G^{(2)}\big|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b'_i)^{(9)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right)) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t), t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^{0}e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}}G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}}G_{13}^{0}e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)},,(\bar{v}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0$, $(\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)},$ $\boldsymbol{if}$ $(v_0)^{(2)} < (v_1)^{(2)}$ 305

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}$, $\boldsymbol{if}$ $(v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$ 306

and $\boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)},$ $\boldsymbol{if}$ $(\bar{v}_1)^{(2)} < (v_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},$ $\boldsymbol{if}$ $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, $\boldsymbol{if}$ $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},$ $\boldsymbol{if}$ $(\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$ 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^{0}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G_{20}^{0}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, , (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}$, ***if*** $(\nu_0)^{(4)} < (\nu_1)^{(4)}$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, ***if*** $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,
and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, ***if*** $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, ***if*** $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, ***if*** $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,
and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}$, ***if*** $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, ***if*** $(v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, ***if*** $(v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,

and $\boxed{(v_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, ***if*** $(\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big) - (b''_{33})^{(6)}\big((G_{35}),t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}\ (v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}\ (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T^0_{32}}{T^0_{33}}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G^0_{32}e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}}G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}}G^0_{32}e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)}G^0_{32}}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G^0_{34}e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)}G^0_{32}}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G^0_{34}e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t}-e^{-(b'_{34})^{(6)}t}\right]+T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0$, $(\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0$, $(\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}$, ***if*** $(\nu_0)^{(7)} < (\nu_1)^{(7)}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}$, $\boldsymbol{if}$ $(\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}$, $\boldsymbol{if}$ $(u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}$, $\boldsymbol{if}$ $(u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)}$,

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}$ $(\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \leq G_{38}(t) \leq \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^{0}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G_{40}^{0}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)}=(v_0)^{(9)},(m_1)^{(9)}=(v_1)^{(9)},$ $\boldsymbol{if}$ $(v_0)^{(9)}<(v_1)^{(9)}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(\bar{v}_1)^{(9)}$, $\boldsymbol{if}$ $(v_1)^{(9)}<(v_0)^{(9)}<(\bar{v}_1)^{(9)}$,

and $\boxed{(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(v_0)^{(9)},$ $\boldsymbol{if}$ $(\bar{v}_1)^{(9)}<(v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)}=(u_0)^{(9)},(\mu_1)^{(9)}=(u_1)^{(9)},$ $\boldsymbol{if}$ $(u_0)^{(9)}<(u_1)^{(9)}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)}<(u_0)^{(9)}<(\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)}=\frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(u_0)^{(9)},\boldsymbol{if}$ $(\bar{u}_1)^{(9)}<(u_0)^{(9)}$ where $(u_1)^{(9)},(\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{44}(t)\le G_{44}^0e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{45}(t)\le\frac{1}{(m_2)^{(9)}}G_{44}^0e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^0e^{-(S_2)^{(9)}t}\le G_{46}(t)\le\right.$$
$$\left.\frac{(a_{46})^{(9)}G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}]+\ G_{46}^0e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le\frac{1}{(\mu_2)^{(9)}}T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq\nu^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \le v^{(3)}(t) \le (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also**

defines $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le \nu^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le \nu^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le \nu^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $v^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\quad \boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq v^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \qquad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\,, \qquad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \qquad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(j) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504

that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi({G_{39}}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}{}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b'_{13})^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b'_{14})^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b'_{15})^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^{*}) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a'_{16})^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a'_{17})^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^{*}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a'_{18})^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^{*}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}\mathrm{T}^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}\mathrm{T}^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}\mathrm{T}^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right)$ 586

G 587

The characteristic equation of this system is

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION ELEVEN

Loophole-Free Bell Inequality, Local Realism, Universal Wave Function, Quantum And Classical Bell-Type Inequalities, Universal Wave Function And Approaches To Quantum Description Of Macroscopic World, With Emphasis On Dissipation, Etal. : Abundans Cautela Non Nocet -Abyssus Abyssum Invocate- Gemutlichkeit-Ansatz Model

INTRODUCTION—VARIABLES USED

Proposal for a loophole-free Bell inequality experiment: Paul G. Kwiat, Philippe H. Eberhard, Aephraim M. Steinberg, and Raymond Y. Chiao

(1) The nonlocality inherent in quantum mechanics has been tested experimentally, but the nonlocal interpretation of all the relevant experiments can be challenged. All past tests of Bell's inequalities **have (e) required** supplementary assumptions, without which the experimental results satisfy the inequalities, in agreement with the notion of local realism.

(2) There are basically three loopholes that need to be closed; neither the cascade sources nor the parametric down-conversion sources of correlation photons used to date are capable of closing all of them, even if the detectors used had been 100% efficient. Authors propose a two-crystal down-conversion source, relying on type-II collinear phase matching, **which should (eb) permit a** violation of Bell's inequalities without the need for supplementary assumptions. As the source can produce a true singlet like state, it is also relevant for quantum cryptographic applications.

Quantum/classical correspondence in the light of Bell's inequalities: Leonid A. Khalfin, Boris S. Tsirelson

(3) Instead of the usual asymptotic passage from quantum mechanics to classical mechanics when a parameter tended to infinity, a sharp boundary is obtained for the domain of existence of classical reality. Classical mechanics is treated as separable empirical reality following d'Espagnat, **described (eb) by a** mathematical superstructure over quantum dynamics for the universal wave function.

(4) Being empirical, this reality is **constructed (eb) in terms** of both fundamental notions and characteristics of observers. It is presupposed that considered observers perception of the world **as a** system of collective degrees of freedom that are inherently dissipative **because of**

interaction with thermal degrees of freedom.

(5) Relevant problems of foundation of statistical physics are considered. A feasible example is given of a macroscopic system not admitting classical reality (problems of foundation of statistical physics).

(i) quantum and classical Bell-type inequalities;

(ii) universal wave function and approaches to quantum description of macroscopic world, with emphasis on dissipation;

(iii) Spontaneous reduction models and experimental tests of the universal validity of the quantum theory.

The Einstein-Podolsky-Rosen Argument and the Bell Inequalities Laszlo E. Szabo

(6) In 1935 Einstein, Podolsky, and Rosen (EPR) published an important paper in which they claimed that the whole formalism of quantum mechanics together with what they called "Reality Criterion" imply that quantum mechanics cannot be complete. That is, there must exist some elements of reality that are not described by quantum mechanics. There must be, they concluded, a more complete description of physical reality behind quantum mechanics. There must be a state, a hidden variable, characterizing the state of affairs in the world in more details than the quantum mechanical state, something that also reflects the missing elements of reality. From this we (authors of the main paper) derive that perception is not reality, but for both to be equal it is necessary to add or subtract certain variables, which we have called "cosmic consciousness" which determines the state of reality. Perception **(e&eb)** Reality. Both the positive and negative cases like augmented reality and dissipated reality could be considered.

NOTATION

Module One

G_{13} : Category one of supplementary assumptions, without which the experimental results satisfy the inequalities, in agreement with the notion of local realism

G_{14} : Category two of supplementary assumptions, without which the experimental results satisfy the inequalities, in agreement with the notion of local realism

G_{15} : Category three of supplementary assumptions, without which the experimental results satisfy the inequalities, in agreement with the notion of local realism

T_{13} : Category one of nonlocality inherent in quantum mechanics has been tested experimentally, but the nonlocal interpretation of all the relevant experiments can be challenged , past tests of Bell's inequalities

T_{14} : Category two of nonlocality inherent in quantum mechanics has been tested experimentally, but the nonlocal interpretation of all the relevant experiments can be challenged , past tests of Bell's inequalities

T_{15} : Category three of nonlocality inherent in quantum mechanics has been tested experimentally, but the nonlocal interpretation of all the relevant experiments can be challenged , past tests of Bell's inequalities

Module Two

G_{16} : Category one of two-crystal down-conversion source, relying on type-II collinear phase matching

G_{17} : Category two of two-crystal down-conversion source, relying on type-II collinear phase matching

G_{18} : Category three of two-crystal down-conversion source, relying on type-II collinear phase matching

T_{16} : Category one of violation of Bell's inequalities without the need for supplementary assumptions. As the source can produce a true singlet like state, it is also relevant for quantum cryptographic applications

T_{17} : Category two of violation of Bell's inequalities without the need for supplementary assumptions. As the source can produce a true singlet like state, it is also relevant for quantum cryptographic applications

T_{18} : Category three of violation of Bell's inequalities without the need for supplementary assumptions. As the source can produce a true singlet like state, it is also relevant for quantum cryptographic applications

Module three

G_{20} : Category one of Classical mechanics as separable empirical reality following d'Espagnat

G_{21} : Category two of Classical mechanics as separable empirical reality following d'Espagnat

G_{22} : Category three of Classical mechanics as separable empirical reality following d'Espagnat

T_{20} : Category one of mathematical superstructure over quantum dynamics for the universal wave function

T_{21} : Category two of mathematical superstructure over quantum dynamics for the universal wave function

T_{22} : Category three of mathematical superstructure over quantum dynamics for the universal wave function

Module four

G_{24} : Category one of both fundamental notions and characteristics of observers

G_{25} : Category two of both fundamental notions and characteristics of observers

G_{26} : Category three of both fundamental notions and characteristics of observers

T_{24} : Category one of empirical reality

T_{25} : Category two of empirical reality

T_{26} : Category three of empirical reality

Module five

G_{28} : Category one of interaction with thermal degrees of freedom.

G_{29} : Category two of interaction with thermal degrees of freedom.

G_{30} : Category three of interaction with thermal degrees of freedom.

T_{28} : Category one of considered observers perception of the world **as a** system of collective degrees of freedom that are inherently dissipative

T_{29} : Category two of considered observers perception of the world **as a** system of collective degrees of freedom that are inherently dissipative

T_{30} : Category three of considered observers perception of the world **as a** system of collective degrees

of freedom that are inherently dissipative

Module six

G_{32} : Category one of quantum and classical Bell-type inequalities

G_{33} : Category two of quantum and classical Bell-type inequalities

G_{34} : Category three of quantum and classical Bell-type inequalities

T_{32} : Category one of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

T_{33} : Category two of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

T_{34} : Category three of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

Module seven

G_{36} : Category one of universal wave function and approaches to quantum description of macroscopic world, with emphasis on dissipation;

G_{37} : Category two of universal wave function and approaches to quantum description of macroscopic world, with emphasis on dissipation;

G_{38} : Category three of universal wave function and approaches to quantum description of macroscopic world, with emphasis on dissipation;

T_{36} : Category one of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

T_{37} : Category two of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

T_{38} : Category three of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

Module eight

G_{40} : Category one of Spontaneous reduction models and experimental tests of the universal validity of the quantum theory.

G_{41} : Category two of Spontaneous reduction models and experimental tests of the universal validity of the quantum theory.

G_{42} : Category three of Spontaneous reduction models and experimental tests of the universal validity of the quantum theory.

T_{40} : Category one of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

T_{41} : Category two of macroscopic system not admitting classical reality (problems of foundation of

statistical physics).

T_{42} : Category three of macroscopic system not admitting classical reality (problems of foundation of statistical physics).

Module Nine(augmented case)

G_{44} : Category one of Perception

G_{45} : Category two of Perception

G_{46} : Category three of Perception

T_{44} : Category one of Reality

T_{45} : Category two of Reality

T_{46} : Category three of Reality

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad 3$$

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad 4$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad 5$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\big((G_{47}),t\big)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\big((G_{47}),t\big)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\big((G_{47}),t\big)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\big((G_{47}),t\big) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & +(a''_{15})^{(1)}(T_{14},t) & +(a''_{18})^{(2,2,)}(T_{17},t) & +(a''_{22})^{(3,3,)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,)}(T_{33},t) \\ +(a''_{38})^{(7,7)}(T_{37},t) & +(a''_{42})^{(8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15} \quad 57$$

Where $(a''_{13})^{(1)}(T_{14},t)$, $(a''_{14})^{(1)}(T_{14},t)$, $(a''_{15})^{(1)}(T_{14},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,)}(T_{17},t)$, $+(a''_{17})^{(2,2,)}(T_{17},t)$, $+(a''_{18})^{(2,2,)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,)}(T_{21},t)$, $+(a''_{21})^{(3,3,)}(T_{21},t)$, $+(a''_{22})^{(3,3,)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)} & -(b''_{13})^{(1)}(G,t) & -(b''_{16})^{(2,2,)}(G_{19},t) & -(b''_{20})^{(3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,)}(G_{35},t) \\ -(b''_{36})^{(7,7,)}(G_{39},t) & -(b''_{40})^{(8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[\begin{array}{ccc} (b'_{14})^{(1)} & -(b''_{14})^{(1)}(G,t) & -(b''_{17})^{(2,2,)}(G_{19},t) \\ -(b''_{21})^{(3,3,)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,)}(G_{35},t) & -(b''_{37})^{(7,7,)}(G_{39},t) & -(b''_{41})^{(8,8)}(G_{43},t) \\ -(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[\begin{array}{ccc} (b'_{15})^{(1)} & -(b''_{15})^{(1)}(G,t) & -(b''_{18})^{(2,2,)}(G_{19},t) \\ -(b''_{22})^{(3,3,)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,)}(G_{35},t) & -(b''_{38})^{(7,7,)}(G_{39},t) & -(b''_{42})^{(8,8)}(G_{43},t) \\ -(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array}\right] T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) \\ +(a''_{20})^{(3,3,3)}(T_{21},t) & +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) \\ +(a''_{44})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) \\ +(a''_{21})^{(3,3,3)}(T_{21},t) & +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) \\ +(a''_{45})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) \\ +(a''_{22})^{(3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) \\ +(a''_{46})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{18} \quad 63$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\,-(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\,-(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\,-(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)} & +(a''_{21})^{(3)}(T_{21},t) & +(a''_{17})^{(2,2,2)}(T_{17},t) & +(a''_{14})^{(1,1,1,)}(T_{14},t) \\ +(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{37})^{(7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)} & +(a''_{22})^{(3)}(T_{21},t) & +(a''_{18})^{(2,2,2)}(T_{17},t) & +(a''_{15})^{(1,1,1,)}(T_{14},t) \\ +(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t) & \\ +(a''_{38})^{(7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9)}(T_{45},t) & \end{bmatrix} G_{22} \quad 69$$

$+(a''_{20})^{(3)}(T_{21},t)$, $+(a''_{21})^{(3)}(T_{21},t)$, $+(a''_{22})^{(3)}(T_{21},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2)}(T_{17},t)$ are second augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,)}(T_{14},t)$ are third augmentation coefficients for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3

$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3

$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)} & -(b''_{20})^{(3)}(G_{23},t) & -(b''_{16})^{(2,2,2)}(G_{19},t) & -(b''_{13})^{(1,1,1,)}(G,t) \\ -(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{36})^{(7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)} & -(b''_{21})^{(3)}(G_{23},t) & -(b''_{17})^{(2,2,2)}(G_{19},t) & -(b''_{14})^{(1,1,1,)}(G,t) \\ -(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{37})^{(7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\left[-(b''_{24})^{(4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\left[-(b''_{25})^{(4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\left[-(b''_{26})^{(4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\left[+(a''_{28})^{(5)}(T_{29},t)\right] & \left[+(a''_{24})^{(4,4,)}(T_{25},t)\right] & \left[+(a''_{32})^{(6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)\right] & \left[+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)\right] & \left[+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)\right] & \left[+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)\right] & \left[+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)\right] \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$

Eighth augmentation coefficients $\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)} G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) \\ +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & & \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)} T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)} T_{41} - \left[\begin{array}{ccc} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}-(b''_{44})^{(9)}(G_{47},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{13}$$
$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}-(b''_{45})^{(9)}(G_{47},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}-(b''_{46})^{(9)}(G_{47},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t) \\ -(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t) \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad (118)$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\widehat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \quad (122)$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}),t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \quad (123)$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: (124)

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}((G_{31}),t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: (125)

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: (126)

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \quad (127)$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\,\big((G_{35}),t\big)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32{,}33{,}34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\big((G_{35}),t\big)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)},(\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}<1$$

Definition of $(\hat{P}_{32})^{(6)},(\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)},(\hat{A}_{32})^{(6)}and\,(\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)},(a_i')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(p_i)^{(6)},\ (r_i)^{(6)},i=32{,}33{,}34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)}+(a_i')^{(6)}+\ (\hat{A}_{32})^{(6)}+\ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}]<1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)}+(b_i')^{(6)}+\ (\hat{B}_{32})^{(6)}+\ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}]<1$$

Where we suppose

$$(a_i)^{(7)},(a_i')^{(7)},(a_i'')^{(7)},(b_i)^{(7)},(b_i')^{(7)},(b_i'')^{(7)}>0,\qquad i,j=36{,}37{,}38 \qquad 131$$

The functions $(a_i'')^{(7)},(b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)},\ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t)\le(p_i)^{(7)}\le(\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t)\le\ (r_i)^{(7)}\le(b_i')^{(7)}\le(\hat{B}_{36})^{(7)}$$

132

$$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37},t)=(p_i)^{(7)}$$

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36{,}37{,}38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36{,}37{,}38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40{,}41{,}42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)}$ 140

$\lim_{G\to\infty}(b_i'')^{(8)}\left((G_{43}),t\right) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \le (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t}$ 142

$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\left((G_{43}),t\right)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}$, $(r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46$ 146 A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \le (p_i)^{(9)} \le (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \le (r_i)^{(9)} \le (b_i')^{(9)} \le (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$

$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)}and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{G_i(0)=G_i^0>0}$

$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}$, $\boxed{T_i(0)=T_i^0>0}$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^{0} + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^{0} + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^{0} + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^{0} + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^{0} + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^{0} + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^{0} + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^{0} + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166
Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A
which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^{0} + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^{0} + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^{0}) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^{0}\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^{0}}{G_{17}^{0}}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^{0} + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^{0} + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^{0}) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}} \left[\left((\hat{P}_{20})^{(3)} + G_{21}^{0}\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^{0}}{G_{21}^{0}}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)} \left(G_{25}^{0} + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^{0} + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}} \left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_0^t\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\bar{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left(e^{(\bar{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\bar{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left(e^{(\bar{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\bar{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left(e^{(\bar{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\bar{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_{0}^{t}\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_{+}$.

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a''_{i})^{(1)}$ and $(b''_{i})^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15}. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(\left((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)}\right)\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205
unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)}|G_{21}^{(1)} - G_{21}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}|(a_{20}'')^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a_{20}'')^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)|\; e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad 214$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})}, \widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})}, \widetilde{(T_{27})}\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{24})^{(4)}\left|G_{25}^{(1)} - G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)} +$$

$$\int_0^t\{(a_{24}')^{(4)}\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a_{24}'')^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t} \leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a_{24}')^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)}|G_{29}^{(1)} - G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)})|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)}G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29}\ \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{32})^{(6)}|G_{33}^{(1)}-G_{33}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$|(G_{35})^{(1)}-(G_{35})^{(2)}|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If $G_{32} \; is$ bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq$$ 259
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 275

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\ t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})},\widetilde{(T_{47})}:\left(\widetilde{(G_{47})},\widetilde{(T_{47})}\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations

$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)} v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)} u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}$ $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}$ $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}$ $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le\right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously 331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t} \right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)} t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t} \right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}$,, $(\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, $\boldsymbol{if}\ (\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, $\boldsymbol{if}\ (\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}$ $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}$ $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,

and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)},\ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{\left((S_1)^{(7)} - (p_{36})^{(7)}\right)t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\right] + G_{38}^0 e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t}\right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, ***if*** $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, ***if*** $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,

and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, ***if*** $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}$ $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\leq\frac{dv^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

$$\text{it follows } (\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\nu_1)^{(1)}$$

In the same manner , we get 384

$$v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq v^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\leq(\bar{\nu}_1)^{(1)}\leq\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\leq v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\leq v^{(1)}(t)\leq(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\leq u^{(1)}(t)\leq(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $v^{(4)}$:- $\boxed{v^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)}\right) \leq \frac{dv^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_4)^{(4)}v^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)}, (v_0)^{(4)}$:-

For $0 < \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (v_1)^{(4)} < (\bar{v}_1)^{(4)}$

$$v^{(4)}(t) \geq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(v_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (v_2)^{(4)}}}$$

it follows $(v_0)^{(4)} \leq v^{(4)}(t) \leq (v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{v}_1)^{(4)} - (v_0)^{(4)}}{(v_0)^{(4)} - (\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)} \leq v^{(4)}(t) \leq (\bar{v}_1)^{(4)}$

If $0 < (v_1)^{(4)} < (v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)} \leq \frac{(v_1)^{(4)} + (C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)} - (v_2)^{(4)}\right)t\right]}} \leq v^{(4)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (\bar{v}_1)^{(4)}$$

If $0 < (v_1)^{(4)} \leq (\bar{v}_1)^{(4)} \leq \boxed{(v_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)} \leq v^{(4)}(t) \leq \frac{(\bar{v}_1)^{(4)} + (\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)} - (\bar{v}_2)^{(4)}\right)t\right]}} \leq (v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)} \leq v^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{v^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \ge \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \le \nu^{(5)}(t) \le (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \le \nu^{(5)}(t) \le (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}$, $\boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}$, $\boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)\le\frac{dv^{(9)}}{dt}\le-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)},(\nu_0)^{(9)}$:-

For $0<\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}<(\nu_1)^{(9)}<(\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t)\ge\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}\quad,\quad\boxed{(C)^{(9)}=\frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(9)}=\frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)}\le\nu^{(9)}(t)\le(\bar{\nu}_1)^{(9)}$

If $0<(\nu_1)^{(9)}<(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}<(\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)}\le\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}\le\nu^{(9)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\bar{\nu}_1)^{(9)}$$

If $0<(\nu_1)^{(9)}\le(\bar{\nu}_1)^{(9)}\le\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)}\le\nu^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)}\le\nu^{(9)}(t)\le(m_1)^{(9)},\quad\boxed{\nu^{(9)}(t)=\frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)}\le u^{(9)}(t)\le(\mu_1)^{(9)},\quad\boxed{u^{(9)}(t)=\frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(k) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^* :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]}$, $G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$

$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]}$, $T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]}$, $G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]}$ 513

$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]}$, $T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]}$ 514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]}$, $G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$

$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]}$, $T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$G_{24}^{*} = \frac{(a_{24})^{(4)}G_{25}^{*}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^{*})]}$, $G_{26}^{*} = \frac{(a_{26})^{(4)}G_{25}^{*}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^{*})]}$

$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]}$, $T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]}$ 517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]}$, $G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}\quad,\quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}\quad,\quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}\quad,\quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}\quad,\quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof:_Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i \qquad , T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*})=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G^*_{13}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G^*_{14}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G^*_{15}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a_{24}')^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a_{25}')^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a_{26}')^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b_{24}')^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^*\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b_{25}')^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^*\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b_{26}')^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^*\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \quad , \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a_{28}')^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a_{29}')^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a_{30}')^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b_{28}')^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T_{28}^*\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b_{29}')^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T_{29}^*\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b_{30}')^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T_{30}^*\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \quad , \frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \ , \ \frac{\partial (b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \ , \ \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586 A

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b''_i)^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left((\lambda)^{(1)}\right)^2+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\Big)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\Big]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)} + (b'_{34})^{(6)} - (r_{34})^{(6)})\{((\lambda)^{(6)} + (a'_{34})^{(6)} + (p_{34})^{(6)})$$

$$\left[\left(((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)})(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(q_{32})^{(6)}G^*_{32}\right)\right]$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(33)}T^*_{33} + (b_{33})^{(6)}s_{(32),(33)}T^*_{33}\right)$$

$$+\left(((\lambda)^{(6)} + (a'_{33})^{(6)} + (p_{33})^{(6)})(q_{32})^{(6)}G^*_{32} + (a_{32})^{(6)}(q_{33})^{(6)}G^*_{33}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(32)}T^*_{33} + (b_{33})^{(6)}s_{(32),(32)}T^*_{32}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$\left(((\lambda)^{(6)})^2 + (\,(b'_{32})^{(6)} + (b'_{33})^{(6)} - (r_{32})^{(6)} + (r_{33})^{(6)})\,(\lambda)^{(6)}\right)$$

$$+\left(((\lambda)^{(6)})^2 + (\,(a'_{32})^{(6)} + (a'_{33})^{(6)} + (p_{32})^{(6)} + (p_{33})^{(6)})\,(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)} + (a'_{32})^{(6)} + (p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^*_{33} + (a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^*_{32}\right)$$

$$\left(((\lambda)^{(6)} + (b'_{32})^{(6)} - (r_{32})^{(6)})s_{(33),(34)}T^*_{33} + (b_{33})^{(6)}s_{(32),(34)}T^*_{32}\right)\} = 0$$

$$+$$

$$(\lambda)^{(7)} + (b'_{38})^{(7)} - (r_{38})^{(7)})\{((\lambda)^{(7)} + (a'_{38})^{(7)} + (p_{38})^{(7)})$$

$$\left[\left(((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)})(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(q_{36})^{(7)}G^*_{36}\right)\right]$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(37)}T^*_{37} + (b_{37})^{(7)}s_{(36),(37)}T^*_{37}\right)$$

$$+\left(((\lambda)^{(7)} + (a'_{37})^{(7)} + (p_{37})^{(7)})(q_{36})^{(7)}G^*_{36} + (a_{36})^{(7)}(q_{37})^{(7)}G^*_{37}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(36)}T^*_{37} + (b_{37})^{(7)}s_{(36),(36)}T^*_{36}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$\left(((\lambda)^{(7)})^2 + (\,(b'_{36})^{(7)} + (b'_{37})^{(7)} - (r_{36})^{(7)} + (r_{37})^{(7)})\,(\lambda)^{(7)}\right)$$

$$+\left(((\lambda)^{(7)})^2 + (\,(a'_{36})^{(7)} + (a'_{37})^{(7)} + (p_{36})^{(7)} + (p_{37})^{(7)})\,(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)} + (a'_{36})^{(7)} + (p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37} + (a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left(((\lambda)^{(7)} + (b'_{36})^{(7)} - (r_{36})^{(7)})s_{(37),(38)}T^*_{37} + (b_{37})^{(7)}s_{(36),(38)}T^*_{36}\right)\} = 0$$

$$+$$

$$(\lambda)^{(8)} + (b'_{42})^{(8)} - (r_{42})^{(8)})\{((\lambda)^{(8)} + (a'_{42})^{(8)} + (p_{42})^{(8)})$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION TWELEVE

Spin-Correlation, Quantum Mechanical Probabilities, Mass And Spin-Parity Of The Higgs Boson Candidate, Decay Of The SM Higgs Boson To A Pair Of Z Bosons, Et Al: Age Quod Agis- Agere Sequitur Credere- Agere Sequitur (Esse)- Ad Pondus Omnium (Ad Pond Om)- Ad Referendum Model

(1) Under some further but quite plausible assumptions, this conclusion (That there must be a state, a hidden variable, characterizing the state of affairs in the world in more details than the quantum mechanical state, something that also reflects the missing elements of reality) **implies (eb) that** in some spin-correlation experiments the measured quantum mechanical probabilities should satisfy particular inequalities (Bell-type inequalities).

(2) The paradox consists in the fact that quantum probabilities **do not (e) satisfy** these inequalities. (And this paradoxical fact has been confirmed by several laboratory experiments in the last three decades. The problem is still open and hotly debated among both physicists and philosophers. It has motivated a wide range of research from the most fundamental quantum mechanical experiments through foundations of probability theory to the theory of stochastic causality as well as the metaphysics of free will).


Study of the Mass and Spin-Parity of the Higgs Boson Candidate via Its Decays to Z Boson Pairs: S. Chatrchyan et al. (CMS Collaboration): Published 21 February 2013

(3) A study is presented of the mass and spin-parity of the new boson recently observed at the LHC at a mass near 125 GeV. An integrated luminosity of 17.3 fb-1, collected by the CMS experiment in proton-proton collisions at center-of-mass energies of 7 and 8 TeV, is used. The measured mass in the ZZ channel, where both Z bosons decay to e or μ pairs, **is** 126.2±0.6 (stat) ±0.2 (syst) GeV. The angular distributions of the lepton pairs in this channel **are (e&eb) sensitive** to the spin-parity of the boson.

(4) Under the assumption of spin 0, the present data are **consistent (eb) with** the pure scalar hypothesis, **while (e) disfavoring** the pure pseudoscalar hypothesis.

Published by the American Physical Society under the terms of the Creative Commons Attribution 3.0 License. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. © 2013 CERN, for the CMS Collaboration URL: http://link.aps.org/doi/10.1103/PhysRevLett.110.081803 DOI: 10.1103/PhysRevLett.110.081803 PACS: 14.80.Bn, 12.60.-i, 13.85.Qk, 13.88. +e

New CMS results at Moriond (Electroweak) 2013



(5) HIG-13-002 Event 1: Event recorded with the CMS detector in 2012 at a proton-proton centre-of-mass energy of 8 TeV. The **event (eb) shows** characteristics expected from the decay of the SM Higgs boson to a pair of Z bosons.

(6) One of Z bosons which subsequently **decay(eb) to a** pair of electrons (green lines and green

towers)

(7) Other Z bosons **decays_(eb) to a** pair of muons (red lines). The event could also be due to known Standard Model background processes.

NOTATION

Module One

G_{13} : Category one of That there must be a state, a hidden variable, characterizing the state of affairs in the world in more details than the quantum mechanical state, something that also reflects the missing elements of reality

G_{14} : Category two of That there must be a state, a hidden variable, characterizing the state of affairs in the world in more details than the quantum mechanical state, something that also reflects the missing elements of reality

G_{15} : Category three of That there must be a state, a hidden variable, characterizing the state of affairs in the world in more details than the quantum mechanical state, something that also reflects the missing elements of reality

T_{13} : Category one of spin-correlation experiments the measured quantum mechanical probabilities should satisfy particular inequalities (Bell-type inequalities).

T_{14} : Category two of spin-correlation experiments the measured quantum mechanical probabilities should satisfy particular inequalities (Bell-type inequalities).

T_{15} : Category three of spin-correlation experiments the measured quantum mechanical probabilities should satisfy particular inequalities (Bell-type inequalities).

Module Two

G_{16} : Category one of these inequalities. (And this paradoxical fact has been confirmed by several laboratory experiments in the last three decades. The problem is still open and hotly debated among both physicists and philosophers. It has motivated a wide range of research from the most fundamental quantum mechanical experiments through foundations of probability theory to the theory of stochastic causality as well as the metaphysics of free will).

G_{17} : Category two of these inequalities. (And this paradoxical fact has been confirmed by several laboratory experiments in the last three decades. The problem is still open and hotly debated among both physicists and philosophers. It has motivated a wide range of research from the most fundamental quantum mechanical experiments through foundations of probability theory to the theory of stochastic causality as well as the metaphysics of free will).

G_{18} : Category three of these inequalities. (And this paradoxical fact has been confirmed by several laboratory experiments in the last three decades. The problem is still open and hotly debated among both physicists and philosophers. It has motivated a wide range of research from the most fundamental quantum mechanical experiments through foundations of probability theory to the theory of stochastic causality as well as the metaphysics of free will).

T_{16} : Category one of quantum probabilities

T_{17} : Category two of quantum probabilities

T_{18} : Category three of quantum probabilities

Module three

Relativistic generalization of Bell's inequalities in Wigner's form: N. V. Nikitin, K. S. Toms

A relativistic generalization of Bell's inequalities in Wigner's form **was (eb) obtained** for the decays of a pseudoscalar and a scalar particle to two particles having a nonzero spin (fermions and photons)after considerations of inequalities involving a full anticorrelation of final-particle spins and having a nonrelativistic analog and inequalities involving a full correlation of spins. It is shown that Bohr's complementarity principle **may(eb) be tested** experimentally in the relativistic region.

G_{20} : Category one of decays of a pseudoscalar and a scalar particle to two particles having a nonzero spin (fermions and photons)after considerations of inequalities involving a full anticorrelation of final-particle spins and having a nonrelativistic analog and inequalities involving a full correlation of spins. It is shown that Bohr's complementarity principle may (eb) be tested experimentally in the relativistic region.

G_{21} : Category two of decays of a pseudoscalar and a scalar particle to two particles having a nonzero spin (fermions and photons)after considerations of inequalities involving a full anticorrelation of final-particle spins and having a nonrelativistic analog and inequalities involving a full correlation of spins. It is shown that Bohr's complementarity principle may (eb) be tested experimentally in the relativistic region.

G_{22} : Category three of decays of a pseudoscalar and a scalar particle to two particles having a nonzero spin (fermions and photons)after considerations of inequalities involving a full anticorrelation of final-particle spins and having a nonrelativistic analog and inequalities involving a full correlation of spins. It is shown that Bohr's complementarity principle may (eb) be tested experimentally in the relativistic region.

T_{20} : Category one of relativistic generalization of Bell's inequalities in Wigner's form

T_{21} : Category two of relativistic generalization of Bell's inequalities in Wigner's form

T_{22} : Category three of relativistic generalization of Bell's inequalities in Wigner's form

Module four

G_{24} : Category one of angular distributions of the lepton pairs in this channel (mass and spin-parity of the new boson recently observed at the LHC at a mass near 125 GeV. An integrated luminosity of 17.3 fb-1, collected by the CMS experiment in proton-proton collisions at center-of-mass energies of 7 and 8 TeV, is used. The measured mass in the ZZ channel, where both Z bosons decay to e or μ pairs, **is** 126.2±0.6 (stat) ±0.2 (syst) GeV)

G_{25} : Category two of angular distributions of the lepton pairs in this channel (mass and spin-parity of the new boson recently observed at the LHC at a mass near 125 GeV. An integrated luminosity of 17.3 fb-1, collected by the CMS experiment in proton-proton collisions at center-of-mass energies of 7 and 8 TeV, is used. The measured mass in the ZZ channel, where both Z bosons decay to e or μ pairs, **is** 126.2±0.6 (stat) ±0.2 (syst) GeV)

G_{26} : Category three of angular distributions of the lepton pairs in this channel (mass and spin-parity of the new boson recently observed at the LHC at a mass near 125 GeV. An integrated luminosity of 17.3 fb-1, collected by the CMS experiment in proton-proton collisions at center-of-mass energies of 7 and 8 TeV, is used. The measured mass in the ZZ channel, where both Z bosons decay to e or μ pairs, **is** 126.2±0.6 (stat) ±0.2 (syst) GeV)

T_{24} : Category one of spin-parity of the boson.

T_{25} : Category two of spin-parity of the boson.

T_{26} : Category three of spin-parity of the boson.

Module five

G_{28} : Category one of Under the assumption of spin 0, the present data

G_{29} : Category two of Under the assumption of spin 0, the present data

G_{30} : Category three of Under the assumption of spin 0, the present data

T_{28} : Category one of pure scalar hypothesis

T_{29} : Category two of pure scalar hypothesis

T_{30} : Category three of pure scalar hypothesis

Module six

G_{32} : Category one of Under the assumption of spin 0, the present data

G_{33} : Category two of Under the assumption of spin 0, the present data

G_{34} : Category three of Under the assumption of spin 0, the present data

T_{32} : Category one of disfavors the pure pseudoscalar hypothesis

T_{33} : Category two of disfavors the pure pseudoscalar hypothesis

T_{34} : Category three of disfavors the pure pseudoscalar hypothesis

Module seven

G_{36} : Category one of proton-proton centre-of-mass energy of 8 TeV.(Systemic characterstics are taken in to consideration in the classification procedure)

G_{37} : Category two of proton-proton centre-of-mass energy of 8 TeV.(Systemic characterstics are taken in to consideration in the classification procedure)

G_{38} : Category three of proton-proton centre-of-mass energy of 8 TeV.(Systemic characterstics are taken in to consideration in the classification procedure)

T_{36} : Category one of characteristics expected from the decay of the SM Higgs boson to a pair of Z bosons.

T_{37} : Category two of characteristics expected from the decay of the SM Higgs boson to a pair of Z bosons.

T_{38} : Category three of characteristics expected from the decay of the SM Higgs boson to a pair of Z bosons.

Module eight

G_{40} : Category one of One of Z bosons (In various systems there are many such one of the bosons and the characterstics of the systems studied notwithstanding the unitaries and absoluteness of the fundamental entities have been taken in to consideration in the classification systems)

G_{41} : Category two of One of Z bosons (In various systems there are many such one of the bosons and the characterstics of the systems studied notwithstanding the unitaries and absoluteness of the fundamental entities have been taken in to consideration in the classification systems)

G_{42} : Category three of One of Z bosons (In various systems there are many such one of the bosons and the characterstics of the systems studied notwithstanding the unitaries and absoluteness of the fundamental entities have been taken in to consideration in the classification systems)

T_{40} : Category one of pair of electrons (green lines and green towers) (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{41} : Category two of pair of electrons (green lines and green towers)

T_{42} : Category three of pair of electrons (green lines and green towers)

Module Nine

G_{44} : Category one of Other Z bosons decay (under various systems with various properties)

G_{45} : Category two of Other Z bosons **decay**

G_{46} : Category three of Other Z bosons **decay**

T_{44} : Category one of pair of muons (red lines). The event could also be due to known Standard Model background processes.


T_{45} : Category two of pair of muons (red lines). The event could also be due to known Standard Model background processes.

T_{46} : Category three of pair of muons (red lines). The event could also be due to known Standard Model background processes.

.

SECTION TWELVE A

INTRODUCTION—VARIABLES USED



NOTATION

Module One

G_{13} : Category one of Bosons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{14} : Category two of Bosons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{15} : Category three of Bosons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{13} : Category one of hadrons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{14} : Category two of hadrons

T_{15} : Category three of hadrons

Module Two

G_{16} : Category one of fermions

G_{17} : Category two of fermions

G_{18} : Category three of fermions

T_{16} : Category one of hadrons

T_{17} : Category two of hadrons

T_{18} : Category three of hadrons

Module three

G_{20} : Category one of fermion- fermion interaction

G_{21} : Category two of fermion- fermion interaction

G_{22} : Category three of fermion- fermion interaction

T_{20} : Category one of Z (Systemic classification)

T_{21} : Category two of Z

T_{22} : Category three of Z

Module four

G_{24} : Category one of quark quark interaction

G_{25} : Category two of quark quark interaction

G_{26} : Category three of quark quark interaction

T_{24} : Category one of Gamma or g (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{25} : Category two of Gamma or g

T_{26} : Category three of Gamma or g

Module five

G_{28} : Category one of up quark-down quark interaction

G_{29} : Category two of up quark-down quark interaction

G_{30} : Category three of up quark-down quark interaction

T_{28} : Category one of W (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{29} : Category two of W

T_{30} : Category three of W

Module six

G_{32} : Category one of Lepton-Neutrino reaction

G_{33} : Category two of Lepton-Neutrino reaction


G_{34} : Category three of Lepton-Neutrino reaction

T_{32} : Category one of W

T_{33} : Category two of W

T_{34} : Category three of W

Module seven



G_{36} : Category one of Photon or Z-Boson and W^(+) interaction

G_{37} : Category two of Photon or Z-Boson and W^(+) interaction

G_{38} : Category three of Photon or Z-Boson and W^(+) interaction

T_{36} : Category one of W ^ (-): Classified based on the characterstics of the system under consideration. There may be different experiments like CERN

T_{37} : Category two of W ^ (-)

T_{38} : Category three of W ^ (-)

Module eight

G_{40} : Category one of Interaction between electroweak interaction: Charge is conserved

G_{41} : Category two of Interaction between electroweak interaction Charge is conserved


G_{42} : Category three of Interaction between electroweak interaction Charge is conserved

T_{40} : Category one of W^ (+) and W ^ (-)

T_{41} : Category two of W^ (+) and W ^ (-)

T_{42} : Category three of W^ (+) and W ^ (-)

Module Nine



G_{44} : Category one of g and g Charge is conserved

G_{45} : Category two of g and g Charge is conserved

G_{46} : Category three of g and g Charge is conserved

T_{44} : Category one of g and g Charge is conserved

T_{45} : Category two of g and g Charge is conserved

T_{46} : Category three of g and g Charge is conserved

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}$: $(a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}$, $(b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$

$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}, (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30}$ 27

$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28}$ 28

$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29}$ 29

$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30}$ 30

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32}$ 31

$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33}$ 32

$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34}$ 33

$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32}$ 34

$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33}$ 35

$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34}$ 36

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36}$ 37

$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37}$ *38*

$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38}$ *39*

$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36}$ *40*

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[\begin{array}{ccc} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[\begin{array}{ccc} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} \quad \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right]G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7)}(T_{37},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8)}(T_{41},t)$ $+(a''_{41})^{(8,8)}(T_{41},t)$ $+(a''_{42})^{(8,8)}(T_{41},t)$ are eight augmentation coefficient for 1,2,3

$+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\left[-(b''_{13})^{(1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\left[-(b''_{14})^{(1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\left[-(b''_{15})^{(1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{15} \quad 60$$

Where $-(b''_{13})^{(1)}(G,t)$, $-(b''_{14})^{(1)}(G,t)$, $-(b''_{15})^{(1)}(G,t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,)}(G_{19},t)$, $-(b''_{17})^{(2,2,)}(G_{19},t)$, $-(b''_{18})^{(2,2,)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) \\ +(a''_{20})^{(3,3,3)}(T_{21},t) & +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{32})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) \\ +(a''_{44})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) \\ +(a''_{21})^{(3,3,3)}(T_{21},t) & +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{33})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) \\ +(a''_{45})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) \\ +(a''_{22})^{(3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6)}(T_{33},t) & +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) \\ +(a''_{46})^{(9,9)}(T_{45},t) & & \end{array}\right] G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) \\ -(b''_{20})^{(3,3,3,)}(G_{23},t) & -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) \\ -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) & -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) \\ -(b''_{44})^{(9,9)}(G_{47},t) & & \end{array}\right] T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[\begin{array}{ccc} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) \\ -(b''_{21})^{(3,3,3,)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) & -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) \\ -(b''_{45})^{(9,9)}(G_{47},t) & & \end{array}\right] T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[\begin{array}{ccc} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) \\ -(b''_{22})^{(3,3,3,)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) & -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) \\ -(b''_{46})^{(9,9)}(G_{47},t) & & \end{array}\right] T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for

category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \tag{67}$$

$$(a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \tag{68}$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \tag{69}$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation

coefficients for category 1, 2 and 3

$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)} & \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$\boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}$ *are fourth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}$ *are fifth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)}$ *are sixth detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}$ $\boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{24} \qquad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{25} \qquad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{26} \quad 75$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}, \boxed{(a''_{25})^{(4)}(T_{25},t)}, \boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad 76$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad 77$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad 78$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{matrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{matrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{matrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{matrix} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{matrix}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{30} \quad 84$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
are first detrition coefficients for category 1, 2 *and* 3
$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3
$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3
$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad 85$$
$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \quad 86$$
$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \quad 87$$
$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[\begin{matrix} (a_{36}')^{(7)}\boxed{+(a_{36}'')^{(7)}(T_{37},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{13} \quad (91)$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[\begin{matrix} (a_{37}')^{(7)}\boxed{+(a_{37}'')^{(7)}(T_{37},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{14} \quad (92)$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[\begin{matrix} (a_{38}')^{(7)}\boxed{+(a_{38}'')^{(7)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{matrix}\right] G_{15} \quad (93)$$

Where $\boxed{(a_{36}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{37}'')^{(7)}(T_{37},t)}$, $\boxed{(a_{38}'')^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b'_{36})^{(7)}\boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} \qquad 95$$

$$= (a_{40})^{(8)}G_{41}$$

$$- \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[\begin{array}{ccc} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[\begin{array}{ccc} (a'_{42})^{(8)}\boxed{+(a''_{42})^{(8)}(T_{41},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{15}$$

Where $\boxed{+(a''_{40})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8)}(T_{41},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[\begin{array}{ccc} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[\begin{array}{ccc} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt}$$ *96*
$$= (a_{44})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$
$$\frac{dG_{45}}{dt}$$
$$= (a_{45})^{(9)}G_{44}$$
$$-\begin{bmatrix} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$
$$\frac{dG_{46}}{dt}$$
$$= (a_{46})^{(9)}G_{45}$$
$$-\begin{bmatrix} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $+(a''_{44})^{(9)}(T_{45},t)$, $+(a''_{45})^{(9)}(T_{45},t)$, $+(a''_{46})^{(9)}(T_{37},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\left[\begin{array}{ccc} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{14}$$

$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\left[\begin{array}{ccc} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{array}\right] T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b_{34}'')^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \leq (p_i)^{(1)} \leq (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \leq (r_i)^{(1)} \leq (b_i')^{(1)} \leq (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together With $(\widehat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants $(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}, i = 13,14,15,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)} (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)} |T_{17} - T_{17}'| e^{-(\widehat{M}_{16})^{(2)} t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)} ||(G_{19}) - (G_{19})'|| e^{-(\widehat{M}_{16})^{(2)} t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\widehat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\widehat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants

$(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)} (\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)} (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \qquad 118$$

$$\lim_{G \to \infty} (b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)} |T_{25} - T_{25}'| e^{-(\hat{M}_{24})^{(4)} t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}((G_{27}), t)| < (\hat{k}_{24})^{(4)} ||(G_{27}) - (G_{27})'|| e^{-(\hat{M}_{24})^{(4)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and $(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i,j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \leq (p_i)^{(5)} \leq (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \leq (r_i)^{(5)} \leq (b_i')^{(5)} \leq (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \leq (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$ and$(a_i'')^{(5)}(T_{29}, t)$. (T_{29}', t) and (T_{29}, t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[\ (b_i)^{(5)} + (b_i')^{(5)} + \ (\hat{B}_{28})^{(5)} + \ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le \ (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(6)}\ (T_{33}, t) = (p_i)^{(6)} \qquad 128$$

$$\lim_{G \to \infty}(b_i'')^{(6)}\ \big((G_{35}), t\big) = \ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)}|T_{33} - T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}\big((G_{35}), t\big)| < (\hat{k}_{32})^{(6)}||(G_{35}) - (G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and$(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)}, (\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ , \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, \ (r_i)^{(6)}, i = 32,33,34,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (a_i)^{(6)} + (a_i')^{(6)} + \ (\hat{A}_{32})^{(6)} + \ (\hat{P}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)} + (b_i')^{(6)} + \ (\hat{B}_{32})^{(6)} + \ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i,j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \leq (p_i)^{(7)} \leq (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \leq (r_i)^{(7)} \leq (b_i')^{(7)} \leq (\hat{B}_{36})^{(7)} \qquad 132$$

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}\big((G_{39}), t\big) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\big((G_{39}), t\big)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\widehat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i,j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}\ (T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}\ \big((G_{43}), t\big) = \ (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - \ T_{41}'| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}\big((G_{43}), t\big)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\widehat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $\big[(\hat{k}_{40})^{(8)}, (\widehat{M}_{40})^{(8)}\big]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\ , \frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\widehat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (a_i)^{(8)} + (a_i')^{(8)} + \ (\hat{A}_{40})^{(8)} + \ (\hat{P}_{40})^{(8)}\ (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\widehat{M}_{40})^{(8)}}[\ (b_i)^{(8)} + (b_i')^{(8)} + \ (\hat{B}_{40})^{(8)} + \ (\hat{Q}_{40})^{(8)}\ \ (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}\,(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}\,(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - \ T_{45}'| e^{-(\widehat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}\big((G_{47}), t\big)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\widehat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and$(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)}, (\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)}t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\widehat{M}_{28})^{(5)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}\,, \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}\left(s_{(13)}\right) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}\left(s_{(13)}\right) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}\left(T_{14}\left(s_{(13)}\right), s_{(13)}\right)\right) G_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{13}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}\left(s_{(13)}\right) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{14}\left(s_{(13)}\right)\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}\left(s_{(13)}\right) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}\left(G\left(s_{(13)}\right), s_{(13)}\right)\right) T_{15}\left(s_{(13)}\right)\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)} (T_{21}(s_{(20)}), s_{(20)}) \right) G_{22}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{20}(s_{(20)}) \right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{21}(s_{(20)}) \right] ds_{(20)}$$

$$\overline{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^{0} + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)} (G_{23}(s_{(20)}), s_{(20)}) \right) T_{22}(s_{(20)}) \right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)} (T_{25}(s_{(24)}), s_{(24)}) \right) G_{26}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{24}(s_{(24)}) \right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{25}(s_{(24)}) \right] ds_{(24)}$$

$$\overline{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^{0} + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)} (G_{27}(s_{(24)}), s_{(24)}) \right) T_{26}(s_{(24)}) \right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\overline{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^{0} + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^{0} + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^{0} + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^{0} + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^{0} + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^{0} + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^{0} + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^{0} + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{44})^{(9)}$, $T_i^0 \le (\hat{Q}_{44})^{(9)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{P}_{13})^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} \left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}} \left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} \left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that 172

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}}\right)\right] ds_{(36)} =$$
$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1\right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0\right) e^{\left(-\frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0}\right)} + (\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \quad 180$$
$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8
Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$
$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{Q}_{13})^{(1)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad (184)$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric (185)

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \quad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\,(t)=0$ *and* $T_i\,(t)=0$

From 19 to 24 it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(1)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a_{14}')^{(1)}G_{14}$ and by integrating

$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$

In the same way , one can obtain

$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\,(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}\,(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}\,(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}},\widetilde{T_{19}}$: $\left(\widetilde{G_{19}},\widetilde{T_{19}}\right)=\mathcal{A}^{(2)}(G_{19},T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{16})^{(2)}|G_{17}^{(1)}-G_{17}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)}+$$

$$\int_0^t\{(a'_{16})^{(2)}|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)|G_{16}^{(1)}-G_{16}^{(2)}|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}+$$

$$G_{16}^{(2)}|(a''_{16})^{(2)}\left(T_{17}^{(1)},s_{(16)}\right)-(a''_{16})^{(2)}\left(T_{17}^{(2)},s_{(16)}\right)|\;e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0,t]$ 197

From the hypotheses it follows

$$|(G_{19})^{(1)}-(G_{19})^{(2)}|e^{-(\widehat{M}_{16})^{(2)}t}\le$$
$$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)}+(a'_{16})^{(2)}+(\widehat{A}_{16})^{(2)}+(\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)},(T_{19})^{(1)};\;(G_{19})^{(2)},(T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i=16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t)=0$ and $T_i(t)=0$ 200

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(2)}-(a_i'')^{(2)}(T_{17}(s_{(16)}),s_{(16)})\}ds_{(16)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(2)}t)}>0\quad\text{for } t>0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1,\left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16}<(\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt}\le\left((\widehat{M}_{16})^{(2)}\right)_1-(a'_{17})^{(2)}G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t),t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t),t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right)$, $t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t),t\right) = (b'_{18})^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\widehat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{20})^{(3)}\right] \leq (\widehat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})} \right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$\left|\tilde{G}_{20}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{20})^{(3)} \left|G_{21}^{(1)} - G_{21}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\, ds_{(20)} +$$

$$\int_0^t \{(a'_{20})^{(3)} \left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right)\left|G_{20}^{(1)} - G_{20}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}} +$$

$$G_{20}^{(2)} \left|(a''_{20})^{(3)}\left(T_{21}^{(1)}, s_{(20)}\right) - (a''_{20})^{(3)}\left(T_{21}^{(2)}, s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)} s_{(20)}} e^{(\widehat{M}_{20})^{(3)} s_{(20)}}\} ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)} t} \leq \qquad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ *and* $(\widehat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)} G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1 / (a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2 / (a'_{22})^{(3)}$$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)=(b_{21}')^{(3)}$ then $T_{21}\to\infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)}-(b_i'')^{(3)}\left((G_{23})(t),t\right)<\varepsilon_3, T_{20}(t)>(m)^{(3)}$

Then $\frac{dT_{21}}{dt}\geq(a_{21})^{(3)}(m)^{(3)}-\varepsilon_3 T_{21}$ which leads to 220

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t})+T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t}=\frac{1}{2}$ it results

$T_{21}\geq\left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right),\quad t=log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right)=(b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}<1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)}+\left((\hat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\tilde{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{24})^{(4)}\right]\leq(\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$

$sup_i\{max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t},max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\tilde{M}_{24})^{(4)}t}\}$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\tilde{M}_{24})^{(4)}s_{(24)}}e^{(\tilde{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \qquad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\ (G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24{,}25{,}26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 228

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0e^{\left(-(b_i')^{(4)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\leq\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25}\leq\left((\widehat{M}_{24})^{(4)}\right)_2=G_{25}^0+2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26}\leq\left((\widehat{M}_{24})^{(4)}\right)_3=G_{26}^0+2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If $G_{25}\ or\ G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24}\ \ is$ bounded, from below, the same property holds for $G_{25}\ and\ G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}\ ((G_{27})(t),t))=(b'_{25})^{(4)}$ then $T_{25}\ \to\infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{31})}, \widetilde{(T_{31})}$: $\left(\widetilde{(G_{31})}, \widetilde{(T_{31})}\right) = \mathcal{A}^{(5)}\left((G_{31}), (T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{28})^{(5)}\left|G_{29}^{(1)} - G_{29}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a_{28}')^{(5)}\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{-(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a_{28}'')^{(5)}\left(T_{29}^{(1)}, s_{(28)}\right)\left|G_{28}^{(1)} - G_{28}^{(2)}\right|e^{-(\hat{M}_{28})^{(5)}s_{(28)}}e^{(\hat{M}_{28})^{(5)}s_{(28)}} +$$

$G_{28}^{(2)}|(a_{28}'')^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a_{28}'')^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a_{28}')^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};\ (G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a_{28}'')^{(5)}$ and $(b_{28}'')^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 239

it results

$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\geq 0$

$T_i\ (t)\geq T_i^0 e^{(-(b_i')^{(5)}t)}>0\ \ \text{for t}>0$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\leq\left((\widehat{M}_{28})^{(5)}\right)_1-(a_{29}')^{(5)}G_{29}$ and by integrating

$G_{29}\leq\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30}\leq\left((\widehat{M}_{28})^{(5)}\right)_3=G_{30}^0+2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t))=(b_{29}')^{(5)}$ then $T_{29}\ \to\infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ >t_5$

$(b_{29})^{(5)}-(b_i'')^{(5)}((G_{31})(t),t)<\varepsilon_5,T_{28}\ (t)>(m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b''_{30})^{(5)}\left((G_{31})(t), t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{33}, G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\tilde{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(\left((G_{35})^{(1)}, (T_{35})^{(1)}\right), \left((G_{35})^{(2)}, (T_{35})^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}), (T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{32})^{(6)}\left|G_{33}^{(1)} - G_{33}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)} +$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{-(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right)\left|G_{32}^{(1)} - G_{32}^{(2)}\right|e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}} +$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)}, s_{(32)}\right) - (a''_{32})^{(6)}\left(T_{33}^{(2)}, s_{(32)}\right)\right|\ e^{-(\tilde{M}_{32})^{(6)}s_{(32)}}e^{(\tilde{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$\left|(G_{35})^{(1)} - (G_{35})^{(2)}\right| e^{-(\widehat{M}_{32})^{(6)}t} \leq$ 248

$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)} + (a'_{32})^{(6)} + (\widehat{A}_{32})^{(6)} + (\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right) d\left(((G_{35})^{(1)}, (T_{35})^{(1)}; (G_{35})^{(2)}, (T_{35})^{(2)})\right)$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)} e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i = 32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 250

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a'_{33})^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1 / (a'_{33})^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2 / (a'_{34})^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b'_{33})^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b''_{34})^{(6)}\left((G_{35})(t), t(t), t\right) = (b'_{34})^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\tilde{M}_{36})^{(7)}} < 1$ and to choose $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\tilde{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)}, (T_{39})^{(1)}), ((G_{39})^{(2)}, (T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\tilde{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\tilde{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}), (\widetilde{T_{39}}) : \left((\widetilde{G_{39}}), (\widetilde{T_{39}})\right) = \mathcal{A}^{(7)}((G_{39}), (T_{39}))$
It results

$$\left|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{36})^{(7)}\left|G_{37}^{(1)} - G_{37}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{-(\tilde{M}_{36})^{(7)}s_{(36)}} +$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right)\left|G_{36}^{(1)} - G_{36}^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}} +$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)}, s_{(36)}\right) - (a''_{36})^{(7)}\left(T_{37}^{(2)}, s_{(36)}\right)\right|\ e^{-(\tilde{M}_{36})^{(7)}s_{(36)}}e^{(\tilde{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)} - (G_{39})^{(2)}\right|e^{-(\tilde{M}_{36})^{(7)}t} \leq \quad 259$$
$$\frac{1}{(\tilde{M}_{36})^{(7)}}\left((a_{36})^{(7)} + (a'_{36})^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)}(\hat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)}, (T_{39})^{(1)};\ (G_{39})^{(2)}, (T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered 260

as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ and $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i = 36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 261

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} is bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is

unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b''_{38})^{(7)}\left((G_{39})(t),t\right)=(b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}<1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)}+\left((\widehat{P}_{40})^{(8)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{40})^{(8)}\right]\leq(\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $(\widetilde{G_{43}}),(\widetilde{T_{43}})$: $\left((\widetilde{G_{43}}),(\widetilde{T_{43}})\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\leq \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i = 40,41,42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 275

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(8)} - (a_i'')^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(8)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a_{41}')^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a_{41}')^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a_{42}')^{(8)}$$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ \ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t), t)) = (b_{41}')^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t\ > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t), t\right) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279

A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})}\right) = \mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and

respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\ (t) = 0\ and\ T_i\ (t) = 0$

From 99 to 44 it results

$$G_i\ (t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}(T_{45}(s_{(44)}), s_{(44)})\} ds_{(44)}\right]} \geq 0$$

$$T_i\ (t) \geq T_i^0 e^{(-(b_i')^{(9)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a_{45}')^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a_{45}')^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2 / (a_{46}')^{(9)}$$

If $G_{45}\ or\ G_{46}$ is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for $G_{45}\ and\ G_{46}$. The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t), t\right)) = (b_{45}')^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t), t\right) < \varepsilon_9, T_{44}\ (t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right), \quad t = log\ \frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b_{46}'')^{(9)}\left((G_{47})(t), t\right) = (b_{46}')^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \le -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \le -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \le -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \le -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations
$(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G^0_{13}}{G^0_{14}}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T^0_{13}}{T^0_{14}}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G^0_{13} e^{(S_1)^{(1)} t}$

where $(p_i)^{(1)}$ is defined by equation

$\frac{1}{(m_1)^{(1)}} G^0_{13} e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G^0_{13} e^{(S_1)^{(1)} t}$

$\left(\frac{(a_{15})^{(1)} G^0_{13}}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G^0_{15} e^{-(S_2)^{(1)} t} \le G_{15}(t) \le\right.$ 286

$$\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t})$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t}\le T_{13}(t)\le\frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t}\le T_{15}(t)\le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)},(S_2)^{(1)},(R_1)^{(1)},(R_2)^{(1)}$:- 290

Where $(S_1)^{(1)}=(a_{13})^{(1)}(m_2)^{(1)}-(a'_{13})^{(1)}$

$(S_2)^{(1)}=(a_{15})^{(1)}-(p_{15})^{(1)}$

$(R_1)^{(1)}=(b_{13})^{(1)}(\mu_2)^{(1)}-(b'_{13})^{(1)}$

$(R_2)^{(1)}=(b'_{15})^{(1)}-(r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)},(\sigma_2)^{(2)},(\tau_1)^{(2)},(\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)}\le-(a'_{16})^{(2)}+(a'_{17})^{(2)}-(a''_{16})^{(2)}(\mathrm{T}_{17},t)+(a''_{17})^{(2)}(\mathrm{T}_{17},t)\le-(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)}\le-(b'_{16})^{(2)}+(b'_{17})^{(2)}-(b''_{16})^{(2)}\big((G_{19}),t\big)-(b''_{17})^{(2)}\big((G_{19}),t\big)\le-(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)},(\nu_2)^{(2)},(u_1)^{(2)},(u_2)^{(2)}$: 295

By $(\nu_1)^{(2)}>0$, $(\nu_2)^{(2)}<0$ and respectively $(u_1)^{(2)}>0$, $(u_2)^{(2)}<0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_1)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2+(\tau_1)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ and 298

Definition of $(\bar{\nu}_1)^{(2)},,(\bar{\nu}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)}>0$, $(\bar{\nu}_2)^{(2)}<0$ and respectively $(\bar{u}_1)^{(2)}>0$, $(\bar{u}_2)^{(2)}<0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2+(\sigma_2)^{(2)}\nu^{(2)}-(a_{16})^{(2)}=0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2+(\tau_2)^{(2)}u^{(2)}-(b_{16})^{(2)}=0$ 302

Definition of $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)},(m_2)^{(2)},(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)},\ \boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)},$ 306

and $\boxed{(\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$

$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$ 318

$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(\nu_1)^{(3)}, (\nu_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(\nu_1)^{(3)} > 0$, $(\nu_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{\nu}_1)^{(3)} > 0$, $(\bar{\nu}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(\nu^{(3)}\big)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, $\boldsymbol{if}\ (\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, $\boldsymbol{if}\ (\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}\ (u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}\ (u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G^0_{20}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G^0_{20}e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G^0_{20}e^{(S_1)^{(3)}t} \quad 323$$

$$\Big(\frac{(a_{22})^{(3)}G^0_{20}}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\Big[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\Big] + G^0_{22}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \quad 324$$

$$\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+\ G_{22}^{0}e^{-(a'_{22})^{(3)}t})$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t}\leq T_{20}(t)\leq T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t}\leq T_{20}(t)\leq \frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t}\leq T_{22}(t)\leq \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)},(S_2)^{(3)},(R_1)^{(3)},(R_2)^{(3)}$:- 328

Where $(S_1)^{(3)}=(a_{20})^{(3)}(m_2)^{(3)}-(a'_{20})^{(3)}$

$$(S_2)^{(3)}=(a_{22})^{(3)}-(p_{22})^{(3)}$$

$$(R_1)^{(3)}=(b_{20})^{(3)}(\mu_2)^{(3)}-(b'_{20})^{(3)}$$

$$(R_2)^{(3)}=(b'_{22})^{(3)}-(r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$:

$(\sigma_1)^{(4)},(\sigma_2)^{(4)},(\tau_1)^{(4)},(\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)}\leq -(a'_{24})^{(4)}+(a'_{25})^{(4)}-(a''_{24})^{(4)}(T_{25},t)+(a''_{25})^{(4)}(T_{25},t)\leq -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)}\leq -(b'_{24})^{(4)}+(b'_{25})^{(4)}-(b''_{24})^{(4)}\big((G_{27}),t\big)-(b''_{25})^{(4)}\big((G_{27}),t\big)\leq -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)},(v_2)^{(4)},(u_1)^{(4)},(u_2)^{(4)},v^{(4)},u^{(4)}$: 329

By $(v_1)^{(4)}>0$, $(v_2)^{(4)}<0$ and respectively $(u_1)^{(4)}>0$, $(u_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_1)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_1)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$ and

Definition of $(\bar{v}_1)^{(4)},,(\bar{v}_2)^{(4)},(\bar{u}_1)^{(4)},(\bar{u}_2)^{(4)}$: 330

By $(\bar{v}_1)^{(4)}>0$, $(\bar{v}_2)^{(4)}<0$ and respectively $(\bar{u}_1)^{(4)}>0$, $(\bar{u}_2)^{(4)}<0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}=0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2+(\tau_2)^{(4)}u^{(4)}-(b_{24})^{(4)}=0$

Definition of $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)},(v_0)^{(4)}$:-

If we define $(m_1)^{(4)},(m_2)^{(4)},(\mu_1)^{(4)},(\mu_2)^{(4)}$ by

$$(m_2)^{(4)}=(v_0)^{(4)},(m_1)^{(4)}=(v_1)^{(4)},\ \boldsymbol{if}\ (v_0)^{(4)}<(v_1)^{(4)}$$

$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}$, $\boldsymbol{if}$ $(\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)}$,

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}$, $\boldsymbol{if}$ $(\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$

and analogously

331

$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}$, $\boldsymbol{if}$ $(u_0)^{(4)} < (u_1)^{(4)}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}$, $\boldsymbol{if}$ $(u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)}$,

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}$ $(\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{24}(t) \leq G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \leq G_{25}(t) \leq \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t}$$

333

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t}\right] + G_{26}^0 e^{-(S_2)^{(4)} t} \leq G_{26}(t) \leq \right.$$

334

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)} t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \leq T_{24}(t) \leq \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}$$

335

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \leq T_{26}(t) \leq$$

336

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t}\right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)}, , (\bar{v}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (v_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,

and $\boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^{0}e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}}G_{28}^{0}e^{((S_1)^{(5)}-(p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}}G_{28}^{0}e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_1)^{(5)}((S_1)^{(5)}-(p_{28})^{(5)}-(S_2)^{(5)})}\left[e^{((S_1)^{(5)}-(p_{28})^{(5)})t}-e^{-(S_2)^{(5)}t}\right]+G_{30}^{0}e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left.\frac{(a_{30})^{(5)}G_{28}^{0}}{(m_2)^{(5)}((S_1)^{(5)}-(a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t}-e^{-(a'_{30})^{(5)}t}\right]+G_{30}^{0}e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^{0}e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^{0}e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^{0}}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t}-e^{-(b'_{30})^{(5)}t}\right]+T_{30}^{0}e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}((G_{35}), t) - (b''_{33})^{(6)}((G_{35}), t) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}(v^{(6)})^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$

and $(b_{33})^{(6)}(u^{(6)})^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{\nu}_1)^{(6)},,(\bar{\nu}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{\nu}_1)^{(6)} > 0$, $(\bar{\nu}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)} = 0$ and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)},(\nu_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (\nu_0)^{(6)}, (m_1)^{(6)} = (\nu_1)^{(6)},$ ***if*** $(\nu_0)^{(6)} < (\nu_1)^{(6)}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\bar{\nu}_6)^{(6)}$, ***if*** $(\nu_1)^{(6)} < (\nu_0)^{(6)} < (\bar{\nu}_1)^{(6)},$

and $\boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (\nu_1)^{(6)}, (m_1)^{(6)} = (\nu_0)^{(6)},$ ***if*** $(\bar{\nu}_1)^{(6)} < (\nu_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},$ ***if*** $(u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, ***if*** $(u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)},$ ***if*** $(\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G_{34}^0 e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t}-e^{-(R_2)^{(6)}t}\right]+T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)},(S_2)^{(6)},(R_1)^{(6)},(R_2)^{(6)}$:- 359

Where $(S_1)^{(6)}=(a_{32})^{(6)}(m_2)^{(6)}-(a'_{32})^{(6)}$

$(S_2)^{(6)}=(a_{34})^{(6)}-(p_{34})^{(6)}$

$(R_1)^{(6)}=(b_{32})^{(6)}(\mu_2)^{(6)}-(b'_{32})^{(6)}$

$(R_2)^{(6)}=(b'_{34})^{(6)}-(r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$:

$(\sigma_1)^{(7)},(\sigma_2)^{(7)},(\tau_1)^{(7)},(\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)}\le-(a'_{36})^{(7)}+(a'_{37})^{(7)}-(a''_{36})^{(7)}(T_{37},t)+(a''_{37})^{(7)}(T_{37},t)\le-(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)}\le-(b'_{36})^{(7)}+(b'_{37})^{(7)}-(b''_{36})^{(7)}\big((G_{39}),t\big)-(b''_{37})^{(7)}\big((G_{39}),t\big)\le-(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)},(\nu_2)^{(7)},(u_1)^{(7)},(u_2)^{(7)},\nu^{(7)},u^{(7)}$: 361

By $(\nu_1)^{(7)}>0,(\nu_2)^{(7)}<0$ and respectively $(u_1)^{(7)}>0,(u_2)^{(7)}<0$ the roots of the equations
$(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_1)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_1)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$ and

Definition of $(\bar{\nu}_1)^{(7)},,(\bar{\nu}_2)^{(7)},(\bar{u}_1)^{(7)},(\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)}>0,(\bar{\nu}_2)^{(7)}<0$ and respectively $(\bar{u}_1)^{(7)}>0,(\bar{u}_2)^{(7)}<0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2+(\sigma_2)^{(7)}\nu^{(7)}-(a_{36})^{(7)}=0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2+(\tau_2)^{(7)}u^{(7)}-(b_{36})^{(7)}=0$

Definition of $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)},(\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)},(m_2)^{(7)},(\mu_1)^{(7)},(\mu_2)^{(7)}$ by

$(m_2)^{(7)}=(\nu_0)^{(7)},(m_1)^{(7)}=(\nu_1)^{(7)},\ \boldsymbol{if}\ (\nu_0)^{(7)}<(\nu_1)^{(7)}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\bar{\nu}_1)^{(7)},\boldsymbol{if}\ (\nu_1)^{(7)}<(\nu_0)^{(7)}<(\bar{\nu}_1)^{(7)}$,
and $\boxed{(\nu_0)^{(7)}=\frac{G_{36}^{0}}{G_{37}^{0}}}$

$(m_2)^{(7)}=(\nu_1)^{(7)},(m_1)^{(7)}=(\nu_0)^{(7)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)}<(\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)},\ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)}, (v_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, v^{(8)}, u^{(8)}$: 372

By $(v_1)^{(8)} > 0$, $(v_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{v}_1)^{(8)}, , (\bar{v}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)} > 0$, $(\bar{v}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}$, $(m_2)^{(8)}$, $(\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, $\boldsymbol{if}\ (v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, $\boldsymbol{if}\ (v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,
and $\boxed{(v_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,
and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G^0_{40}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G^0_{40}e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G^0_{40}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t}\le G_{42}(t)\le\right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t}\le T_{40}(t)\le\frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t}\le T_{42}(t)\le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)},(S_2)^{(8)},(R_1)^{(8)},(R_2)^{(8)}$:- 381

Where $(S_1)^{(8)}=(a_{40})^{(8)}(m_2)^{(8)}-(a'_{40})^{(8)}$

$(S_2)^{(8)}=(a_{42})^{(8)}-(p_{42})^{(8)}$

$(R_1)^{(8)}=(b_{40})^{(8)}(\mu_2)^{(8)}-(b'_{40})^{(8)}$

$(R_2)^{(8)}=(b'_{42})^{(8)}-(r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$:

$(\sigma_1)^{(9)},(\sigma_2)^{(9)},(\tau_1)^{(9)},(\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)}\le-(a'_{44})^{(9)}+(a'_{45})^{(9)}-(a''_{44})^{(9)}(T_{45},t)+(a''_{45})^{(9)}(T_{45},t)\le-(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)}\le-(b'_{44})^{(9)}+(b'_{45})^{(9)}-(b''_{44})^{(9)}\big((G_{47}),t\big)-(b''_{45})^{(9)}\big((G_{47}),t\big)\le-(\tau_1)^{(9)}$$

Definition of $(\nu_1)^{(9)},(\nu_2)^{(9)},(u_1)^{(9)},(u_2)^{(9)},\nu^{(9)},u^{(9)}$:

By $(\nu_1)^{(9)}>0$, $(\nu_2)^{(9)}<0$ and respectively $(u_1)^{(9)}>0$, $(u_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_1)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{\nu}_1)^{(9)},,(\bar{\nu}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)}>0$, $(\bar{\nu}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2+(\sigma_2)^{(9)}\nu^{(9)}-(a_{44})^{(9)}=0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}, \ \boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}, \boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)},$$

and $\boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$

$$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}, \boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)},$$

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{44}(t) \le G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \le G_{45}(t) \le \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \le G_{46}(t) \le \right.$$

$$\left. \frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \le T_{44}(t) \le \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \le T_{46}(t) \le$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$$

$$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$$

$$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \leq \frac{d\nu^{(1)}}{dt} \leq -\left((a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)} \leq v^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{v^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $v^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(\mathrm{T}_{17}, \mathrm{t})\right) - (a''_{17})^{(2)}(\mathrm{T}_{17}, \mathrm{t})v^{(2)} - (a_{17})^{(2)} v^{(2)}$$

Definition of $v^{(2)}$:- $\boxed{v^{(2)} = \frac{\mathrm{G}_{16}}{\mathrm{G}_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)} v^{(2)} - (a_{16})^{(2)}\right) \leq \frac{\mathrm{d}v^{(2)}}{\mathrm{dt}} \leq -\left((a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_1)^{(2)} v^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$v^{(2)}(t) \geq \frac{(\nu_1)^{(2)} + (\mathrm{C})^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (\mathrm{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\mathrm{C})^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq v^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a_{16}'')^{(2)} = (a_{17}'')^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b_{16}'')^{(2)} = (b_{17}'')^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a_{20}')^{(3)} - (a_{21}')^{(3)} + (a_{20}'')^{(3)}(T_{21}, t)\right) - (a_{21}'')^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_2)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)\leq\frac{dv^{(3)}}{dt}\leq-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2+(\sigma_1)^{(3)}v^{(3)}-(a_{20})^{(3)}\right)$$

400

From which one obtains

For $0<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(v_1)^{(3)}<(\bar{v}_1)^{(3)}$

$$v^{(3)}(t)\geq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_0)^{(3)}\right)t\right]}}\quad,\quad\boxed{(C)^{(3)}=\frac{(v_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(v_2)^{(3)}}}$$

it follows $(v_0)^{(3)}\leq v^{(3)}(t)\leq(v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(3)}=\frac{(\bar{v}_1)^{(3)}-(v_0)^{(3)}}{(v_0)^{(3)}-(\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)}\leq v^{(3)}(t)\leq(\bar{v}_1)^{(3)}$

If $0<(v_1)^{(3)}<(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}<(\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)}\leq\frac{(v_1)^{(3)}+(C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)}-(v_2)^{(3)}\right)t\right]}}\leq v^{(3)}(t)\leq$$

$$\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(\bar{v}_1)^{(3)}$$

If $0<(v_1)^{(3)}\leq(\bar{v}_1)^{(3)}\leq(v_0)^{(3)}=\frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)}\leq v^{(3)}(t)\leq\frac{(\bar{v}_1)^{(3)}+(\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)}-(\bar{v}_2)^{(3)}\right)t\right]}}\leq(v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)}\leq v^{(3)}(t)\leq(m_1)^{(3)},\quad\boxed{v^{(3)}(t)=\frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)}\leq u^{(3)}(t)\leq(\mu_1)^{(3)},\quad\boxed{u^{(3)}(t)=\frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{24}'')^{(4)} = (a_{25}'')^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b_{24}'')^{(4)} = (b_{25}'')^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a_{28}')^{(5)} - (a_{29}')^{(5)} + (a_{28}'')^{(5)}(T_{29}, t)\right) - (a_{29}'')^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\quad \boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}}{5+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)}=\frac{(\nu_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{5+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)}=\frac{(\bar{\nu}_1)^{(5)}-(\nu_0)^{(5)}}{(\nu_0)^{(5)}-(\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)}=\frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t)=\frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t)=\frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{dv^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right) - (a''_{33})^{(6)}(T_{33},t)v^{(6)} - (a_{33})^{(6)}v^{(6)}$$

Definition of $v^{(6)}$:- $\boxed{v^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)}\right) \le \frac{dv^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$v^{(6)}(t) \ge \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\nu_1)^{(6)}$

In the same manner , we get 413

$$v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \le v^{(6)}(t) \le (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \le \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \le v^{(6)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \le (\bar{\nu}_1)^{(6)} \le \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \le v^{(6)}(t) \le \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \le (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \le v^{(6)}(t) \le (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain
Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .
Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then
$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{d\nu^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)\nu^{(7)} - (a_{37})^{(7)}\nu^{(7)}$$

Definition of $\nu^{(7)}$:- $\quad \boxed{\nu^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right) \leq \frac{d\nu^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$\nu^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get 417

$$\nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)} e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \leq \nu^{(7)}(t) \leq (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \leq \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \leq \nu^{(7)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \leq (\bar{\nu}_1)^{(7)} \leq \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \leq \nu^{(7)}(t) \leq \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \leq (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also**

defines $(\nu_0)^{(8)}$ for the special case .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right) - (a''_{45})^{(9)}(T_{45},t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \le (\bar{\nu}_1)^{(9)} \le \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$, we obtain

$$(\nu_1)^{(9)} \le \nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)} \leq v^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{v^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $v^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\ ,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \quad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \quad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\ , \quad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \quad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(l) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows

that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G^*_{17} given by $\varphi((G_{19})^*) = 0$, T^*_{17} given by $f(T^*_{17}) = 0$ and 512

$$G^*_{16} = \frac{(a_{16})^{(2)}G^*_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G^*_{18} = \frac{(a_{18})^{(2)}G^*_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \quad 513$$

$$T^*_{16} = \frac{(b_{16})^{(2)}T^*_{17}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T^*_{18} = \frac{(b_{18})^{(2)}T^*_{17}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G^*_{21} given by $\varphi((G_{23})^*) = 0$, T^*_{21} given by $f(T^*_{21}) = 0$ and

$$G^*_{20} = \frac{(a_{20})^{(3)}G^*_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G^*_{22} = \frac{(a_{22})^{(3)}G^*_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

$$T^*_{20} = \frac{(b_{20})^{(3)}T^*_{21}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T^*_{22} = \frac{(b_{22})^{(3)}T^*_{21}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G^*_{25} given by $\varphi(G_{27}) = 0$, T^*_{25} given by $f(T^*_{25}) = 0$ and

$$G^*_{24} = \frac{(a_{24})^{(4)}G^*_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G^*_{26} = \frac{(a_{26})^{(4)}G^*_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

$$T^*_{24} = \frac{(b_{24})^{(4)}T^*_{25}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T^*_{26} = \frac{(b_{26})^{(4)}T^*_{25}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(\,G^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a_{13}')^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^*\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a_{14}')^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^*\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a_{15}')^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^*\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b_{13}')^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^*\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b_{14}')^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^*\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b_{15}')^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^*\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^*) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}(\,(G_{19})^*\,) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a_{16}')^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^*\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a_{17}')^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^*\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a_{18}')^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^*\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\big((b_{16}')^{(2)} - (r_{16})^{(2)}\big)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\big(s_{(16)(j)}T_{16}^*\mathbb{G}_j\big) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}\ and\ (b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \ , \ \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \ , \ \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}} (T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j} ((G_{43})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)} G_{40}^* \mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)} G_{41}^* \mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)} G_{42}^* \mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)} T_{40}^* \mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)} T_{41}^* \mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)} T_{42}^* \mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}} (T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j} ((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)} G_{44}^* \mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)} G_{45}^* \mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)} G_{46}^* \mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)} T_{44}^* \mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)} T_{45}^* \mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)} T_{46}^* \mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is

$$(\lambda)^{(1)}+(b'_{15})^{(1)}-(r_{15})^{(1)})\{((\lambda)^{(1)}+(a'_{15})^{(1)}+(p_{15})^{(1)})$$

$$\left[\left(((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)})(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)\right]$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(14)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(14)}T^{*}_{14}\right)$$

$$+\left(((\lambda)^{(1)}+(a'_{14})^{(1)}+(p_{14})^{(1)})(q_{13})^{(1)}G^{*}_{13}+(a_{13})^{(1)}(q_{14})^{(1)}G^{*}_{14}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(13)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(13)}T^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$\left(((\lambda)^{(1)})^{2}+(\,(b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)})\,(\lambda)^{(1)}\right)$$

$$+\left(((\lambda)^{(1)})^{2}+(\,(a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)})\,(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^{*}_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^{*}_{13}\right)$$

$$\left(((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)})s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)})\{((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)})$$

$$\left[\left(((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)})(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)})(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$\left(((\lambda)^{(2)})^{2}+(\,(b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)})\,(\lambda)^{(2)}\right)$$

$$+\left(((\lambda)^{(2)})^{2}+(\,(a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)})\,(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)})s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)})\{((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)})$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\Big)$$

$$+\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\Big)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\Big)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right]$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\Big)$$

$$+\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\Big)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\Big)$$

$$+\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\Big)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\Big)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^2+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^2+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^*_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^*_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^*_{37}+(b_{37})^{(7)}s_{(36),(38)}T^*_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^*_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^*_{41}+(b_{41})^{(8)}s_{(40),(41)}T^*_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^*_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^*_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^*_{41}+(b_{41})^{(8)}s_{(40),(40)}T^*_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^2+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^2+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^*_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^*_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^*_{41}+(b_{41})^{(8)}s_{(40),(42)}T^*_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^*_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^*_{45}+(b_{45})^{(9)}s_{(44),(45)}T^*_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^*_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^*_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^*_{45}+(b_{45})^{(9)}s_{(44),(44)}T^*_{44}\Big)$$

$$\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\Big)$$

$$\left(\left((\lambda)^{(9)}\right)^2 + \left((b'_{44})^{(9)} + (b'_{45})^{(9)} - (r_{44})^{(9)} + (r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2 + \left((a'_{44})^{(9)} + (a'_{45})^{(9)} + (p_{44})^{(9)} + (p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)} + (a'_{44})^{(9)} + (p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45} + (a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)} + (b'_{44})^{(9)} - (r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45} + (b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\} = 0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.


SECTION THIRTEEN

Electromagnetic, Weak, And Strong Nuclear Interactions, Particle Masses And Coupling Constants, W Bosons: Ad Pondus Omnium (Ad Pond Om)- Ad Referendum- Ab Ovo Usque Ad Mala


INTRODUCTION: VARIABLES TAKEN IN TO CONSIDERATION:

We take in to consideration that the following propositions to be true and give models:

(1) Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles **gives rise to** the Standard Model of particle physics
(2) Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles without the incorporation of gravity **is the reason for** particle masses and coupling constants, have the values that we measure?
(3) Comprehensive understanding of the relationship between the generations of the leptons may eventually explain the ratio of masses of the fundamental particles, and shed further light on the nature of mass generally from a quantum perspective and subtle modifications of precision electroweak observables that extra generations would induce **leads to** three generations of particles
(4) Matter **dissipates** antimatter in the universe which explains the predominance of the former
(5) Dark Matter **is(= eb or e) a** new particle
(6) HIG-13-003 Event 1: Event recorded with the CMS detector in 2012 at a proton-proton centre-of-mass energy of 8 TeV. The event shows characteristics expected from the decay of the SM Higgs boson to(eb) a pair of W bosons,
(7) Each of pair of W bosons subsequently decays(eb)to a muon (red lines)



HIG-13-002 Figure 1: Distribution of the four-lepton reconstructed mass for the sum of the 4e, 4μ, and 2e2μ channels. Points represent the data, shaded histograms represent the background and un-shaded histogram the signal expectations. The distributions are presented as stacked histograms. The measurements are presented for the sum of the data collected at centre-of-mass energies of 7 TeV and 8 TeV.



New CMS results at Moriond (Electroweak) 2013

NOTATION

Module One

G_{13} : Category one of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles

G_{14} : Category two of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles

G_{15} : Category three of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles

T_{13} : Category one of Standard Model of particle physics

T_{14} : Category two of Standard Model of particle physics

T_{15} : Category three of Standard Model of particle physics

Module Two

G_{16} : Category one of particle masses and coupling constants, have the values that we measure

G_{17} : Category two of particle masses and coupling constants, have the values that we measure

G_{18} : Category three of particle masses and coupling constants, have the values that we measure

T_{16} : Category one of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles without the incorporation of gravity

T_{17} : Category two of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles without the incorporation of gravity

T_{18} : Category three of Electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles without the incorporation of gravity

Module three

G_{20} : Category one of three generations of particles

G_{21} : Category two of three generations of particles

G_{22} : Category three of three generations of particles

T_{20} : Category one of Comprehensive understanding of the relationship between the generations of the leptons may eventually explain the ratio of masses of the fundamental particles, and shed further light on the nature of mass generally from a quantum perspective and subtle modifications of precision

electroweak observables that extra generations would induce

T_{21} : Category two of Comprehensive understanding of the relationship between the generations of the leptons may eventually explain the ratio of masses of the fundamental particles, and shed further light on the nature of mass generally from a quantum perspective and subtle modifications of precision electroweak observables that extra generations would induce

T_{22} : Category three of Comprehensive understanding of the relationship between the generations of the leptons may eventually explain the ratio of masses of the fundamental particles, and shed further light on the nature of mass generally from a quantum perspective and subtle modifications of precision electroweak observables that extra generations would induce

Module four

G_{24} : Category one of antimatter

G_{25} : Category two of antimatter

G_{26} : Category three of antimatter

T_{24} : Category one of Matter

T_{25} : Category two of Matter

T_{26} : Category three of Matter

Module five

G_{28} : Category one of new particle

G_{29} : Category two of new particle

G_{30} : Category three of new particle

T_{28} : Category one of Dark Matter

T_{29} : Category two of Dark Matter

T_{30} : Category three of Dark Matter

Module six

G_{32} : Category one of SM Higgs boson

G_{33} : Category two of SM Higgs boson

G_{34} : Category three of SM Higgs boson

T_{32} : Category one of pair of W bosons

T_{33} : Category two of pair of W bosons

T_{34} : Category three of pair of W bosons

Module seven

G_{36} : Category one of Each of pair of W bosons

G_{37} : Category two of Each of pair of W bosons

G_{38} : Category three of Each of pair of W bosons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{36} : Category one of muon (red lines)

T_{37} : Category two of muon (red lines)

T_{38} : Category three of muon (red lines) (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

Module eight

Each of pair of W bosons them **decays(eb) in to** undetected neutrino (the missing energy from the neutrino is shown by the red arrow).

G_{40} : Category one of Each of pair of W bosons

G_{41} : Category two of Each of pair of W bosons

G_{42} : Category three of Each of pair of W bosons

T_{40} : Category one of undetected neutrino (the missing energy from the neutrino is shown by the red arrow).

T_{41} : Category two of undetected neutrino (the missing energy from the neutrino is shown by the red arrow).

T_{42} : Category three of undetected neutrino (the missing energy from the neutrino is shown by the red arrow).

Module Nine

Each of pair of W bosons them decays (eb) in to undetected neutrino (the missing energy from the neutrino is shown by the red arrow) could also be(e) due to known Standard Model background processes

G_{44} : Category one of Standard Model background processes

G_{45} : Category two of Standard Model background processes

G_{46} : Category three of Standard Model background processes

T_{44} : Category one of Each of pair of W bosons them decays (eb) in to undetected neutrino (the missing energy from the neutrino is shown by the red arrow) (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{45} : Category two of Each of pair of W bosons them decays (eb) in to undetected neutrino (the missing energy from the neutrino is shown by the red arrow)

T_{46} : Category three of Each of pair of W bosons them decays (eb) in to undetected neutrino (the missing energy from the neutrino is shown by the red arrow)

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$

$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15}$ 6

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16}$ 7

$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17}$ 8

$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18}$ 9

$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\left((G_{19}),t\right)\right]T_{16}$ 10

$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\left((G_{19}),t\right)\right]T_{17}$ 11

$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\left((G_{19}),t\right)\right]T_{18}$ 12

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\left((G_{19}),t\right) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20}$ 13

$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21}$ 14

$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22}$ 15

$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20}$ 16

$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21}$ 17

$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22}$ 18

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24}$ 19

$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25}$ 20

$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26}$ 21

$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}),t\right)\right]T_{24}$ 22

$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}),t\right)\right]T_{25}$ 23

$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}),t\right)\right]T_{26}$ 24

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}),t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28}$ 25

$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29}$ 26

$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30}$ 27

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$
$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$
$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$
$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**
$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$
$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$
$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$
$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$
$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$
$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$
$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$
$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$
$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$
$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$
$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$
$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$
$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$
$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$
$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$
$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$
$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}),t)\right]T_{41} \quad 47$$
$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}),t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$
$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$
$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$
$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}),t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation

coefficient for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3
$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3
$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3
$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad (65)$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad (66)$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3
$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3
$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3
$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3
$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3
$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3
$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3
$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3
$-(b''_{44})^{(9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad (67)$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)} & +(a''_{20})^{(3)}(T_{21},t) & +(a''_{16})^{(2,2,2)}(T_{17},t) & +(a''_{13})^{(1,1,1,)}(T_{14},t) \\ +(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{36})^{(7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9)}(T_{45},t) \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)}\ \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\ \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\ \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\ \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\ \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\ \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\ \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\ \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\ \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\ \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\ \boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\ \boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\ \boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\ \boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\ \boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients*

for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) \quad +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[\begin{array}{ccc} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) \quad +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[\begin{array}{ccc} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) \quad +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{array}\right] G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* $1, 2\ 3$

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* $1, 2$ *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$ *are fourth augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$ *are fifth augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$ *are sixth augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$ *are seventh augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$ *are eighth augmentation coefficients for category* $1, 2$ *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients

for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)} & -(b''_{24})^{(4)}(G_{27},t) & -(b''_{28})^{(5,5,)}(G_{31},t) & -(b''_{32})^{(6,6,)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)} & -(b''_{25})^{(4)}(G_{27},t) & -(b''_{29})^{(5,5,)}(G_{31},t) & -(b''_{33})^{(6,6,)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)} & -(b''_{26})^{(4)}(G_{27},t) & -(b''_{30})^{(5,5,)}(G_{31},t) & -(b''_{34})^{(6,6,)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1)}(G,t) & -(b''_{18})^{(2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9)}(G_{47},t) \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3

$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3

$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3

$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3

$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3

$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3

$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3

$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)} & +(a''_{28})^{(5)}(T_{29},t) & +(a''_{24})^{(4,4,)}(T_{25},t) & +(a''_{32})^{(6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)} & +(a''_{29})^{(5)}(T_{29},t) & +(a''_{25})^{(4,4,)}(T_{25},t) & +(a''_{33})^{(6,6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)} & +(a''_{30})^{(5)}(T_{29},t) & +(a''_{26})^{(4,4,)}(T_{25},t) & +(a''_{34})^{(6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{30} \quad (81)$$

Where $+(a''_{28})^{(5)}(T_{29},t)$, $+(a''_{29})^{(5)}(T_{29},t)$, $+(a''_{30})^{(5)}(T_{29},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}\ ,\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}\ ,\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}\ ,\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}\ ,\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}\ ,\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}\ ,\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8\ ,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}\ ,\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}\ ,\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt}=(b_{28})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{28} \qquad 82$$

$$\frac{dT_{29}}{dt}=(b_{29})^{(5)}T_{28}-\left[\begin{array}{ccc}(b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{29} \qquad 83$$

$$\frac{dT_{30}}{dt}=(b_{30})^{(5)}T_{29}-\left[\begin{array}{ccc}(b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}\end{array}\right]T_{30} \qquad 84$$

$where\ \boxed{-(b''_{28})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{29})^{(5)}(G_{31},t)}\ ,\boxed{-(b''_{30})^{(5)}(G_{31},t)}$
$are\ first\ detrition\ coefficients\ \ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}\ ,\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}\ ,\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}\ are\ \ third\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)}\ ,\boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)}$ are fourth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)}\ ,\boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)}\ ,\boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)}\ ,\boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)}\ ,\boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1,2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39},t)} \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39},t)} \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second

detrition coefficients for category 1, 2 and 3

$-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)$, $-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a_{40}')^{(8)} & +(a_{40}'')^{(8)}(T_{41},t) & +(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{44}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a_{41}')^{(8)} & +(a_{41}'')^{(8)}(T_{41},t) & +(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{45}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a_{42}')^{(8)} & +(a_{42}'')^{(8)}(T_{41},t) & +(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a_{46}'')^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a_{40}'')^{(8)}(T_{41},t)$, $+(a_{41}'')^{(8)}(T_{41},t)$, $+(a_{42}'')^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)}\boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$lim_{T_2\to\infty}(a_i'')^{(1)}\,(T_{14},t)=(p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}\,(G,t)=\ \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)},(p_i)^{(1)},\ \ (r_i)^{(1)}}$ are positive constants and $\boxed{i=13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},$ are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\ ,\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)}+(a_i')^{(1)}+\ (\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)}+(b_i')^{(1)}+\ (\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le\left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le\ \ (r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}\,(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=\ \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with 116
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26$$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$ 118

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29}, t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}), t\big) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(5)}(T_{29}, t) = (p_i)^{(5)} \qquad 123$$

$$\lim_{G \to \infty}(b_i'')^{(5)}(G_{31}, t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}', t) - (a_i'')^{(5)}(T_{29}, t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})', t) - (b_i'')^{(5)}\big((G_{31}), t\big)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}', t)$

and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}<1$$

Definition of $(\hat{P}_{28})^{(5)},(\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)},(\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)},(a_i')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(p_i)^{(5)},\ (r_i)^{(5)},i=28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)}+(a_i')^{(5)}+\ (\hat{A}_{28})^{(5)}+\ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}]<1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)}+(b_i')^{(5)}+\ (\hat{B}_{28})^{(5)}+\ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}]<1$$

Where we suppose

$$(a_i)^{(6)},(a_i')^{(6)},(a_i'')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(b_i'')^{(6)}>0,\quad i,j=32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)},(b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t)\le(p_i)^{(6)}\le(\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t)\le\ (r_i)^{(6)}\le(b_i')^{(6)}\le(\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}),t\right)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\left((G_{35}),t\right)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\widehat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$$ 131

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$ 132

$$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$$

$$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It

is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+(\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)}+(b_i')^{(8)}+(\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)}+(a_i')^{(9)}+(\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)}+(b_i')^{(9)}+(\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}(\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0),T_i(0)$:

$$G_i(t)\le(\hat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le(\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0),T_i(0)$

$$G_i(t)\le(\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ G_i(0)=G_i^0>0$$

$$T_i(t)\le(\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\ T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le(\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ G_i(0)=G_i^0>0$$

$$T_i(t)\le(\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\ T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0),T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$$G_i(0) = G_i^0\ ,\ T_i(0) = T_i^0\ ,\ G_i^0 \le (\hat{P}_{20})^{(3)}\ , T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i\ ,\ T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\ ,\ T_i(0) = T_i^0\ ,\ G_i^0 \le (\hat{P}_{24})^{(4)}\ , T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{28})^{(5)}$, $T_i^0 \le (\hat{Q}_{28})^{(5)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\overline{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious 167 that

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that (172)

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that (174)

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

) The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$G_{13}^{(2)}|(a_{13}'')^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a_{13}'')^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a_{13}')^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a_{13}'')^{(1)}$ and $(b_{13}'')^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b_i')^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a_{14}')^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\ (G(t),t))=(b_{14}')^{(1)}$ then $T_{14}\ \to\infty$. (189)

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1,T_{13}\ (t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$

$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205

unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b''_{18})^{(2)}\left((G_{19})(t),t\right)=(b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}<1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 215

$(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\leq \tag{226}$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2 \; and \; \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25} \; and \; G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25} \; or \; G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24} \; is$ bounded, from below, the same property holds for $G_{25} \; and \; G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (239)

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad (245)$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. 249

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 250

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 261

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{40})^{(8)}|G_{41}^{(1)}-G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}|(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})-(a''_{40})^{(8)}(T_{41}^{(2)},s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$|(G_{43})^{(1)}-(G_{43})^{(2)}|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)})d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(8)}t)}>0 \quad \text{for t} > 0$$

Definition of $((\widehat{M}_{40})^{(8)})_1,((\widehat{M}_{40})^{(8)})_2$ *and* $((\widehat{M}_{40})^{(8)})_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40}<(\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt}\le((\widehat{M}_{40})^{(8)})_1-(a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})} \right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{44})^{(9)} \left|G_{45}^{(1)} - G_{45}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\, ds_{(44)} +$$
$$\int_0^t \{(a'_{44})^{(9)} \left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$G_{44}^{(2)} \left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\} ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)}\right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ *and* $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44,45,46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(9)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$ are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, $\boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, $\boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, $\boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, $\boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16} e^{((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)})\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16} e^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$

$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

332

$$G^0_{24} e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24} e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)}-(a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)},$ ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)},$ ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)},$ ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)},$ ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)},(S_2)^{(5)},(R_1)^{(5)},(R_2)^{(5)}$:- 348

Where $(S_1)^{(5)}=(a_{28})^{(5)}(m_2)^{(5)}-(a'_{28})^{(5)}$

$(S_2)^{(5)}=(a_{30})^{(5)}-(p_{30})^{(5)}$

$(R_1)^{(5)}=(b_{28})^{(5)}(\mu_2)^{(5)}-(b'_{28})^{(5)}$

$(R_2)^{(5)}=(b'_{30})^{(5)}-(r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$:

$(\sigma_1)^{(6)},(\sigma_2)^{(6)},(\tau_1)^{(6)},(\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)}\le-(a'_{32})^{(6)}+(a'_{33})^{(6)}-(a''_{32})^{(6)}(T_{33},t)+(a''_{33})^{(6)}(T_{33},t)\le-(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)}\le-(b'_{32})^{(6)}+(b'_{33})^{(6)}-(b''_{32})^{(6)}\big((G_{35}),t\big)-(b''_{33})^{(6)}\big((G_{35}),t\big)\le-(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)},(v_2)^{(6)},(u_1)^{(6)},(u_2)^{(6)},v^{(6)},u^{(6)}$: 350

By $(v_1)^{(6)}>0$, $(v_2)^{(6)}<0$ and respectively $(u_1)^{(6)}>0$, $(u_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2+(\sigma_1)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2+(\tau_1)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$ and

Definition of $(\bar{v}_1)^{(6)},,(\bar{v}_2)^{(6)},(\bar{u}_1)^{(6)},(\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)}>0$, $(\bar{v}_2)^{(6)}<0$ and respectively $(\bar{u}_1)^{(6)}>0$, $(\bar{u}_2)^{(6)}<0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2+(\sigma_2)^{(6)}v^{(6)}-(a_{32})^{(6)}=0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2+(\tau_2)^{(6)}u^{(6)}-(b_{32})^{(6)}=0$

Definition of $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)},(v_0)^{(6)}$:-

If we define $(m_1)^{(6)},(m_2)^{(6)},(\mu_1)^{(6)},(\mu_2)^{(6)}$ by

$(m_2)^{(6)}=(v_0)^{(6)},(m_1)^{(6)}=(v_1)^{(6)}$, $\boldsymbol{if}\ (v_0)^{(6)}<(v_1)^{(6)}$

$(m_2)^{(6)}=(v_1)^{(6)},(m_1)^{(6)}=(\bar{v}_6)^{(6)}$, $\boldsymbol{if}\ (v_1)^{(6)}<(v_0)^{(6)}<(\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)}=\frac{G_{32}^{0}}{G_{33}^{0}}}$

$(m_2)^{(6)}=(v_1)^{(6)},(m_1)^{(6)}=(v_0)^{(6)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(6)}<(v_0)^{(6)}$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0$, $(v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0$, $(\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G^0_{36} e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t}\leq G_{38}(t)\leq\right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\leq T_{38}(t)\leq \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$$

$$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$$

$$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)}\leq-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\leq-(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)}\leq-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\leq-(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0$, $(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0$, $(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0$, $(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0$, $(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}$, $\boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}$, $\boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)}$,

and $\boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, $\boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, $\boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}\left((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)}\right)}\left[e^{\left((S_1)^{(8)} - (p_{40})^{(8)}\right)t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}\left((S_1)^{(8)} - (a'_{42})^{(8)}\right)}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}\left((R_1)^{(8)} - (b'_{42})^{(8)}\right)}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}\left((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)}\right)}\left[e^{\left((R_1)^{(8)} + (r_{40})^{(8)}\right)t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$

Definition of $(\nu_1)^{(9)}, (\nu_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, \nu^{(9)}, u^{(9)}$:

By $(\nu_1)^{(9)} > 0$, $(\nu_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{\nu}_1)^{(9)}, , (\bar{\nu}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{\nu}_1)^{(9)} > 0$, $(\bar{\nu}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(\nu^{(9)}\big)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (\nu_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (\nu_0)^{(9)}, (m_1)^{(9)} = (\nu_1)^{(9)}$, $\boldsymbol{if}\ (\nu_0)^{(9)} < (\nu_1)^{(9)}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\bar{\nu}_1)^{(9)}$, $\boldsymbol{if}\ (\nu_1)^{(9)} < (\nu_0)^{(9)} < (\bar{\nu}_1)^{(9)}$,
and $\boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (\nu_1)^{(9)}, (m_1)^{(9)} = (\nu_0)^{(9)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(9)} < (\nu_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \leq \frac{d\nu^{(1)}}{dt} \leq -\left((a_{14})^{(1)}(\nu^{(1)})^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also

defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $v^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq v^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)\le\frac{dv^{(4)}}{dt}\le-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_4)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)},(v_0)^{(4)}$:-

For $0<\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}<(v_1)^{(4)}<(\bar{v}_1)^{(4)}$

$$v^{(4)}(t)\ge\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}\quad,\quad\boxed{(C)^{(4)}=\frac{(v_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(v_2)^{(4)}}}$$

it follows $(v_0)^{(4)}\le v^{(4)}(t)\le(v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(4)}=\frac{(\bar{v}_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)}\le v^{(4)}(t)\le(\bar{v}_1)^{(4)}$

If $0<(v_1)^{(4)}<(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}<(\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)}\le\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}\le v^{(4)}(t)\le$$

$$\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(\bar{v}_1)^{(4)}$$

If $0<(v_1)^{(4)}\le(\bar{v}_1)^{(4)}\le\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)}\le v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)}\le v^{(4)}(t)\le(m_1)^{(4)},\quad\boxed{v^{(4)}(t)=\frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)}\le u^{(4)}(t)\le(\mu_1)^{(4)},\quad\boxed{u^{(4)}(t)=\frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \le \frac{dv^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \ge \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \le v^{(7)}(t) \le (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \le \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \le v^{(7)}(t) \le (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \le \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \le v^{(7)}(t) \le$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \le (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \le (\bar{v}_1)^{(7)} \le \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \le v^{(7)}(t) \le \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \le (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \le v^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:-

420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain

421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get

422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case,

423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0$ 477

$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0$ 478

$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0$ 479

$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0$ 480

$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0$ 481

$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0$ 482

$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0$ 483

$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0$ 484

$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0$ 484 A

$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$

$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$

$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(m) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502
$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{\left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T^*_{14})\right]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{\left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T^*_{14})\right]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{\left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)\right]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{\left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)\right]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \tag{513}$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*} = \frac{(b_{44})^{(9)} T_{45}^{*}}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*} = \frac{(b_{46})^{(9)} T_{45}^{*}}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial (a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b''_i)^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^* \,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\,(G_{31})^*\,) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(6)}$ *and* $(b''_i)^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b''_i)^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(7)}$ *and* $(b''_i)^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\right)\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\Big(\big((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\big)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\Big)$$

$$\Big(\big((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\big)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\Big)$$

$$\Big(\big((\lambda)^{(8)}\big)^{2}+\big((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\big)(\lambda)^{(8)}\Big)$$

$$\Big(\big((\lambda)^{(8)}\big)^{2}+\big((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\big)(\lambda)^{(8)}\Big)$$

$$+\Big(\big((\lambda)^{(8)}\big)^{2}+\big((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\big)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\big((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\big)\big((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\big)$$

$$\Big(\big((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\big)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\big((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\big)$$

$$\Big[\Big(\big((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\big)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\Big)\Big]$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\Big(\big((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\big)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\Big(\big((\lambda)^{(9)}\big)^{2}+\big((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\big)(\lambda)^{(9)}\Big)$$

$$\Big(\big((\lambda)^{(9)}\big)^{2}+\big((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\big)(\lambda)^{(9)}\Big)$$

$$+\Big(\big((\lambda)^{(9)}\big)^{2}+\big((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\big)(\lambda)^{(9)}\Big)(q_{46})^{(9)}G_{46}$$

$$+\big((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\big)\big((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\big)$$

$$\Big(\big((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\big)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\Big)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.


SECTION FOURTEEN

Bekenstein–Hawking Entropy, A Muon (Red Lines On The Right) And Neutrinos, Charged Hadron And A Neutrino, B-Quarks And An LSP, Etal.: Felix Qui Potuit Rerum Cognoscere Causas - Forsan Et Haec Olim Meminisse Iuvabit Model

INTRODUCTION—VARIABLES USED


CERN WEB SITE NEWS AND BLOGS RECEIVED

(1) The formula for the Bekenstein–Hawking entropy (S) of a black hole, which depends on the area of the black hole (A). The constants are the speed of light (c), the Boltzmann constant (k), Newton's constant (G), and the reduced Planck constant (ħ). $$S = \frac{1}{4}\frac{c^3 k}{G\hbar}A$$




HIG-12-004 Event 1: Event recorded with the CMS detector in 2012 at a proton-proton centre-of-mass energy of 8 TeV.

(2) The event shows characteristics expected from the decay of the SM Higgs boson **to(eb) a** pair of τ leptons. Such an event is characterised **by (eb) the production of** two forward-going jets, seen here in opposite end caps.
(3) One of the τs decays(eb) to a muon (red lines on the right) and neutrinos
(4) Other τ **decays(eb) into** a charged hadron and a neutrino.




(5) SUS-12-024 Figure 1: In a model where the gluino **decays(eb) t**o two b-quarks and an LSP, CMS has excluded gluinos with a mass up to 1.15 TeV, for LSP masses up to 500 GeV.




(6) EXO-12-026 Figure 1: Distribution of energy loss (Y axis) versus particle momentum (X axis) for data, together with the Monte Carlo (MC) simulation of HSCP candidates with single,

fractional, and multiple charges. The separation expected between HSCP candidates and typical events **provides (eb) a** powerful discriminator for the search.

Katzenjammerian slovenly inept schleps, schlock trash, sucker and hookers, spatio temporal spallationalistic spalpleen, swindler mountebank who would not hesitate to use God's name to sell themselves, badgering badinage who put forward women as shields, lollypop hoodlum lowly gags, nocturnal fossa cryptoprocta ferox fossickers, looming Looney, mealy mouthed meddle, murky monads, corybantic crows, muttatis mutandis mutton heads, are some of the epithets , alliterations and metaphors that could be used for such fellows. God made them to suffer and they shall!



(7) EXO-12-026 Figure 2: Lower limits on the mass of new fractionally or multiply charged particles. The plot shows the lower mass limit (95% confidence level) versus electric charge. Results are compared also with previous searches.



(8) EXO-12-060 Figure 1: The transverse mass distribution of events with one electron and missing energy. The points with error bars represent the data. The histograms represent the expectations from various Standard Model processes, including 'di-boson' contributions from WW, ZZ and WZ. Simulated signal distributions for W′ masses of 0.5 and 2.5 TeV are also shown.

Experimental Test of Local Hidden-Variable Theories: Edward S. Fry and Randall C. Thompson

(9) Authors have measured the linear polarization correlation between the two photons from the 73S1→63P1→61S0 cascade of Hg200. The results were used to evaluate Freedman's version of the Bell inequality, δ<~0. Our result is δexp=+0.046±0.014, in **clear violation of the** inequality and in excellent agreement with the quantum mechanical prediction, δQM=+0.044±0.007. An important feature of the experiment was the explicit measurement of the initial density matrix for the cascading atoms. © 1976 The American Physical Society URL: http://link.aps.org/doi/10.1103/PhysRevLett.37.465 DOI: 10.1103/PhysRevLett.37.465

NOTATION

Module One

$$S = \frac{1}{4}\frac{c^3 k}{G\hbar}A$$

G_{13} : Category one of RHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

G_{14} : Category two of RHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

G_{15} : Category three of RHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

T_{13} : Category one of LHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

T_{14} : Category two of LHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

T_{15} : Category three of LHS of the above equation (Classification is based on the characterstics of the systems that study the relation and are under investigation)

Module Two

G_{16} : Category one of SM Higgs boson

G_{17} : Category two of SM Higgs boson

G_{18} : Category three of SM Higgs boson

T_{16} : Category one of pair of τ leptons (characterised by forward-going jets, seen here in opposite end caps).

T_{17} : Category two of pair of τ leptons (characterised by forward-going jets, seen here in opposite end caps).

T_{18} : Category three of pair of τ leptons (characterised by forward-going jets, seen here in opposite end caps).

Module three

G_{20} : Category one of One of the τs decays (There are lot of systems and so there are lot of 'one of taus decaying)

G_{21} : Category two of One of the τs decays(There are lot of systems and so there are lot of 'one of taus decaying)

G_{22} : Category three of One of the τs decays(There are lot of systems and so there are lot of 'one of taus decaying)

T_{20} : Category one of muon (red lines on the right) and neutrinos

T_{21} : Category two of muon (red lines on the right) and neutrinos

T_{22} : Category three of muon (red lines on the right) and neutrinos

Module four

G_{24} : Category one of Other τ decays

G_{25} : Category two of Other τ decays

G_{26} : Category three of Other τ decays

T_{24} : Category one of charged hadron and a neutrino (For various systems there are very many such charged hadron and neutrino)

T_{25} : Category two of charged hadron and a neutrino

T_{26} : Category three of charged hadron and a neutrino

Module five

G_{28} : Category one of gluino (Lot of investigating systems would have gluinos and they shall decay. We shall utilize the characterstics of the systems under consideration to be fundamental basis for the classification purposes)

G_{29} : Category two of gluino (Lot of investigating systems would have gluinos and they shall decay. We shall utilize the characterstics of the systems under consideration to be fundamental basis for the classification purposes)

G_{30} : Category three of gluino (Lot of investigating systems would have gluinos and they shall decay. We shall utilize the characterstics of the systems under consideration to be fundamental basis for the classification purposes)

T_{28} : Category one of two b-quarks and an LSP, CMS has excluded gluinos with a mass up to 1.15 TeV, for LSP masses up to 500 GeV

T_{29} : Category two of two b-quarks and an LSP, CMS has excluded gluinos with a mass up to 1.15 TeV, for LSP masses up to 500 GeV

T_{30} : Category three of two b-quarks and an LSP, CMS has excluded gluinos with a mass up to 1.15 TeV, for LSP masses up to 500 GeV

Module six

G_{32} : Category one of separation expected between HSCP candidates and typical events

G_{33} : Category two of separation expected between HSCP candidates and typical events

G_{34} : Category three of separation expected between HSCP candidates and typical events

T_{32} : Category one of powerful discriminator for the search

T_{33} : Category two of powerful discriminator for the search

T_{34} : Category three of powerful discriminator for the search

Module seven

G_{36} : Category one of lower mass limit (95% confidence level)

G_{37} : Category two of lower mass limit (95% confidence level)

G_{38} : Category three of lower mass limit (95% confidence level)

T_{36} : Category one of electric charge

T_{37} : Category two of electric charge

T_{38} : Category three of electric charge

Module eight

G_{40} : Category one of transverse mass distribution of events with one electron and missing energy

G_{41} : Category two of transverse mass distribution of events with one electron and missing energy

G_{42} : Category three of transverse mass distribution of events with one electron and missing energy

T_{40} : Category one of expectations from various Standard Model processes, including 'di-boson' contributions from WW, ZZ and WZ

T_{41} : Category two of expectations from various Standard Model processes, including 'di-boson' contributions from WW, ZZ and WZ

T_{42} : Category three of expectations from various Standard Model processes, including 'di-boson' contributions from WW, ZZ and WZ

Module Nine

G_{44} : Category one of violates Bell's inequalities but is consonance with the quantum mechanical projection, prediction, and prognostication

G_{45} : Category two of violates Bell's inequalities but is consonance with the quantum mechanical projection, prediction, and prognostication

G_{46} : Category three of violates Bell's inequalities but is consonance with the quantum mechanical projection, prediction, and prognostication (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

T_{44} : Category one of measurement of the linear polarization correlation between the two photons from the 73S1→63P1→61S0 cascade of Hg200

T_{45} : Category two of measurement of the linear polarization correlation between the two photons from the 73S1→63P1→61S0 cascade of Hg200

T_{46} : Category three of measurement of the linear polarization correlation between the two photons from the 73S1→63P1→61S0 cascade of Hg200

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}, (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big)\right]T_{16} \quad 10$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}), t\big)\right]T_{17} \quad 11$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}), t\big)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor
$-(b''_{16})^{(2)}\big((G_{19}), t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27}), t\right)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27}), t\right)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27}), t\right)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}\left((G_{27}), t\right) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}\left((G_{31}), t\right)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}\left((G_{31}), t\right)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}\left((G_{31}), t\right)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}\left((G_{31}), t\right) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}\left((G_{35}), t\right)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}\left((G_{35}), t\right)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}\left((G_{35}),t\right)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}\left((G_{39}),t\right)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}\left((G_{39}),t\right)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}\left((G_{39}),t\right)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}\left((G_{43}),t\right)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & +(a''_{13})^{(1)}(T_{14},t) & +(a''_{16})^{(2,2,)}(T_{17},t) & +(a''_{20})^{(3,3,)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{36})^{(7,7)}(T_{37},t) & +(a''_{40})^{(8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & +(a''_{14})^{(1)}(T_{14},t) & +(a''_{17})^{(2,2,)}(T_{17},t) & +(a''_{21})^{(3,3,)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,)}(T_{33},t) & \\ +(a''_{37})^{(7,7)}(T_{37},t) & +(a''_{41})^{(8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{37})^{(7,7,)}(G_{39},t)$, $-(b''_{36})^{(7,7,)}(G_{39},t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8)}(G_{43},t)$ $-(b''_{41})^{(8,8)}(G_{43},t)$ $-(b''_{42})^{(8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt}=(a_{16})^{(2)}G_{17}-\begin{bmatrix}(a'_{16})^{(2)} & +(a''_{16})^{(2)}(T_{17},t) & +(a''_{13})^{(1,1,)}(T_{14},t) & +(a''_{20})^{(3,3,3)}(T_{21},t)\\ +(a''_{24})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{36})^{(7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9)}(T_{45},t)\end{bmatrix}G_{16} \quad (61)$$

$$\frac{dG_{17}}{dt}=(a_{17})^{(2)}G_{16}-\begin{bmatrix}(a'_{17})^{(2)} & +(a''_{17})^{(2)}(T_{17},t) & +(a''_{14})^{(1,1,)}(T_{14},t) & +(a''_{21})^{(3,3,3)}(T_{21},t)\\ +(a''_{25})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{37})^{(7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9)}(T_{45},t)\end{bmatrix}G_{17} \quad (62)$$

$$\frac{dG_{18}}{dt}=(a_{18})^{(2)}G_{17}-\begin{bmatrix}(a'_{18})^{(2)} & +(a''_{18})^{(2)}(T_{17},t) & +(a''_{15})^{(1,1,)}(T_{14},t) & +(a''_{22})^{(3,3,3)}(T_{21},t)\\ +(a''_{26})^{(4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6)}(T_{33},t)\\ +(a''_{38})^{(7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9)}(T_{45},t)\end{bmatrix}G_{18} \quad (63)$$

Where $+(a''_{16})^{(2)}(T_{17},t)$, $+(a''_{17})^{(2)}(T_{17},t)$, $+(a''_{18})^{(2)}(T_{17},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt}=(b_{16})^{(2)}T_{17}-\begin{bmatrix}(b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t)\\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t)\\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t)\end{bmatrix}T_{16} \quad (64)$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation

coefficients for category 1, 2 and 3
$+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficients for category 1, 2 and 3
$+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficients for category 1, 2 and 3
$+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficients for category 1, 2 and 3
$+(a''_{36})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1, 2 and 3
$+(a''_{40})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8)}(T_{41},t)$ are eight augmentation coefficients for category 1, 2 and 3
$+(a''_{44})^{(9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\left[-(b''_{20})^{(3)}(G_{23},t)\right] & \left[-(b''_{16})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{13})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\left[-(b''_{21})^{(3)}(G_{23},t)\right] & \left[-(b''_{17})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{14})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\left[-(b''_{22})^{(3)}(G_{23},t)\right] & \left[-(b''_{18})^{(2,2,2)}(G_{19},t)\right] & \left[-(b''_{15})^{(1,1,1,)}(G,t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*
$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*
$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*
$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*
$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*
$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*
$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3
$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients

for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt}=(a_{24})^{(4)}G_{25}-\begin{bmatrix}(a'_{24})^{(4)}\boxed{+(a''_{24})^{(4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{24} \quad (73)$$

$$\frac{dG_{25}}{dt}=(a_{25})^{(4)}G_{24}-\begin{bmatrix}(a'_{25})^{(4)}\boxed{+(a''_{25})^{(4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6)}(T_{33},t)}\\ \boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{25} \quad (74)$$

$$\frac{dG_{26}}{dt}=(a_{26})^{(4)}G_{25}-\begin{bmatrix}(a'_{26})^{(4)}\boxed{+(a''_{26})^{(4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}\\ \boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}\\ \boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}\end{bmatrix}G_{26} \quad (75)$$

$\boxed{(a''_{24})^{(4)}(T_{25},t)}$, $\boxed{(a''_{25})^{(4)}(T_{25},t)}$, $\boxed{(a''_{26})^{(4)}(T_{25},t)}$ *are first augmentation coefficients category* 1, 2 3

$\boxed{+(a''_{28})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,)}(T_{29},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1)}(T_{14},t)}$
are fourth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{16})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2)}(T_{17},t)}$
are fifth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{20})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3)}(T_{21},t)}$
are sixth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)}$
are seventh augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)}$
are eighth augmentation coefficients for category 1, 2 *and* 3

$\boxed{+(a''_{46})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9)}(T_{45},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt}=(b_{24})^{(4)}T_{25}-\begin{bmatrix}(b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt}=(b_{25})^{(4)}T_{24}-\begin{bmatrix}(b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt}=(b_{26})^{(4)}T_{25}-\begin{bmatrix}(b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}\\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}\end{bmatrix}T_{26} \quad (78)$$

$Where\ \boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}\ are\ first\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}\ are\ second\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}\ are\ third\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$

$are\ fourth\ \ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$

$are\ fifth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$

$are\ sixth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$

$are\ seventh\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$

$are\ eighth\ detrition\ coefficients\ for\ category\ 1,2\ and\ 3$

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{28} \quad 79$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad 80$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad 81$$

$Where\ \boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}\ are\ first\ augmentation\ coefficients\ for\ category\ 1,2\ and\ 3$

$And\ \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}\ are\ second\ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}\ are\ third\ \ augmentation\ coefficient\ for\ category\ 1,2\ and\ 3$

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh

augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad (84)$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \quad (85)$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)} & +(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ & +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* $1, 2$ *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}, \boxed{-(b''_{33})^{(6)}(G_{35},t)}, \boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}, \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \qquad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \qquad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \qquad 93$$

$$= (a_{38})^{(7)}G_{37} - \begin{bmatrix} (a'_{38})^{(7)}\boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}, \boxed{(a''_{37})^{(7)}(T_{37},t)}, \boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for

category 1, 2 and 3

$+(a_{16}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{17}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$, $+(a_{18}'')^{(2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a_{20}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{21}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$, $+(a_{22}'')^{(3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a_{24}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{25}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$, $+(a_{26}'')^{(4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a_{28}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{29}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$, $+(a_{30}'')^{(5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a_{32}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{33}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$, $+(a_{34}'')^{(6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{13}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$, $+(a_{15}'')^{(1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a_{40}'')^{(8,8,8,8,8,8,8,8,)}(T_{41},t)$ are eighth augmentation coefficient for 1,2,3

$+(a_{46}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{45}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$, $+(a_{44}'')^{(9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \qquad 94$$

$$(b_{36})^{(7)}T_{37} - \begin{bmatrix} (b_{36}')^{(7)}\boxed{-(b_{36}'')^{(7)}(G_{39},t)} & \boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{32}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{13}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{44}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b_{37}')^{(7)}\boxed{-(b_{37}'')^{(7)}(G_{39},t)} & \boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{33}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{14}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{45}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b_{38}')^{(7)}\boxed{-(b_{38}'')^{(7)}(G_{39},t)} & \boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b_{34}'')^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b_{15}'')^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b_{46}'')^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)} & +(a''_{40})^{(8)}(T_{41},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)}\boxed{-(b''_{40})^{(8)}(G_{43},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$
$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)}\boxed{-(b''_{41})^{(8)}(G_{43},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)}\boxed{-(b''_{42})^{(8)}(G_{43},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)} G_{45} - \begin{bmatrix} (a'_{44})^{(9)} \boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)} G_{44} - \begin{bmatrix} (a'_{45})^{(9)} \boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)} G_{45} - \begin{bmatrix} (a'_{46})^{(9)} \boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)} T_{45} - \begin{bmatrix} (b'_{44})^{(9)} \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{14}$$
$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a''_i)^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b''_i)^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}', t) - (a_i'')^{(1)}(T_{14}, t)| \leq (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G', t) - (b_i'')^{(1)}(G, t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}', t)$ and $(a_i'')^{(1)}(T_{14}, t)$. (T_{14}', t) and (T_{14}, t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \ , \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}, \ (r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)} + (a_i')^{(1)} + \ (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)} + (b_i')^{(1)} + \ (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i, j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq \left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq \ (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}\left((G_{19}), t\right) = \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}',t)-(a_i'')^{(2)}(T_{17},t)|\leq(\hat{k}_{16})^{(2)}|T_{17}-T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})',t)-(b_i'')^{(2)}\big((G_{19}),t\big)|<(\hat{k}_{16})^{(2)}||(G_{19})-(G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}',t)$ and$(a_i'')^{(2)}(T_{17},t)$. (T_{17}',t) and (T_{17},t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)},(\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)}=1$ then the function $(a_i'')^{(2)}(T_{17},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},$ are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}},\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}<1$$

Definition of $(\hat{P}_{13})^{(2)},(\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)},(\hat{k}_{16})^{(2)},(\hat{A}_{16})^{(2)}and\ (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)},(a_i')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(p_i)^{(2)},\ (r_i)^{(2)},i=16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)}+(a_i')^{(2)}+(\hat{A}_{16})^{(2)}+(\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)}+(b_i')^{(2)}+(\hat{B}_{16})^{(2)}+(\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}]<1 \quad 111$$

Where we suppose

$$(a_i)^{(3)},(a_i')^{(3)},(a_i'')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(b_i'')^{(3)}>0,\quad i,j=20,21,22 \quad 112$$

The functions $(a_i'')^{(3)},(b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)},\ (\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21},t)\leq(p_i)^{(3)}\leq(\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23},t)\leq\ (r_i)^{(3)}\leq(b_i')^{(3)}\leq(\hat{B}_{20})^{(3)}$$

$$lim_{T_2\to\infty}(a_i'')^{(3)}(T_{21},t)=(p_i)^{(3)} \quad 113$$

$$\lim_{G\to\infty}(b_i'')^{(3)}(G_{23},t)=\ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)},(\hat{B}_{20})^{(3)},(p_i)^{(3)},\ (r_i)^{(3)}}$ are positive constants and $\boxed{i=20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}',t)-(a_i'')^{(3)}(T_{21},t)|\leq(\hat{k}_{20})^{(3)}|T_{21}-T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}{}',t)-(b_i'')^{(3)}(G_{23},t)|<(\hat{k}_{20})^{(3)}||G_{23}-G_{23}{}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}',t)$ and$(a_i'')^{(3)}(T_{21},t)$. (T_{21}',t) And (T_{21},t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)},(\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)}=1$ then the function $(a_i'')^{(3)}(T_{21},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}},\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)},(\hat{k}_{20})^{(3)},(\hat{A}_{20})^{(3)}and\ (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)},(a_i')^{(3)},(b_i)^{(3)},(b_i')^{(3)},(p_i)^{(3)},\ (r_i)^{(3)},i=20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)}+(a_i')^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)}+(b_i')^{(3)}+(\hat{B}_{20})^{(3)}+(\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}]<1$$

Where we suppose

$$(a_i)^{(4)},(a_i')^{(4)},(a_i'')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(b_i'')^{(4)}>0,\quad i,j=24,25,26 \qquad 117$$

The functions $(a_i'')^{(4)},(b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)},\ (r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25},t)\leq(p_i)^{(4)}\leq(\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}),t\big)\leq\ (r_i)^{(4)}\leq(b_i')^{(4)}\leq(\hat{B}_{24})^{(4)}$$

$$lim_{T_2\to\infty}(a_i'')^{(4)}(T_{25},t)=(p_i)^{(4)} \qquad 118$$

$$\lim_{G\to\infty}(b_i'')^{(4)}\big((G_{27}),t\big)=\ (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)},(\hat{B}_{24})^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}}$ are positive constants and $\boxed{i=24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}',t)-(a_i'')^{(4)}(T_{25},t)|\leq(\hat{k}_{24})^{(4)}|T_{25}-T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It

is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{24})^{(4)}=1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$: 120

$(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}},\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}<1$$

Definition of $(\hat{P}_{24})^{(4)},(\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\widehat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)},(\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)},(a_i')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(p_i)^{(4)},\ (r_i)^{(4)}, i=24,25,26$, satisfy the inequalities

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\,(a_i)^{(4)}+(a_i')^{(4)}+\ (\hat{A}_{24})^{(4)}+\ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}]<1$$

$$\frac{1}{(\widehat{M}_{24})^{(4)}}[\ (b_i)^{(4)}+(b_i')^{(4)}+\ (\hat{B}_{24})^{(4)}+\ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}]<1$$

Where we suppose

$$(a_i)^{(5)},(a_i')^{(5)},(a_i'')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(b_i'')^{(5)}>0,\quad i,j=28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)},(b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)},\ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t)\le(p_i)^{(5)}\le(\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big)\le\ (r_i)^{(5)}\le(b_i')^{(5)}\le(\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}\,(T_{29},t)=(p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}\,(G_{31},t)=\ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)},(p_i)^{(5)},\ (r_i)^{(5)}}$ are positive constants and $\boxed{i=28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t)-(a_i'')^{(5)}(T_{29},t)|\le(\hat{k}_{28})^{(5)}|T_{29}-T_{29}'|e^{-(\widehat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t)-(b_i'')^{(5)}\big((G_{31}),t\big)|<(\hat{k}_{28})^{(5)}||(G_{31})-(G_{31})'||e^{-(\widehat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and $(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\big[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\big]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)} (\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i, j = 32,33,34$$ 127

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33}, t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}), t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2 \to \infty} (a_i'')^{(6)}(T_{33}, t) = (p_i)^{(6)}$$ 128

$$\lim_{G \to \infty} (b_i'')^{(6)}((G_{35}), t) = (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)}, (\hat{B}_{32})^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}}$ are positive constants and $\boxed{i = 32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}', t) - (a_i'')^{(6)}(T_{33}, t)| \le (\hat{k}_{32})^{(6)} |T_{33} - T_{33}'| e^{-(\widehat{M}_{32})^{(6)} t}$$

$$|(b_i'')^{(6)}((G_{35})', t) - (b_i'')^{(6)}((G_{35}), t)| < (\hat{k}_{32})^{(6)} ||(G_{35}) - (G_{35})'|| e^{-(\widehat{M}_{32})^{(6)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}', t)$ and $(a_i'')^{(6)}(T_{33}, t)$. (T_{33}', t) and (T_{33}, t) are points belonging to the interval $[(\hat{k}_{32})^{(6)}, (\widehat{M}_{32})^{(6)}]$. It is to be noted that $(a_i'')^{(6)}(T_{33}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)} = 1$ then the function $(a_i'')^{(6)}(T_{33}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\widehat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)}(\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

$$(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38 \qquad 131$$

The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)} \qquad 132$$

$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42$ 136

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)}$ 138

$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)}$ 139

$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)}$ 140

$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t}$ 142

$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It is to be noted that $(a_i'')^{(8)}(T_{41}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}} \ , \frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, \ (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)} (\hat{k}_{40})^{(8)}] < 1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i, j = 44,45,46 \qquad 146\text{A}$$

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}, \ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45}, t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47}, t) \leq \ (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$$

$$\lim_{T_2 \to \infty} (a_i'')^{(9)}(T_{45}, t) = (p_i)^{(9)}$$

$$\lim_{G \to \infty} (b_i'')^{(9)}(G_{47}, t) = \ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)}, (\hat{B}_{44})^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}}$ are positive constants and $\boxed{i = 44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}', t) - (a_i'')^{(9)}(T_{45}, t)| \leq (\hat{k}_{44})^{(9)} |T_{45} - T_{45}'| e^{-(\hat{M}_{44})^{(9)} t}$$

$$|(b_i'')^{(9)}((G_{47})', t) - (b_i'')^{(9)}((G_{47}), t)| < (\hat{k}_{44})^{(9)} ||(G_{47}) - (G_{47})'|| e^{-(\hat{M}_{44})^{(9)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}', t)$ and $(a_i'')^{(9)}(T_{45}, t)$. (T_{45}', t) and (T_{45}, t) are points belonging to the interval $[(\hat{k}_{44})^{(9)}, (\hat{M}_{44})^{(9)}]$. It is to be noted that $(a_i'')^{(9)}(T_{45}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)} = 1$ then the function $(a_i'')^{(9)}(T_{45}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} \ , \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$$

Definition of $(\hat{P}_{44})^{(9)}, (\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)}, (\hat{k}_{44})^{(9)}, (\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)}, (a_i')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (p_i)^{(9)}, \ (r_i)^{(9)}, i = 44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(a_i)^{(9)} + (a_i')^{(9)} + (\hat{A}_{44})^{(9)} + (\hat{P}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[(b_i)^{(9)} + (b_i')^{(9)} + (\hat{B}_{44})^{(9)} + (\hat{Q}_{44})^{(9)} (\hat{k}_{44})^{(9)}] < 1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{13})^{(1)} e^{(\widehat{M}_{13})^{(1)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0), T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\widehat{M}_{20})^{(3)} t} \ , \ T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le (\hat{P}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\widehat{M}_{24})^{(4)} t} \ , \ \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0), T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \ , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy 154

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \quad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{\mathrm{T}}_{15}(\mathrm{t}) = \mathrm{T}_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{16})^{(2)}$, $T_i^0 \le (\hat{Q}_{16})^{(2)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t}$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{24})^{(4)} \text{ , } T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\bar{M}_{24})^{(4)} t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\bar{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{32})^{(6)}$, $T_i^0 \le (\hat{Q}_{32})^{(6)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\bar{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)} (T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)} (G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\overline{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^{0} + \int_0^t \left[(b_{42})^{(8)} T_{41}\left(s_{(40)}\right) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}\left(G_{43}\left(s_{(40)}\right), s_{(40)}\right)\right) T_{42}\left(s_{(40)}\right)\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0 \,,\; T_i(0) = T_i^0 \,,\; G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} \,, T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\,\hat{P}_{44}\,)^{(9)} e^{(\,\hat{M}_{44}\,)^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\,\hat{Q}_{44}\,)^{(9)} e^{(\,\hat{M}_{44}\,)^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}\left(s_{(44)}\right) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}\left(T_{45}\left(s_{(44)}\right), s_{(44)}\right)\right) G_{44}\left(s_{(44)}\right)\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}\left(s_{(44)}\right) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}\left(T_{45}\left(s_{(44)}\right), s_{(44)}\right)\right) G_{45}\left(s_{(44)}\right)\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}\left(s_{(44)}\right) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}\left(T_{45}\left(s_{(44)}\right), s_{(44)}\right)\right) G_{46}\left(s_{(44)}\right)\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}\left(s_{(44)}\right) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}\left(G_{47}\left(s_{(44)}\right), s_{(44)}\right)\right) T_{44}\left(s_{(44)}\right)\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}\left(s_{(44)}\right) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}\left(G_{47}\left(s_{(44)}\right), s_{(44)}\right)\right) T_{45}\left(s_{(44)}\right)\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}\left(s_{(44)}\right) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}\left(G_{47}\left(s_{(44)}\right), s_{(44)}\right)\right) T_{46}\left(s_{(44)}\right)\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\,\hat{P}_{13}\,)^{(1)} e^{(\,\hat{M}_{13}\,)^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\,\hat{P}_{13}\,)^{(1)}}{(\,\hat{M}_{13}\,)^{(1)}} \left(e^{(\,\hat{M}_{13}\,)^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^0) e^{-(\,\hat{M}_{13}\,)^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\,\hat{M}_{13}\,)^{(1)}} \left[\left((\,\hat{P}_{13}\,)^{(1)} + G_{14}^0\right) e^{\left(-\frac{(\,\hat{P}_{13}\,)^{(1)} + G_{14}^0}{G_{14}^0}\right)} + (\,\hat{P}_{13}\,)^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for G_{14} , $G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^0 + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^0 + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^0 + \frac{(a_{16})^{(2)} (\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t) - G_{16}^0) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^0\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^0}{G_{17}^0}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t) \le G_{20}^0 + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^0 + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^0 + \frac{(a_{20})^{(3)} (\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that (172)

$$(G_{20}(t) - G_{20}^0) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^0\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^0}{G_{21}^0}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t) \le G_{24}^0 + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^0 + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^0 + \frac{(a_{24})^{(4)} (\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that (174)

$$(G_{24}(t) - G_{24}^0) e^{-(\hat{M}_{24})^{(4)} t} \le \frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{P}_{24})^{(4)} + G_{25}^0\right) e^{\left(-\frac{(\hat{P}_{24})^{(4)} + G_{25}^0}{G_{25}^0}\right)} + (\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \le G_{28}^0 + \int_0^t \left[(a_{28})^{(5)}\left(G_{29}^0 + (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} s_{(28)}}\right)\right] ds_{(28)} =$$

$$\left(1 + (a_{28})^{(5)} t\right) G_{29}^0 + \frac{(a_{28})^{(5)} (\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)} t} - 1\right)$$

From which it follows that (175)

$$(G_{28}(t) - G_{28}^0) e^{-(\hat{M}_{28})^{(5)} t} \le \frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[\left((\hat{P}_{28})^{(5)} + G_{29}^0\right) e^{\left(-\frac{(\hat{P}_{28})^{(5)} + G_{29}^0}{G_{29}^0}\right)} + (\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \le G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} \left(G_{33}^0 + (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} s_{(32)}} \right) \right] ds_{(32)} =$$

$$\left(1 + (a_{32})^{(6)} t\right) G_{33}^0 + \frac{(a_{32})^{(6)} (\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left(e^{(\hat{M}_{32})^{(6)} t} - 1 \right)$$

From which it follows that 177

$$(G_{32}(t) - G_{32}^0) e^{-(\hat{M}_{32})^{(6)} t} \le \frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}} \left[\left((\hat{P}_{32})^{(6)} + G_{33}^0 \right) e^{\left(- \frac{(\hat{P}_{32})^{(6)} + G_{33}^0}{G_{33}^0} \right)} + (\hat{P}_{32})^{(6)} \right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \le G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} \left(G_{37}^0 + (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} s_{(36)}} \right) \right] ds_{(36)} =$$

$$\left(1 + (a_{36})^{(7)} t\right) G_{37}^0 + \frac{(a_{36})^{(7)} (\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left(e^{(\hat{M}_{36})^{(7)} t} - 1 \right)$$

From which it follows that

$$(G_{36}(t) - G_{36}^0) e^{-(\hat{M}_{36})^{(7)} t} \le \frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}} \left[\left((\hat{P}_{36})^{(7)} + G_{37}^0 \right) e^{\left(- \frac{(\hat{P}_{36})^{(7)} + G_{37}^0}{G_{37}^0} \right)} + (\hat{P}_{36})^{(7)} \right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}} \right) \right] ds_{(40)} = \quad 180$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1 \right)$$

From which it follows that 181

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0 \right) e^{\left(- \frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0} \right)} + (\hat{P}_{40})^{(8)} \right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}} \right) \right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1 \right)$$

From which it follows that

$$(G_{44}(t)-G_{44}^{0})e^{-(\hat{M}_{44})^{(9)}t}\leq\frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}}\left[((\hat{P}_{44})^{(9)}+G_{45}^{0})e^{\left(-\frac{(\hat{P}_{44})^{(9)}+G_{45}^{0}}{G_{45}^{0}}\right)}+(\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[(\hat{P}_{13})^{(1)}+((\hat{P}_{13})^{(1)}+G_j^0)e^{-\left(\frac{(\hat{P}_{13})^{(1)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{Q}_{13})^{(1)}+T_j^0)e^{-\left(\frac{(\hat{Q}_{13})^{(1)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{13})^{(1)}\right]\leq(\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)},T^{(1)}\right),\left(G^{(2)},T^{(2)}\right)\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{13})^{(1)}t}\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:(\tilde{G},\tilde{T})=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_0^t\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{-(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\hat{M}_{13})^{(1)}s_{(13)}}e^{(\hat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\hat{M}_{13})^{(1)}t}\leq \quad 186$$
$$\frac{1}{(\hat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a'_{13})^{(1)}+(\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\,G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ *and* $(\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(1)}$ and $(b''_i)^{(1)}, i = 13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 19 to 24 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(1)} - (a''_i)^{(1)}(T_{14}(s_{(13)}), s_{(13)})\} ds_{(13)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(1)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1, \left((\widehat{M}_{13})^{(1)}\right)_2$ *and* $\left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also G_{14} *and* G_{15} . indeed if

$G_{13} < (\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt} \leq \left((\widehat{M}_{13})^{(1)}\right)_1 - (a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14} \leq \left((\widehat{M}_{13})^{(1)}\right)_2 = G_{14}^0 + 2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15} \leq \left((\widehat{M}_{13})^{(1)}\right)_3 = G_{15}^0 + 2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If G_{14} *or* G_{15} is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15} . The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(1)}(G(t), t)) = (b'_{14})^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$$(b_{14})^{(1)} - (b''_i)^{(1)}(G(t), t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b''_{15})^{(1)}(G(t), t) = (b'_{15})^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}$, $\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $\left(\widetilde{G_{19}}, \widetilde{T_{19}}\right) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a'_{16})^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a''_{16})^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a''_{16})^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\hat{M}_{16})^{(2)}t} \leq$$
$$\frac{1}{(\hat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}\right)d\left(((G_{19})^{(1)}, (T_{19})^{(1)};\ (G_{19})^{(2)}, (T_{19})^{(2)})\right)$$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\hat{P}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ and $(\hat{Q}_{16})^{(2)}e^{(\hat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and

respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$$

In the same way , one can obtain

$$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$ 205

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\widehat{P}_{20})^{(3)}$ and $(\widehat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[(\widehat{P}_{20})^{(3)} + \left((\widehat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\widehat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad (209)$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself (210)

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric (211)

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{20})^{(3)}t}\}$$

Indeed if we denote (212)

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results (213)

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\widehat{M}_{20})^{(3)}s_{(20)}}e^{(\widehat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\widehat{M}_{20})^{(3)}t}\leq \quad (214)$$
$$\frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\widehat{A}_{20})^{(3)}+(\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. (215)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i=20,21,22$ depend only on T_{21} and respectively on (G_{23})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (216)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(3)}-(a_i'')^{(3)}(T_{21}(s_{(20)}),s_{(20)})\}ds_{(20)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2$ *and* $\left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also G_{21} *and* G_{22} . indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)} G_{21}$ and by integrating

$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$

In the same way , one can obtain

$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$

If G_{21} *or* G_{22} is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for G_{21} *and* G_{22} . The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t),t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t),t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t),t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[(\hat{P}_{24})^{(4)} + \left((\hat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}\left[\left((\hat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{24})^{(4)}\right] \leq (\hat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{24})^{(4)}\left|G_{25}^{(1)}-G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)\left|G_{24}^{(1)}-G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}\left|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)\right|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)}-(G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t}\le \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};\,(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}\ and\ (\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on $(G_{27})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 228

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(4)}-(a_i'')^{(4)}(T_{25}(s_{(24)}),s_{(24)})\}ds_{(24)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{(-(b_i')^{(4)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1,\left((\widehat{M}_{24})^{(4)}\right)_2\ and\ \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25}\ and\ G_{26}$. indeed if

$G_{24}<(\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt}\le\left((\widehat{M}_{24})^{(4)}\right)_1-(a'_{25})^{(4)}G_{25}$ and by integrating

$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$

In the same way , one can obtain

$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t),t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t),t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t),t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}} < 1$ and to choose 233

$(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[(\hat{P}_{28})^{(5)} + \left((\hat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\bar{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{28})^{(5)}\right] \leq (\hat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\le \qquad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}\ and\ (\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. 238

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)},i=28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 239

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(5)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1,\left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28}<(\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt}\le\left((\widehat{M}_{28})^{(5)}\right)_1-(a'_{29})^{(5)}G_{29}$ and by integrating

$$G_{29}\le\left((\widehat{M}_{28})^{(5)}\right)_2=G_{29}^0+2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a'_{29})^{(5)}$$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a'_{30})^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t),t)) = (b'_{29})^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b'_{30})^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[(\hat{P}_{32})^{(6)} + \left((\hat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{32})^{(6)} \quad 245$$

$$\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{32})^{(6)}\right] \leq (\hat{Q}_{32})^{(6)} \quad 246$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric 247

$$d\left(\left((G_{35})^{(1)},(T_{35})^{(1)}\right),\left((G_{35})^{(2)},(T_{35})^{(2)}\right)\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\hat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{35})}, \widetilde{(T_{35})}$: $\left(\widetilde{(G_{35})}, \widetilde{(T_{35})}\right) = \mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_{0}^{t}\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \qquad 248$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};\ (G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_{+}$. 249

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on $(G_{35})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 250

it results

$$G_i\,(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$$T_i\,(t)\geq T_i^0e^{(-(b_i')^{(6)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1,\left((\widehat{M}_{32})^{(6)}\right)_2\ and\ \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33}\ and\ G_{34}$. indeed if

$G_{32}<(\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt}\leq\left((\widehat{M}_{32})^{(6)}\right)_1-(a'_{33})^{(6)}G_{33}$ and by integrating

$$G_{33}\leq\left((\widehat{M}_{32})^{(6)}\right)_2=G_{33}^0+2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a'_{33})^{(6)}$$

In the same way , one can obtain

$$G_{34}\leq\left((\widehat{M}_{32})^{(6)}\right)_3=G_{34}^0+2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a'_{34})^{(6)}$$

If $G_{33}\ or\ G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33}\ and\ G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\big((G_{35})(t),t\big) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{37}, G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}} < 1$ and to choose
$(\widehat{P}_{36})^{(7)}$ and $(\widehat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[(\widehat{P}_{36})^{(7)} + \left((\widehat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\widehat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{36})^{(7)}\right] \leq (\widehat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})}, \widetilde{(T_{39})} : \left(\widetilde{(G_{39})}, \widetilde{(T_{39})}\right) = \mathcal{A}^{(7)}((G_{39}),(T_{39}))$
It results

$$|\tilde{G}_{36}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{36})^{(7)}|G_{37}^{(1)} - G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)} +$$
$$\int_0^t\{(a_{36}')^{(7)}|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}} +$$
$$(a_{36}'')^{(7)}(T_{37}^{(1)}, s_{(36)})|G_{36}^{(1)} - G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}} +$$

$G_{36}^{(2)}|(a_{36}'')^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a_{36}'')^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\leq \frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a_{36}')^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right) \quad (259)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a_{36}'')^{(7)}$ and $(b_{36}'')^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. (260)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0\ and\ T_i(t)=0$ (261)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$$T_i(t)\geq T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2\ and\ \left((\widehat{M}_{36})^{(7)}\right)_3$: (262)

Remark 33: if G_{36} is bounded, the same property have also $G_{37}\ and\ G_{38}$. indeed if

$G_{36}<(\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt}\leq\left((\widehat{M}_{36})^{(7)}\right)_1-(a_{37}')^{(7)}G_{37}$ and by integrating

$$G_{37}\leq\left((\widehat{M}_{36})^{(7)}\right)_2=G_{37}^0+2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$$

In the same way , one can obtain

$$G_{38}\leq\left((\widehat{M}_{36})^{(7)}\right)_3=G_{38}^0+2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$$

If $G_{37}\ or\ G_{38}$ is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If $G_{36}\ is$ bounded, from below, the same property holds for $G_{37}\ and\ G_{38}$. The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. (263)

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t))=(b_{37}')^{(7)}$ then $T_{37}\to\infty$. (264)

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t), t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t), t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)}, (T_{43})^{(1)}), ((G_{43})^{(2)}, (T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})}, \widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})}, \widetilde{(T_{43})}\right) = \mathcal{A}^{(8)}((G_{43}), (T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{40})^{(8)}|G_{41}^{(1)} - G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)} +$$
$$\int_0^t\{(a_{40}')^{(8)}|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$(a_{40}'')^{(8)}(T_{41}^{(1)}, s_{(40)})|G_{40}^{(1)} - G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}} +$$
$$G_{40}^{(2)}|(a_{40}'')^{(8)}(T_{41}^{(1)}, s_{(40)}) - (a_{40}'')^{(8)}(T_{41}^{(2)}, s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0, t]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)} - (G_{43})^{(2)}\right| e^{-(\widehat{M}_{40})^{(8)}t} \leq \frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)} + (a'_{40})^{(8)} + (\widehat{A}_{40})^{(8)} + (\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right) d\left(((G_{43})^{(1)}, (T_{43})^{(1)};\ (G_{43})^{(2)}, (T_{43})^{(2)})\right) \quad 273$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)} e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(8)}$ and $(b''_i)^{(8)}, i = 40{,}41{,}42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 275

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(8)} - (a''_i)^{(8)}(T_{41}(s_{(40)}), s_{(40)})\} ds_{(40)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b'_i)^{(8)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1, \left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)} G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1 / (a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2 / (a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(8)}((G_{43})(t), t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\big((G_{43})(t), t\big) < \varepsilon_8, T_{40}\,(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b_{42}'')^{(8)}\left((G_{43})(t), t(t), t\right) = (b_{42}')^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\hat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\hat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{44})^{(9)}\right] \leq (\hat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i, T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)}, (T_{47})^{(1)}), ((G_{47})^{(2)}, (T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}), (\widetilde{T_{47}}) : \left((\widetilde{G_{47}}), (\widetilde{T_{47}})\right) = \mathcal{A}^{(9)}\big((G_{47}), (T_{47})\big)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{44})^{(9)}\left|G_{45}^{(1)} - G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)} +$$
$$\int_0^t\{(a_{44}')^{(9)}\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}} +$$
$$G_{44}^{(2)}\left|(a_{44}'')^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a_{44}'')^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t} \leq$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ *and* $(\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t) = 0$ *and* $T_i\,(t) = 0$

From 99 to 44 it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b'_i)^{(9)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)}G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right)) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}\,(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t), t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G, t) - (b''_{14})^{(1)}(G, t) \leq -(\tau_1)^{(1)}$

Definition of $(\nu_1)^{(1)}, (\nu_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, \nu^{(1)}, u^{(1)}$: 281

By $(\nu_1)^{(1)} > 0$, $(\nu_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{\nu}_1)^{(1)}, , (\bar{\nu}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{\nu}_1)^{(1)} > 0$, $(\bar{\nu}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(\nu^{(1)}\right)^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by
$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)}$, $if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)}$,

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)}$, $if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)}$, $if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)}$,

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^{0}e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}}G_{13}^{0}e^{((S_1)^{(1)}-(p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}}G_{13}^{0}e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_1)^{(1)}((S_1)^{(1)}-(p_{13})^{(1)}-(S_2)^{(1)})}\left[e^{((S_1)^{(1)}-(p_{13})^{(1)})t}-e^{-(S_2)^{(1)}t}\right]+G_{15}^{0}e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left.\frac{(a_{15})^{(1)}G_{13}^{0}}{(m_2)^{(1)}((S_1)^{(1)}-(a'_{15})^{(1)})}[e^{(S_1)^{(1)}t}-e^{-(a'_{15})^{(1)}t}]+\ G_{15}^{0}e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}}T_{13}^{0}e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}}T_{13}^{0}e^{((R_1)^{(1)}+(r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)}T_{13}^{0}}{(\mu_1)^{(1)}((R_1)^{(1)}-(b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t}-e^{-(b'_{15})^{(1)}t}\right]+T_{15}^{0}e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)}T_{13}^{0}}{(\mu_2)^{(1)}((R_1)^{(1)}+(r_{13})^{(1)}+(R_2)^{(1)})}\left[e^{((R_1)^{(1)}+(r_{13})^{(1)})t}-e^{-(R_2)^{(1)}t}\right]+T_{15}^{0}e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(T_{17}, t) + (a''_{17})^{(2)}(T_{17}, t) \le -(\sigma_1)^{(2)} \quad 293$$

$$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)} \quad 294$$

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{v}_1)^{(2)},,(\bar{v}_2)^{(2)},(\bar{u}_1)^{(2)},(\bar{u}_2)^{(2)}$: 299

By $(\bar{v}_1)^{(2)} > 0$, $(\bar{v}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\left(v^{(2)}\right)^2 + (\sigma_2)^{(2)}v^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\left(u^{(2)}\right)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}$, $(m_2)^{(2)}$, $(\mu_1)^{(2)},(\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (v_0)^{(2)}, (m_1)^{(2)} = (v_1)^{(2)},\ \boldsymbol{if}\ (v_0)^{(2)} < (v_1)^{(2)}$ 305

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (\bar{v}_1)^{(2)}, \boldsymbol{if}\ (v_1)^{(2)} < (v_0)^{(2)} < (\bar{v}_1)^{(2)},$ 306

and $\boxed{(v_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}}$

$(m_2)^{(2)} = (v_1)^{(2)}, (m_1)^{(2)} = (v_0)^{(2)},\ \boldsymbol{if}\ (\bar{v}_1)^{(2)} < (v_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)},\ \boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)},$

and $\boxed{(u_0)^{(2)} = \frac{T_{16}^0}{T_{17}^0}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)}t}$

$(p_i)^{(2)}$ is defined by equation

$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)}t}$ 311

$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)}t}\right] + G_{18}^0 e^{-(S_2)^{(2)}t} \le G_{18}(t) \le\right.$ 312

$\left.\frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)}t} - e^{-(a'_{18})^{(2)}t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)}t}\right)$

$\boxed{T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}}$ 313

$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}$ 314

$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)}t} - e^{-(b'_{18})^{(2)}t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le$ 315

$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)}t}\right] + T_{18}^0 e^{-(R_2)^{(2)}t}$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)}$$ 318

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, ***if*** $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, ***if*** $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, ***if*** $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T^0_{20}}{T^0_{21}}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^{0}e^{(S_1)^{(3)}t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}}G_{20}^{0}e^{((S_1)^{(3)}-(p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}}G_{20}^{0}e^{(S_1)^{(3)}t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_1)^{(3)}((S_1)^{(3)}-(p_{20})^{(3)}-(S_2)^{(3)})}\left[e^{((S_1)^{(3)}-(p_{20})^{(3)})t}-e^{-(S_2)^{(3)}t}\right]+G_{22}^{0}e^{-(S_2)^{(3)}t} \le G_{22}(t) \le \right. \quad 324$$

$$\left.\frac{(a_{22})^{(3)}G_{20}^{0}}{(m_2)^{(3)}((S_1)^{(3)}-(a'_{22})^{(3)})}[e^{(S_1)^{(3)}t}-e^{-(a'_{22})^{(3)}t}]+G_{22}^{0}e^{-(a'_{22})^{(3)}t}\right)$$

$$\boxed{T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}}T_{20}^{0}e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}}T_{20}^{0}e^{((R_1)^{(3)}+(r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)}T_{20}^{0}}{(\mu_1)^{(3)}((R_1)^{(3)}-(b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t}-e^{-(b'_{22})^{(3)}t}\right]+T_{22}^{0}e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le \quad 327$$

$$\frac{(a_{22})^{(3)}T_{20}^{0}}{(\mu_2)^{(3)}((R_1)^{(3)}+(r_{20})^{(3)}+(R_2)^{(3)})}\left[e^{((R_1)^{(3)}+(r_{20})^{(3)})t}-e^{-(R_2)^{(3)}t}\right]+T_{22}^{0}e^{-(R_2)^{(3)}t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(v_1)^{(4)}, (v_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, v^{(4)}, u^{(4)}$: 329

By $(v_1)^{(4)} > 0$, $(v_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(v^{(4)}\big)^2 + (\sigma_1)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$

and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{v}_1)^{(4)}, , (\bar{v}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\left(v^{(4)}\right)^2 + (\sigma_2)^{(4)}v^{(4)} - (a_{24})^{(4)} = 0$ and $(b_{25})^{(4)}\left(u^{(4)}\right)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$, $(\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}$, $(m_2)^{(4)}$, $(\mu_1)^{(4)}$, $(\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}\ , \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)} \tag{331}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}\ , \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T_{24}^0}{T_{25}^0}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)}t} \tag{332}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \tag{333}$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})}\left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \tag{334}$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \tag{335}$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \tag{336}$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})}\left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$

$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$

Definition of $(\nu_1)^{(5)}, (\nu_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, \nu^{(5)}, u^{(5)}$: 339

By $(\nu_1)^{(5)} > 0$, $(\nu_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},, (\bar{\nu}_2)^{(5)}, (\bar{u}_1)^{(5)}, (\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(\nu^{(5)}\big)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$
and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}, (\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}, (m_2)^{(5)}, (\mu_1)^{(5)}, (\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)}$, ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)}$,
and $\boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)}$, ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}$ $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}$ $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{28}(t) \leq G_{28}^0 e^{(S_1)^{(5)} t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \leq G_{29}(t) \leq \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)} t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)} t}\right] + G_{30}^0 e^{-(S_2)^{(5)} t} \leq G_{30}(t) \leq \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)} t} - e^{-(a'_{30})^{(5)} t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)} t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)} t} \leq T_{28}(t) \leq \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)} - (b'_{30})^{(5)})}\left[e^{(R_1)^{(5)} t} - e^{-(b'_{30})^{(5)} t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)} t} \leq T_{30}(t) \leq \quad 347$$

$$\frac{(a_{30})^{(5)} T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)} + (r_{28})^{(5)} + (R_2)^{(5)})}\left[e^{((R_1)^{(5)} + (r_{28})^{(5)})t} - e^{-(R_2)^{(5)} t}\right] + T_{30}^0 e^{-(R_2)^{(5)} t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \leq -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \leq -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}),t\big) - (b''_{33})^{(6)}\big((G_{35}),t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}$, $(m_2)^{(6)}$, $(\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}\ (v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}\ (v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}$, $\boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}$, $\boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)}$,
and $\boxed{(u_0)^{(6)} = \frac{T^0_{32}}{T^0_{33}}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G^0_{32}e^{(S_1)^{(6)}t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}}G^0_{32}e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}}G^0_{32}e^{(S_1)^{(6)}t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)}G^0_{32}}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)}t}\right] + G^0_{34}e^{-(S_2)^{(6)}t} \le G_{34}(t) \le \right. \quad 355$$
$$\left. \frac{(a_{34})^{(6)}G^0_{32}}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)}t} - e^{-(a'_{34})^{(6)}t}\right] + G^0_{34}e^{-(a'_{34})^{(6)}t}\right)$$

$$\boxed{T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}}T_{32}^{0}e^{(R_1)^{(6)}t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}}T_{32}^{0}e^{((R_1)^{(6)}+(r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)}T_{32}^{0}}{(\mu_1)^{(6)}((R_1)^{(6)}-(b'_{34})^{(6)})}\left[e^{(R_1)^{(6)}t} - e^{-(b'_{34})^{(6)}t}\right] + T_{34}^{0}e^{-(b'_{34})^{(6)}t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)}T_{32}^{0}}{(\mu_2)^{(6)}((R_1)^{(6)}+(r_{32})^{(6)}+(R_2)^{(6)})}\left[e^{((R_1)^{(6)}+(r_{32})^{(6)})t} - e^{-(R_2)^{(6)}t}\right] + T_{34}^{0}e^{-(R_2)^{(6)}t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0$, $(\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)}, , (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0$, $(\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\left(\nu^{(7)}\right)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\left(u^{(7)}\right)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}$, ***if*** $(\nu_0)^{(7)} < (\nu_1)^{(7)}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}$, $\boldsymbol{if}$ $(\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)}$,

and $\boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$

$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}$, $\boldsymbol{if}$ $(\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$

and analogously 363

$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}$, $\boldsymbol{if}$ $(u_0)^{(7)} < (u_1)^{(7)}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}$, $\boldsymbol{if}$ $(u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)}$,

and $\boxed{(u_0)^{(7)} = \frac{T_{36}^0}{T_{37}^0}}$

$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}$ $(\bar{u}_1)^{(7)} < (u_0)^{(7)}$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{36}(t) \leq G_{36}^0 e^{(S_1)^{(7)} t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \leq G_{37}(t) \leq \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)} t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)} - (p_{36})^{(7)} - (S_2)^{(7)})}\left[e^{((S_1)^{(7)} - (p_{36})^{(7)})t} - e^{-(S_2)^{(7)} t}\right] + G_{38}^0 e^{-(S_2)^{(7)} t} \leq G_{38}(t) \leq \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)} - (a'_{38})^{(7)})}\left[e^{(S_1)^{(7)} t} - e^{-(a'_{38})^{(7)} t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)} t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)} t} \leq T_{36}(t) \leq \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)} + (r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)} - (b'_{38})^{(7)})}\left[e^{(R_1)^{(7)} t} - e^{-(b'_{38})^{(7)} t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)} t} \leq T_{38}(t) \leq \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)} + (r_{36})^{(7)} + (R_2)^{(7)})}\left[e^{((R_1)^{(7)} + (r_{36})^{(7)})t} - e^{-(R_2)^{(7)} t}\right] + T_{38}^0 e^{-(R_2)^{(7)} t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0$, $(\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0$, $(u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0$, $(\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0$, $(\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$$(m_2)^{(8)} = (\nu_0)^{(8)}, (m_1)^{(8)} = (\nu_1)^{(8)}, \ \boldsymbol{if}\ (\nu_0)^{(8)} < (\nu_1)^{(8)}$$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)} < (\nu_0)^{(8)} < (\bar{\nu}_1)^{(8)},$$

and $\boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}}$

$$(m_2)^{(8)} = (\nu_1)^{(8)}, (m_1)^{(8)} = (\nu_0)^{(8)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)} < (\nu_0)^{(8)}$$

and analogously 374

$$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}, \ \boldsymbol{if}\ (u_0)^{(8)} < (u_1)^{(8)}$$

$$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)},$$

and $\boxed{(u_0)^{(8)} = \frac{T^0_{40}}{T^0_{41}}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^{0}e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G_{40}^{0}e^{((S_1)^{(8)}-(p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}}G_{40}^{0}e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_1)^{(8)}((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)})}\left[e^{((S_1)^{(8)}-(p_{40})^{(8)})t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^{0}e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^{0}}{(m_2)^{(8)}((S_1)^{(8)}-(a'_{42})^{(8)})}[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}]+\ G_{42}^{0}e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^{0}e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}}T_{40}^{0}e^{((R_1)^{(8)}+(r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^{0}}{(\mu_1)^{(8)}((R_1)^{(8)}-(b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^{0}e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^{0}}{(\mu_2)^{(8)}((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)})}\left[e^{((R_1)^{(8)}+(r_{40})^{(8)})t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^{0}e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$

and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_1)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$ and

Definition of $(\bar{v}_1)^{(9)},,(\bar{v}_2)^{(9)},(\bar{u}_1)^{(9)},(\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)}>0$, $(\bar{v}_2)^{(9)}<0$ and respectively $(\bar{u}_1)^{(9)}>0$, $(\bar{u}_2)^{(9)}<0$ the roots of the equations $(a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}=0$ and $(b_{45})^{(9)}\left(u^{(9)}\right)^2+(\tau_2)^{(9)}u^{(9)}-(b_{44})^{(9)}=0$

Definition of $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)},(v_0)^{(9)}$:-

If we define $(m_1)^{(9)}$, $(m_2)^{(9)}$, $(\mu_1)^{(9)},(\mu_2)^{(9)}$ by

$(m_2)^{(9)}=(v_0)^{(9)},(m_1)^{(9)}=(v_1)^{(9)},$ $\boldsymbol{if}$ $(v_0)^{(9)}<(v_1)^{(9)}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(\bar{v}_1)^{(9)}$, $\boldsymbol{if}$ $(v_1)^{(9)}<(v_0)^{(9)}<(\bar{v}_1)^{(9)}$,

and $\boxed{(v_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$

$(m_2)^{(9)}=(v_1)^{(9)},(m_1)^{(9)}=(v_0)^{(9)},$ $\boldsymbol{if}$ $(\bar{v}_1)^{(9)}<(v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)}=(u_0)^{(9)},(\mu_1)^{(9)}=(u_1)^{(9)},$ $\boldsymbol{if}$ $(u_0)^{(9)}<(u_1)^{(9)}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)}<(u_0)^{(9)}<(\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)}=\frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)}=(u_1)^{(9)},(\mu_1)^{(9)}=(u_0)^{(9)},\boldsymbol{if}$ $(\bar{u}_1)^{(9)}<(u_0)^{(9)}$ where $(u_1)^{(9)},(\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{44}(t)\le G_{44}^0e^{(S_1)^{(9)}t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}}G_{44}^0e^{((S_1)^{(9)}-(p_{44})^{(9)})t}\le G_{45}(t)\le\frac{1}{(m_2)^{(9)}}G_{44}^0e^{(S_1)^{(9)}t}$$

$$\left(\frac{(a_{46})^{(9)}G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)}-(p_{44})^{(9)}-(S_2)^{(9)})}\left[e^{((S_1)^{(9)}-(p_{44})^{(9)})t}-e^{-(S_2)^{(9)}t}\right]+G_{46}^0e^{-(S_2)^{(9)}t}\le G_{46}(t)\le\right.$$
$$\left.\frac{(a_{46})^{(9)}G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)}-(a'_{46})^{(9)})}[e^{(S_1)^{(9)}t}-e^{-(a'_{46})^{(9)}t}]+\ G_{46}^0e^{-(a'_{46})^{(9)}t}\right)$$

$$\boxed{T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}}T_{44}^0e^{(R_1)^{(9)}t}\le T_{44}(t)\le\frac{1}{(\mu_2)^{(9)}}T_{44}^0e^{((R_1)^{(9)}+(r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)}T_{44}^{0}}{(\mu_1)^{(9)}((R_1)^{(9)}-(b'_{46})^{(9)})}\left[e^{(R_1)^{(9)}t}-e^{-(b'_{46})^{(9)}t}\right]+T_{46}^{0}e^{-(b'_{46})^{(9)}t}\leq T_{46}(t)\leq$$

$$\frac{(a_{46})^{(9)}T_{44}^{0}}{(\mu_2)^{(9)}((R_1)^{(9)}+(r_{44})^{(9)}+(R_2)^{(9)})}\left[e^{((R_1)^{(9)}+(r_{44})^{(9)})t}-e^{-(R_2)^{(9)}t}\right]+T_{46}^{0}e^{-(R_2)^{(9)}t}$$

Definition of $(S_1)^{(9)},(S_2)^{(9)},(R_1)^{(9)},(R_2)^{(9)}$:-

Where $(S_1)^{(9)}=(a_{44})^{(9)}(m_2)^{(9)}-(a'_{44})^{(9)}$

$(S_2)^{(9)}=(a_{46})^{(9)}-(p_{46})^{(9)}$

$(R_1)^{(9)}=(b_{44})^{(9)}(\mu_2)^{(9)}-(b'_{44})^{(9)}$

$(R_2)^{(9)}=(b'_{46})^{(9)}-(r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt}=(a_{13})^{(1)}-\left((a'_{13})^{(1)}-(a'_{14})^{(1)}+(a''_{13})^{(1)}(T_{14},t)\right)-(a''_{14})^{(1)}(T_{14},t)\nu^{(1)}-(a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)}=\frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^2+(\sigma_2)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)\leq\frac{d\nu^{(1)}}{dt}\leq-\left((a_{14})^{(1)}(\nu^{(1)})^2+(\sigma_1)^{(1)}\nu^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_0)^{(1)})t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)})t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)})t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq\nu^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^{0}}{G_{14}^{0}}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_2)^{(1)})t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}((\nu_1)^{(1)}-(\nu_2)^{(1)})t\right]}}\leq\nu^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{13}'')^{(1)} = (a_{14}'')^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)}G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b_{13}'')^{(1)} = (b_{14}'')^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a_{16}')^{(2)} - (a_{17}')^{(2)} + (a_{16}'')^{(2)}(T_{17}, t)\right) - (a_{17}'')^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \leq \frac{d\nu^{(2)}}{dt} \leq -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \geq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)}=\frac{(\nu_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)}=\frac{(\bar{\nu}_1)^{(2)}-(\nu_0)^{(2)}}{(\nu_0)^{(2)}-(\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \leq \nu^{(2)}(t) \leq (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \leq \frac{(\nu_1)^{(2)}+(C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}}{1+(C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)}-(\nu_2)^{(2)}\right)t\right]}} \leq \nu^{(2)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \leq (\bar{\nu}_1)^{(2)} \leq (\nu_0)^{(2)} = \frac{G_{16}^0}{G_{17}^0}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t)=\frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t)=\frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right) - (a''_{21})^{(3)}(T_{21},t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \le \frac{dv^{(3)}}{dt} \le -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (v_1)^{(3)} < (\bar{v}_1)^{(3)}$

$$v^{(3)}(t) \ge \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(v_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (v_2)^{(3)}}}$$

it follows $(v_0)^{(3)} \le v^{(3)}(t) \le (v_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{v}_1)^{(3)} - (v_0)^{(3)}}{(v_0)^{(3)} - (\bar{v}_2)^{(3)}}}$$

Definition of $(\bar{v}_1)^{(3)}$:-

From which we deduce $(v_0)^{(3)} \le v^{(3)}(t) \le (\bar{v}_1)^{(3)}$

If $0 < (v_1)^{(3)} < (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{v}_1)^{(3)}$ we find like in the previous case, 402

$$(v_1)^{(3)} \le \frac{(v_1)^{(3)} + (C)^{(3)}(v_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}}{1 + (C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((v_1)^{(3)} - (v_2)^{(3)}\right)t\right]}} \le v^{(3)}(t) \le$$

$$\frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (\bar{v}_1)^{(3)}$$

If $0 < (v_1)^{(3)} \le (\bar{v}_1)^{(3)} \le (v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(v_1)^{(3)} \le v^{(3)}(t) \le \frac{(\bar{v}_1)^{(3)} + (\bar{C})^{(3)}(\bar{v}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{v}_1)^{(3)} - (\bar{v}_2)^{(3)}\right)t\right]}} \le (v_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(3)}(t)$:-

$$(m_2)^{(3)} \le v^{(3)}(t) \le (m_1)^{(3)}, \quad \boxed{v^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)} G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$ if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)} T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \le \frac{(\nu_1)^{(4)}+(C)^{(4)}(\nu_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)}-(\nu_2)^{(4)}\right)t\right]}} \le \nu^{(4)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \le (\bar{\nu}_1)^{(4)} \le \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain

407

$$(\nu_1)^{(4)} \le \nu^{(4)}(t) \le \frac{(\bar{\nu}_1)^{(4)}+(\bar{C})^{(4)}(\bar{\nu}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)}-(\bar{\nu}_2)^{(4)}\right)t\right]}} \le (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$(m_2)^{(4)} \le \nu^{(4)}(t) \le (m_1)^{(4)}$, $\boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$(\mu_2)^{(4)} \le u^{(4)}(t) \le (\mu_1)^{(4)}$, $\boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)}G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)}T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \le \frac{d\nu^{(5)}}{dt} \le -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also**

defines $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b''_{28})^{(5)} = (b''_{29})^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a'_{32})^{(6)} - (a'_{33})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right) - (a''_{33})^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)} - (\nu_0)^{(6)}}{(\nu_0)^{(6)} - (\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)} + (C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}}{1 + (C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)} - (\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G^0_{32}}{G^0_{33}}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)} + (\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}}{1 + (\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)} - (\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(6)}(t)$:-

$$(m_2)^{(6)} \leq v^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{v^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $v^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)} G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)} T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\quad \boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$$

$$v^{(7)}(t) \geq \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)} e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \leq v^{(7)}(t) \leq (\nu_1)^{(7)}$

In the same manner , we get

417

$$\nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)}-(\nu_0)^{(7)}}{(\nu_0)^{(7)}-(\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le \nu^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)}+(C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}}{1+(C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)}-(\nu_2)^{(7)}\right)t\right]}} \le \nu^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G_{36}^0}{G_{37}^0}}$, we obtain 419

$$(\nu_1)^{(7)} \le \nu^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)}+(\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}}{1+(\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)}-(\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have
Definition of $\nu^{(7)}(t)$:-

$$(m_2)^{(7)} \le \nu^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \le u^{(7)}(t) \le (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)}G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)}T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right) - (a''_{41})^{(8)}(T_{41},t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $v^{(8)}$:- $\boxed{v^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)}\right) \leq \frac{dv^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_1)^{(8)}v^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(8)}, (v_0)^{(8)}$:-

For $0 < \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}} < (v_1)^{(8)} < (\bar{v}_1)^{(8)}$

$$v^{(8)}(t) \geq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(v_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (v_2)^{(8)}}}$$

it follows $(v_0)^{(8)} \leq v^{(8)}(t) \leq (v_1)^{(8)}$

In the same manner , we get 422

$$v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{v}_1)^{(8)} - (v_0)^{(8)}}{(v_0)^{(8)} - (\bar{v}_2)^{(8)}}}$$

From which we deduce $(v_0)^{(8)} \leq v^{(8)}(t) \leq (\bar{v}_8)^{(8)}$

If $0 < (v_1)^{(8)} < (v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{v}_1)^{(8)}$ we find like in the previous case, 423

$$(v_1)^{(8)} \leq \frac{(v_1)^{(8)} + (C)^{(8)}(v_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}}{1 + (C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((v_1)^{(8)} - (v_2)^{(8)}\right)t\right]}} \leq v^{(8)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (\bar{v}_1)^{(8)}$$

If $0 < (v_1)^{(8)} \leq (\bar{v}_1)^{(8)} \leq \boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(v_1)^{(8)} \leq v^{(8)}(t) \leq \frac{(\bar{v}_1)^{(8)} + (\bar{C})^{(8)}(\bar{v}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{v}_1)^{(8)} - (\bar{v}_2)^{(8)}\right)t\right]}} \leq (v_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(8)}(t)$:-

$$(m_2)^{(8)} \leq v^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{v^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)} G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)} T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424

Proof : From 99,20,44,22,23,44 we obtain A

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \le \frac{d\nu^{(9)}}{dt} \le -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \ge \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \le \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)} - (\nu_0)^{(9)}}{(\nu_0)^{(9)} - (\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \le \nu^{(9)}(t) \le (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \le \frac{(\nu_1)^{(9)} + (C)^{(9)}(\nu_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}}{1 + (C)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)} - (\nu_2)^{(9)}\right)t\right]}} \le \nu^{(9)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \le (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)} + (\bar{C})^{(9)}(\bar{\nu}_2)^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}}{1 + (\bar{C})^{(9)} e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)} - (\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$$

$$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0\,,$$

$$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$$

$with\ \ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0 \qquad 427$$

$$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0 \qquad 428$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0\,, \qquad 429$$

$$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0 \qquad 430$$

$with\ \ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(3)} and\ (b_i'')^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ \ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a_i'')^{(4)} and\ (b_i'')^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0$ 437

$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0$ 438

$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0$ 439

$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0$ 440

$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0$ 441

$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0$ 442

has a unique positive solution , which is an equilibrium solution for the system

$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0$ 443

$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0$ 444

$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0$ 445

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27})\big)\,]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27})\big)\,]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\big((G_{27})\big)\,]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})\,]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})\,]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})\,]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})\,]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})\,]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})\,]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$

$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$

Proof: 486

(n) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]}$$ 495

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T_{33}^* :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T_{33}^* for which $f(T_{33}^*) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)} G_{33}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)} G_{33}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

Definition and uniqueness of T_{37}^* :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T_{37}^* for which $f(T_{37}^*) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)} G_{37}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)} G_{37}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504

that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{14}^{*} such that $\varphi((G_{19})^{*}) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{21}^{*} such that $\varphi((G_{23})^{*}) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{25}^{*} such that $\varphi((G_{27})^{*}) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507

$$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{29}^{*} such that $\varphi((G_{31})^{*}) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508

$$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{33}^{*} such that $\varphi({G_{35}}^{*}) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509

$$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$$

$$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G_{37}^{*} such that $\varphi({G_{39}}^{*}) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510

$$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$$

$$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{41}^* such that $\varphi(G_{43}{}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{45}^* such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G_{14}^* given by $\varphi(G^*) = 0$, T_{14}^* given by $f(T_{14}^*) = 0$ and

$$G_{13}^* = \frac{(a_{13})^{(1)}G_{14}^*}{[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}^*)]} \quad , \quad G_{15}^* = \frac{(a_{15})^{(1)}G_{14}^*}{[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}^*)]}$$

$$T_{13}^* = \frac{(b_{13})^{(1)}T_{14}^*}{[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G^*)]} \quad , \quad T_{15}^* = \frac{(b_{15})^{(1)}T_{14}^*}{[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)}G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)}G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \quad 513$$

$$T_{16}^* = \frac{(b_{16})^{(2)}T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)}T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \quad 514$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)}G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)}G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)}T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)}T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)}G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)}G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)}T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)}T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \quad 517$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$$G_{28}^{*} = \frac{(a_{28})^{(5)} G_{29}^{*}}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^{*})]} \quad , \quad G_{30}^{*} = \frac{(a_{30})^{(5)} G_{29}^{*}}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^{*})]}$$

$$T_{28}^{*} = \frac{(b_{28})^{(5)} T_{29}^{*}}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^{*})]} \quad , \quad T_{30}^{*} = \frac{(b_{30})^{(5)} T_{29}^{*}}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^{*})]} \qquad 519$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*} = \frac{(a_{32})^{(6)} G_{33}^{*}}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^{*})]} \quad , \quad G_{34}^{*} = \frac{(a_{34})^{(6)} G_{33}^{*}}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*} = \frac{(b_{32})^{(6)} T_{33}^{*}}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^{*})]} \quad , \quad T_{34}^{*} = \frac{(b_{34})^{(6)} T_{33}^{*}}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*}) = 0$, T_{37}^{*} given by $f(T_{37}^{*}) = 0$ and

$$G_{36}^{*} = \frac{(a_{36})^{(7)} G_{37}^{*}}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^{*})]} \quad , \quad G_{38}^{*} = \frac{(a_{38})^{(7)} G_{37}^{*}}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*} = \frac{(b_{36})^{(7)} T_{37}^{*}}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^{*})]} \quad , \quad T_{38}^{*} = \frac{(b_{38})^{(7)} T_{37}^{*}}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*}) = 0$, T_{41}^{*} given by $f(T_{41}^{*}) = 0$ and

$$G_{40}^{*} = \frac{(a_{40})^{(8)} G_{41}^{*}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^{*})]} \quad , \quad G_{42}^{*} = \frac{(a_{42})^{(8)} G_{41}^{*}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*} = \frac{(b_{40})^{(8)} T_{41}^{*}}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^{*})]} \quad , \quad T_{42}^{*} = \frac{(b_{42})^{(8)} T_{41}^{*}}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*}) = 0$, T_{45}^{*} given by $f(T_{45}^{*}) = 0$ and

$$G_{44}^{*} = \frac{(a_{44})^{(9)} G_{45}^{*}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^{*})]} \quad , \quad G_{46}^{*} = \frac{(a_{46})^{(9)} G_{45}^{*}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]} \quad , \quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}\ and\ (b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{14}'')^{(1)}}{\partial T_{14}}(T_{14}^{*}) = (q_{14})^{(1)} \quad , \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*}) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G_{13}^{*}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G_{14}^{*}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G_{15}^{*}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\big((b'_{13})^{(1)} - (r_{13})^{(1)}\big)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(13)(j)}T_{13}^{*}\mathbb{G}_j\big) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\big((b'_{14})^{(1)} - (r_{14})^{(1)}\big)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\big(s_{(14)(j)}T_{14}^{*}\mathbb{G}_j\big) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\big((b'_{15})^{(1)} - (r_{15})^{(1)}\big)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\big(s_{(15)(j)}T_{15}^{*}\mathbb{G}_j\big) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(2)}$ and $(b_i'')^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^{*} + \mathbb{G}_i \qquad , T_i = T_i^{*} + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a_{17}'')^{(2)}}{\partial T_{17}}(T_{17}^{*}) = (q_{17})^{(2)} \quad , \frac{\partial(b_i'')^{(2)}}{\partial G_j}((G_{19})^{*}) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\big((a'_{16})^{(2)} + (p_{16})^{(2)}\big)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G_{16}^{*}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\big((a'_{17})^{(2)} + (p_{17})^{(2)}\big)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G_{17}^{*}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\big((a'_{18})^{(2)} + (p_{18})^{(2)}\big)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G_{18}^{*}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(3)}$ *and* $(b_i'')^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \quad , \frac{\partial(b_i'')^{(5)}}{\partial G_j}\left((G_{31})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b_i'')^{(6)}}{\partial G_j}\left((G_{35})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(7)}$ *and* $(b''_i)^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a''_{37})^{(7)}}{\partial T_{37}}(T^*_{37}) = (q_{37})^{(7)} \quad , \frac{\partial(b''_i)^{(7)}}{\partial G_j}\left((G_{39})^{**}\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a'_{36})^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G^*_{36}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a'_{37})^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G^*_{37}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a'_{38})^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G^*_{38}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b'_{36})^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T^*_{36}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b'_{37})^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T^*_{37}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b'_{38})^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T^*_{38}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(8)}$ *and* $(b''_i)^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \quad , \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad (581)$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad (582)$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad (583)$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^*\mathbb{G}_j\right) \quad (584)$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^*\mathbb{G}_j\right) \quad (585)$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^*\mathbb{G}_j\right) \quad (586)$$

(586 A)

ASYMPTOTIC STABILITY ANALYSIS

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}\left((G_{47})^*\right) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right)$ 586

G 587

The characteristic equation of this system is

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b'_{13})^{(1)} + (b'_{14})^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14} + (b_{14})^{(1)}s_{(13),(15)}T^*_{13}\right)\} = 0$$

$$+$$

$$(\lambda)^{(2)} + (b'_{18})^{(2)} - (r_{18})^{(2)}\{\left((\lambda)^{(2)} + (a'_{18})^{(2)} + (p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\right)$$

$$+\left(\left((\lambda)^{(2)} + (a'_{17})^{(2)} + (p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16} + (a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^2 + \left((b'_{16})^{(2)} + (b'_{17})^{(2)} - (r_{16})^{(2)} + (r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^2 + \left((a'_{16})^{(2)} + (a'_{17})^{(2)} + (p_{16})^{(2)} + (p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)} + (a'_{16})^{(2)} + (p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17} + (a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left(\left((\lambda)^{(2)} + (b'_{16})^{(2)} - (r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17} + (b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\right)\} = 0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\Big)$$

$$+\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\Big)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\Big)$$

$$+\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\Big)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\Big)\}=0$$

$$+$$

$$(\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\Big)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\Big)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\Big)$$

$$+\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\Big)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\Big)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\Big)\right]$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\Big)$$

$$+\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\Big)$$

$$\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\Big)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^2+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^2+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^*_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^*_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^*_{45}+(b_{45})^{(9)}s_{(44),(46)}T^*_{44}\right)\}=0$$


And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION FIFTEEN

Proton-Collision Event With A High Di-Muon Mass (1697 Gev), As Seen From The Direction Of The Proton Beams: Age Quod Agis- Agere Sequitur Credere- Agere Sequitur (Esse)- Credula Vitam Spes Fovet Et Melius Cras Fore Semper Dicit Model


INTRODUCTION—VARIABLES USED

(1) EXO-12-061 Event 1: A proton-collision event with a high di-muon mass (1697 GeV), as seen from the direction of the proton beams. Events such as these **form (eb) the** background for Z′




searches.

(2) EXO-12-061 Event 2: A 3D view of a proton-collision event **with a** high di-muon mass (1697 GeV). Events such as these form the background for Z′ searches.




(3) EXO-12-061 Figure 1: The invariant mass spectrum of μ+μ− events. The points with error bars represent the data. The histograms represent the expectations from various Standard Model processes, and include the Z peak at 91 GeV. No evidence for new physics observed which


would manifest itself as a cluster of data above the expected background.




(4) EXO-12-059 Event 1: An event with two wide jets, with the highest mass recorded for the wide-jet pair (5.15 TeV), **as seen (e) from the** direction of the proton beams.




(5) EXO-12-059 Figure 1: Di-jet mass spectrum from wide jets. (The data are compared to a smooth fit and predictions from quantum chromodynamics Monte Carlo (QCD MC) simulation). Simulation of two representatives new physics models, W′ **and** axigluon/colouron (A/C), are shown to indicate the expected scale of contribution from new physics processes. The shaded band **shows (eb) the** contribution from the systematic uncertainty in the jet energy scale (JES).



(6) TOP-12-035 Event 1: An eμ event candidate selected in 8 TeV data, as seen from the direction of the proton beams. The kinematics of the main objects used in the event selection is highlighted: two isolated leptons **and** two particle-flow jets. (The reconstructed missing transverse energy is also displayed for reference).



TOP-12-035 Figure 1: Measuring R from the data. The variation observed in the di-lepton channels is depicted by the dashed lines. The inset shows the inclusive b-tag multiplicity distribution and the result of the fit.



TOP-12-035 Figure 2: Chronological summary of the different measurements of R performed at the Tevatron and the LHC. The shaded region represents the lower endpoint for the confidence interval at 95% confidence level on R obtained after imposing the physical constraint R≤1.

CMS is presenting a wide range of results for the annual winter conference, Rencontres. The first week of Moriond — 3–8 March — is devoted primarily to electroweak (EW) studies. Among the CMS results are further studies of the recently discovered Higgs boson candidate, searches for new physics beyond the Standard Model and precision measurements of the properties of the top quark. We present below brief highlights from a few of the many excellent analyses that will be shown at the conference.

(7) Higgs **decays (eb) into** pairs of Z bosons, W bosons or τ (tau) leptons

In July 2012, CMS and ATLAS particle consistent with the predictions of the Standard Model (SM) Higgs boson. However, in order to confirm whether the particle is indeed the expected Higgs boson requires a more thorough understanding of its properties. The following results all take into account around 5 fb−1 and 20 fb−1 (inverse femtobarn) of proton-collision data recorded by CMS at centre-of-mass energies of 7 and 8 TeV, respectively.

Decays to two Z bosons: HIG-13-002

(8) The study of the new boson's decay **into(eb) two** Z particles is crucial to the determination of two of its quantum numbers, spin and parity

(9) Standard Model **predicts (eb) that the** Higgs boson should have zero spin and positive parity (represented as 0+). CMS recently published a study showing that the data were consistent with 0+ and excluded 0− at 97% confidence level (Phys. Rev. Lett. 110, 081803 (2013)).

NOTATION

Module One

G_{13} : Category one of Z' searches (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

G_{14} : Category two of Z' searches

G_{15} : Category three of Z' searches

T_{13} : Category one of proton-collision event with a high di-muon mass (1697 GeV), as seen from the direction of the proton beams

T_{14} : Category two of proton-collision event with a high di-muon mass (1697 GeV), as seen from the direction of the proton beams

T_{15} : Category three of proton-collision event with a high di-muon mass (1697 GeV), as seen from the direction of the proton beams

Module Two

G_{16} : Category one of invariant mass spectrum of μ+μ− events

G_{17} : Category two of invariant mass spectrum of μ+μ− events

G_{18} : Category three of invariant mass spectrum of μ+μ− events

T_{16} : Category one of expectations from various Standard Model processes (and includes the Z peak at 91 GeV. No evidence for new physics observed which would manifest itself as a cluster of data above the expected background)

T_{17} : Category two of expectations from various Standard Model processes (and includes the Z peak at 91 GeV. No evidence for new physics observed which would manifest itself as a cluster of data above the expected background)

T_{18} : Category three of expectations from various Standard Model processes (and includes the Z peak at 91 GeV. No evidence for new physics observed which would manifest itself as a cluster of data above the expected background)

Module three

G_{20} : Category one of event with two wide jets, with the highest mass recorded for the wide-jet pair (5.15 TeV). Classification is based on various characterstics of the investigating systems such as these. Note this is not the only experiment and there shall be and have been various such experiments' and the characterstics form the bastion of the classification scheme.

G_{21} : Category two of event with two wide jets, with the highest mass recorded for the wide-jet pair (5.15 TeV),

G_{22} : Category three of event with two wide jets, with the highest mass recorded for the wide-jet pair (5.15 TeV),

T_{20} : Category one of direction of the proton beams (For different investigating experiments there would be different direction of proton beams and the concomitant characterstics form the basis of the classification)

T_{21} : Category two of direction of the proton beams

T_{22} : Category three of direction of the proton beams

Module four

G_{24} : Category one of Di-jet mass spectrum from wide jets. (The data are compared to a smooth fit and predictions from quantum chromodynamics Monte Carlo (QCD MC) simulation). Simulation of two representatives new physics models,

G_{25} : Category two of Di-jet mass spectrum from wide jets. (The data are compared to a smooth fit and predictions from quantum chromodynamics Monte Carlo (QCD MC) simulation). Simulation of two representatives new physics models,

G_{26} : Category three of Di-jet mass spectrum from wide jets. (The data are compared to a smooth fit and predictions from quantum chromodynamics Monte Carlo (QCD MC) simulation). Simulation of two representatives new physics models,

T_{24} : Category one of axigluon/colouron (A/C), are shown to indicate the expected scale of contribution from new physics processes. The shaded band **shows (eb) the** contribution from the systematic uncertainty in the jet energy scale (JES).

T_{25} : Category two of axigluon/colouron (A/C), are shown to indicate the expected scale of contribution from new physics processes. The shaded band **shows (eb) the** contribution from the systematic uncertainty in the jet energy scale (JES).

T_{26} : Category three of axigluon/colouron (A/C), are shown to indicate the expected scale of contribution from new physics processes. The shaded band **shows (eb) the** contribution from the systematic uncertainty in the jet energy scale (JES).

Module five

G_{28} : Category one of two isolated leptons

G_{29} : Category two of two isolated leptons

G_{30} : Category three of two isolated leptons

T_{28} : Category one of two particle-flow jets. (The reconstructed missing transverse energy is also displayed for reference).

T_{29} : Category two of two particle-flow jets. (The reconstructed missing transverse energy is also displayed for reference).

T_{30} : Category three of two particle-flow jets. (The reconstructed missing transverse energy is also displayed for reference).

Module six

G_{32} : Category one of Higgs (Higgs is the same. In different experiments' you shall find Higgs. Experimental characterstics and environment form the basis of classification. It should not be confused that there are three types of Higgs or there are three Higgs)

G_{33} : Category two of Higgs

G_{34} : Category three of Higgs

T_{32} : Category one of pairs of Z bosons, W bosons or τ (tau) leptons

T_{33} : Category two of pairs of Z bosons, W bosons or τ (tau) leptons

T_{34} : Category three of pairs of Z bosons, W bosons or τ (tau) leptons

Module seven

G_{36} : Category one of new boson's decay

G_{37} : Category two of new boson's decay

G_{38} : Category three of new boson's decay

T_{36} : Category one of two Z particles (There are many such "two particles' in different experiments. Experimental characterstics form the basis for the stratification strategy)

T_{37} : Category two of two Z particles

T_{38} : Category three of two Z particles

Module eight

G_{40} : Category one of Result of Module seven. There are various results for different experiments.

G_{41} : Category two of Result of Module seven. There are various results for different experiments

G_{42} : Category three of Result of Module seven. There are various results for different experiments

T_{40} : Category one of determination of two of its quantum numbers, spin and parity
T_{41} : Category two of determination of two of its quantum numbers, spin and parity
T_{42} : Category three of determination of two of its quantum numbers, spin and parity

Module Nine

G_{44} : Category one of Standard Model. The Standard Model of particle physics is a theory concerning the electromagnetic, weak, and strong nuclear interactions, which mediate the dynamics of the known subatomic particles. Developed throughout the mid to late 20th century, the Standard Model is truly "a tapestry woven by many hands", sometimes driven forward by new experimental discoveries, sometimes by theoretical advances. It was a collaborative effort in the largest sense, spanning continents and decades. The current formulation was finalized in the mid-1970s upon experimental confirmation of the existence of quarks. Since then, discoveries of the bottom quark (1977), the top quark (1995), and the tau neutrino (2000) have given further credence to the Standard Model. More recently, (2011–2012) the detection of the Higgs boson completes the set of predicted particles. Because of its success in explaining a wide variety of experimental results, the Standard Model is sometimes regarded as a "theory of almost everything".

G_{45} : Category two of Standard Model. Again it is to be said that we are talking of the systems for which Standard model hold. These systems have certain characterstics. These characterstics form the basis of the classification.

G_{46} : Category three of Standard Model

T_{44} : Category one of Higgs boson should have zero spin and positive parity (represented as 0+). CMS recently published a study showing that the data were consistent with 0+ and excluded 0− at 97% confidence level (Phys. Rev. Lett. 110, 081803 (2013)). Spin and positive parity is indeed same for various experiments. We again emphasise that the characterstics of the experiment form the basis for the classification scheme.

T_{45} : Category two of Higgs boson should have zero spin and positive parity (represented as 0+). CMS recently published a study showing that the data were consistent with 0+ and excluded 0− at 97% confidence level (Phys. Rev. Lett. 110, 081803 (2013)).

T_{46} : Category three of Higgs boson should have zero spin and positive parity (represented as 0+). CMS recently published a study showing that the data were consistent with 0+ and excluded 0− at 97% confidence level (Phys. Rev. Lett. 110, 081803 (2013)).

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right]G_{13} \quad 1$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}, t)\right]G_{14} \quad 2$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}, t)\right]G_{15} \quad 3$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G, t)\right]T_{13} \quad 4$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G, t)\right]T_{14} \quad 5$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G, t)\right]T_{15} \quad 6$$

$+(a''_{13})^{(1)}(T_{14}, t) =$ First augmentation factor
$-(b''_{13})^{(1)}(G, t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right]G_{16} \quad 7$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}, t)\right]G_{17} \quad 8$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}, t)\right]G_{18} \quad 9$$

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19}), t)\right]T_{16} \quad 10$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}((G_{19}), t)\right]T_{17} \quad 11$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19}), t)\right]T_{18} \quad 12$$

$+(a''_{16})^{(2)}(T_{17}, t) =$ First augmentation factor

$-(b''_{16})^{(2)}((G_{19}), t) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right]G_{20} \quad 13$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}, t)\right]G_{21} \quad 14$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}, t)\right]G_{22} \quad 15$$

$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}, t)\right]T_{20} \quad 16$$

$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}, t)\right]T_{21} \quad 17$$

$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}, t)\right]T_{22} \quad 18$$

$+(a''_{20})^{(3)}(T_{21}, t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23}, t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right]G_{24} \quad 19$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}, t)\right]G_{25} \quad 20$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}, t)\right]G_{26} \quad 21$$

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27}), t)\right]T_{24} \quad 22$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}((G_{27}), t)\right]T_{25} \quad 23$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}), t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25}, t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}), t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}, t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}, t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}), t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}), t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}), t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29}, t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}), t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}, t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}, t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}, t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}), t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}), t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}), t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33}, t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}, t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}, t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}), t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}), t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}), t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37}, t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}, t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}, t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}), t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}((G_{43}), t)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43}), t)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}, t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}, t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47}), t)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}((G_{47}), t)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47}), t)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45}, t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}((G_{47}), t) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)}\boxed{+(a''_{13})^{(1)}(T_{14},t)}\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)}\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad (55)$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)}\boxed{+(a''_{14})^{(1)}(T_{14},t)}\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad (56)$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)}\boxed{+(a''_{15})^{(1)}(T_{14},t)}\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad (57)$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)}\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad (58)$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)}\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad (59)$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)}\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad (60)$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{array}\right] G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[\begin{array}{ccc} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{array}\right] G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[\begin{array}{ccc} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{array}\right] G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,)}(T_{14},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)}$ are eight augmentation coefficient

for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)}\boxed{-(b''_{16})^{(2)}(G_{19},t)} & \boxed{-(b''_{13})^{(1,1,)}(G,t)} & \boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)}\boxed{-(b''_{17})^{(2)}(G_{19},t)} & \boxed{-(b''_{14})^{(1,1,)}(G,t)} & \boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)}\boxed{-(b''_{18})^{(2)}(G_{19},t)} & \boxed{-(b''_{15})^{(1,1,)}(G,t)} & \boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9)}(G_{47},t)} \end{bmatrix} T_{18} \quad 66$$

where $\boxed{-(b''_{16})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2)}(G_{19},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{13})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,)}(G,t)}$ are second detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{20})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{36})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{40})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \quad 67$$

$$(a_{20})^{(3)}G_{21} - \begin{bmatrix} (a'_{20})^{(3)}\boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \begin{bmatrix} (a'_{21})^{(3)}\boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{21} \quad 68$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \begin{bmatrix} (a'_{22})^{(3)}\boxed{+(a''_{22})^{(3)}(T_{21},t)}\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{bmatrix} G_{22} \quad 69$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \quad 70$$

$$(b_{20})^{(3)}T_{21} - \begin{bmatrix} (b'_{20})^{(3)}\boxed{-(b''_{20})^{(3)}(G_{23},t)}\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{20}$$

$$\frac{dT_{21}}{dt} = \quad 71$$

$$(b_{21})^{(3)}T_{20} - \begin{bmatrix} (b'_{21})^{(3)}\boxed{-(b''_{21})^{(3)}(G_{23},t)}\boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{21}$$

$$\frac{dT_{22}}{dt} = \quad 72$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)}\boxed{-(b''_{22})^{(3)}(G_{23},t)}\boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}\boxed{-(b''_{15})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)}\boxed{-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)}\boxed{-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7)}(G_{39},t)}\boxed{-(b''_{42})^{(8,8,8,8)}(G_{43},t)}\boxed{-(b''_{46})^{(9,9,9)}(G_{47},t)} \end{bmatrix} T_{22}$$

$\boxed{-(b''_{20})^{(3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3)}(G_{23},t)}$ *are first detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2)}(G_{19},t)}$ *are second detrition coefficients for category 1, 2 and 3*

$\boxed{-(b''_{13})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,)}(G,t)}$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ & +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) \\ & +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3

$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\boxed{-(b''_{24})^{(4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\boxed{-(b''_{25})^{(4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\boxed{-(b''_{26})^{(4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{26} \quad (78)$$

Where $\boxed{-(b''_{24})^{(4)}(G_{27},t)}, \boxed{-(b''_{25})^{(4)}(G_{27},t)}, \boxed{-(b''_{26})^{(4)}(G_{27},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{29})^{(5,5,)}(G_{31},t)}, \boxed{-(b''_{30})^{(5,5,)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{32})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{33})^{(6,6,)}(G_{35},t)}, \boxed{-(b''_{34})^{(6,6,)}(G_{35},t)}$ *are third detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{14})^{(1,1,1,1)}(G,t)}, \boxed{-(b''_{15})^{(1,1,1,1)}(G,t)}$
are fourth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{16})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{17})^{(2,2,2,2)}(G_{19},t)}, \boxed{-(b''_{18})^{(2,2,2,2)}(G_{19},t)}$
are fifth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{20})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{21})^{(3,3,3,3)}(G_{23},t)}, \boxed{-(b''_{22})^{(3,3,3,3)}(G_{23},t)}$
are sixth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)}, \boxed{-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)}$
are seventh detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)}, \boxed{-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)}$
are eighth detrition coefficients for category 1, 2 *and* 3

$\boxed{-(b''_{46})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{45})^{(9,9,9,9)}(G_{47},t)}, \boxed{-(b''_{44})^{(9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\boxed{+(a''_{28})^{(5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \begin{bmatrix} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \begin{bmatrix} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation*

$coefficient\ for\ category\ 1,2\ and\ 3$

$+(a''_{32})^{(6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)$ are fourth augmentation coefficients for category 1,2, and 3

$+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)$ are fifth augmentation coefficients for category 1,2,and 3

$+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)$ are sixth augmentation coefficients for category 1,2, 3

$+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)$ are seventh augmentation coefficients for category 1,2, 3

$+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)$ are eighth augmentation coefficients for category 1,2, 3

$+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \begin{bmatrix} (b'_{28})^{(5)} & -(b''_{28})^{(5)}(G_{31},t) & -(b''_{24})^{(4,4,)}(G_{27},t) & -(b''_{32})^{(6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1)}(G,t) & -(b''_{16})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{28} \quad 82$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \begin{bmatrix} (b'_{29})^{(5)} & -(b''_{29})^{(5)}(G_{31},t) & -(b''_{25})^{(4,4,)}(G_{27},t) & -(b''_{33})^{(6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1)}(G,t) & -(b''_{17})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{29} \quad 83$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \begin{bmatrix} (b'_{30})^{(5)} & -(b''_{30})^{(5)}(G_{31},t) & -(b''_{26})^{(4,4,)}(G_{27},t) & -(b''_{34})^{(6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,)}(G,t) & -(b''_{18})^{(2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3)}(G_{23},t) \\ & -(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{30} \quad 84$$

where $-(b''_{28})^{(5)}(G_{31},t)$, $-(b''_{29})^{(5)}(G_{31},t)$, $-(b''_{30})^{(5)}(G_{31},t)$ *are first detrition coefficients for category* 1, 2 *and* 3

$-(b''_{24})^{(4,4,)}(G_{27},t)$, $-(b''_{25})^{(4,4,)}(G_{27},t)$, $-(b''_{26})^{(4,4,)}(G_{27},t)$ *are second detrition coefficients for category* 1,2 *and* 3

$-(b''_{32})^{(6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6)}(G_{35},t)$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition

coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}\boxed{+(a''_{32})^{(6)}(T_{33},t)} & \boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}\boxed{+(a''_{33})^{(6)}(T_{33},t)} & \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}\boxed{+(a''_{34})^{(6)}(T_{33},t)} & \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{34}$$

$\boxed{+(a''_{32})^{(6)}(T_{33},t)}, \boxed{+(a''_{33})^{(6)}(T_{33},t)}, \boxed{+(a''_{34})^{(6)}(T_{33},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{28})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{29})^{(5,5,5)}(T_{29},t)}, \boxed{+(a''_{30})^{(5,5,5)}(T_{29},t)}$ *are second augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{24})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{25})^{(4,4,4,)}(T_{25},t)}, \boxed{+(a''_{26})^{(4,4,4,)}(T_{25},t)}$ *are third augmentation coefficients for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)}, \boxed{+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)}$ - are fourth augmentation coefficients

$\boxed{+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)}, \boxed{+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)}$ - fifth augmentation coefficients

$\boxed{+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)}, \boxed{+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)}$ sixth augmentation coefficients

$\boxed{+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)}, \boxed{+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)}$ seventh augmentation coefficients

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)}, \boxed{+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)}$ Eighth augmentation coefficients

$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}, \boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \quad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \quad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)}G_{37} - \begin{bmatrix} (a'_{36})^{(7)}\boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)}G_{36} - \begin{bmatrix} (a'_{37})^{(7)}\boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$\frac{dG_{38}}{dt}$ 93

$= (a_{38})^{(7)} G_{37}$

$$- \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$\frac{dT_{36}}{dt} =$ 94

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$\frac{dT_{37}}{dt} =$

$$(b_{37})^{(7)} T_{36} - \begin{bmatrix} (b'_{37})^{(7)} \boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$\frac{dT_{38}}{dt} =$

$$(b_{38})^{(7)} T_{37} - \begin{bmatrix} (b'_{38})^{(7)} \boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second

detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)$, $-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)} & +(a''_{40})^{(8)}(T_{41},t) & +(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)} & +(a''_{41})^{(8)}(T_{41},t) & +(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \begin{bmatrix} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) \\ & +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) & +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) \\ & +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)}T_{40} -$$
$$\begin{bmatrix} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)}T_{41} -$$
$$\begin{bmatrix} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) & -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) \\ & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) \\ & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) & -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) \end{bmatrix} T_{15}$$

Where $-(b''_{36})^{(7)}(G_{39},t)$, $-(b''_{37})^{(7)}(G_{39},t)$, $-(b''_{38})^{(7)}(G_{39},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,)}(G_{35},t)$, $-(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t)$ are sixth detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t)$, $-(b''_{38})^{(7,7,)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t)$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45}$$

$$-\left[\begin{array}{ccc} (a'_{44})^{(9)}\boxed{+(a''_{44})^{(9)}(T_{45},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{13}$$

$$\frac{dG_{45}}{dt}$$

$$= (a_{45})^{(9)}G_{44}$$

$$-\left[\begin{array}{ccc} (a'_{45})^{(9)}\boxed{+(a''_{45})^{(9)}(T_{45},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{14}$$

$$\frac{dG_{46}}{dt}$$

$$= (a_{46})^{(9)}G_{45}$$

$$-\left[\begin{array}{ccc} (a'_{46})^{(9)}\boxed{+(a''_{46})^{(9)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)} \end{array}\right]G_{15}$$

Where $\boxed{+(a''_{44})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)}$ are Seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$

$$(b_{44})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{44})^{(9)} & \boxed{-(b''_{44})^{(9)}(G_{47},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$

$$(b_{45})^{(9)}T_{44} -$$

$$\begin{bmatrix} (b'_{45})^{(9)} & \boxed{-(b''_{45})^{(9)}(G_{47},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$

$$(b_{46})^{(9)}T_{45} -$$

$$\begin{bmatrix} (b'_{46})^{(9)} & \boxed{-(b''_{46})^{(9)}(G_{47},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)} & \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)} & \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{44})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9)}(G_{47},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a'_i)^{(1)}, (a''_i)^{(1)}, (b_i)^{(1)}, (b'_i)^{(1)}, (b''_i)^{(1)} > 0, \quad i,j = 13,14,15 \qquad (97)$$

The functions $(a''_i)^{(1)}, (b''_i)^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a''_i)^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b''_i)^{(1)}(G,t) \le (r_i)^{(1)} \le (b'_i)^{(1)} \le (\hat{B}_{13})^{(1)}$$

$lim_{T_2\to\infty}(a_i'')^{(1)}\,(T_{14},t)=(p_i)^{(1)}$ 98

$\lim_{G\to\infty}(b_i'')^{(1)}\,(G,t)=\ \ (r_i)^{(1)}$

Definition of $(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)},(\hat{B}_{13})^{(1)},(p_i)^{(1)},\ \ (r_i)^{(1)}}$ are positive constants and $\boxed{i=13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t)-(a_i'')^{(1)}(T_{14},t)|\le(\hat{k}_{13})^{(1)}|T_{14}-T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t)-(b_i'')^{(1)}(G,t)|<(\hat{k}_{13})^{(1)}||G-G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and $(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $\left[(\hat{k}_{13})^{(1)},(\hat{M}_{13})^{(1)}\right]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)}=1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},$ are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}\ ,\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}}<1$$

Definition of $(\hat{P}_{13})^{(1)},(\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)},(\hat{k}_{13})^{(1)},(\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)},(a_i')^{(1)},(b_i)^{(1)},(b_i')^{(1)},(p_i)^{(1)},\ (r_i)^{(1)},i=13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\,(a_i)^{(1)}+(a_i')^{(1)}+\ (\hat{A}_{13})^{(1)}+(\hat{P}_{13})^{(1)}\,(\hat{k}_{13})^{(1)}]<1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[\ (b_i)^{(1)}+(b_i')^{(1)}+\ (\hat{B}_{13})^{(1)}+(\hat{Q}_{13})^{(1)}\ (\hat{k}_{13})^{(1)}]<1$$

Where we suppose

$$(a_i)^{(2)},(a_i')^{(2)},(a_i'')^{(2)},(b_i)^{(2)},(b_i')^{(2)},(b_i'')^{(2)}>0,\quad i,j=16,17,18$$

The functions $(a_i'')^{(2)},(b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(\mathrm{p_i})^{(2)}$, $(\mathrm{r_i})^{(2)}$:

$$(a_i'')^{(2)}(T_{17},t)\le(p_i)^{(2)}\le\left(\hat{A}_{16}\right)^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19},t)\le\ \ (r_i)^{(2)}\le(b_i')^{(2)}\le(\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2\to\infty}(a_i'')^{(2)}\,(T_{17},t)=(p_i)^{(2)} \quad 104$$

$$\lim_{G\to\infty}(b_i'')^{(2)}\left((G_{19}),t\right)=\ \ (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and$(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $\left[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}\right]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, \ (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(b_i)^{(2)} + (b_i')^{(2)} + (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 111$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i, j = 20,21,22 \quad 112$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}(T_{21}, t) = (p_i)^{(3)} \quad 113$$

$$\lim_{G \to \infty}(b_i'')^{(3)}(G_{23}, t) = (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: 114

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: 115

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$, are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and (\hat{B}_{20})^{(3)}$ and the constants $(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, (r_i)^{(3)}, i = 20,21,22,$ satisfy the inequalities 116

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(a_i)^{(3)} + (a_i')^{(3)} + (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[(b_i)^{(3)} + (b_i')^{(3)} + (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}(\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i, j = 24,25,26$$ 117

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}((G_{27}), t) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)}$$ 118

$$\lim_{G \to \infty}(b_i'')^{(4)}((G_{27}), t) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: 119

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})',t)-(b_i'')^{(4)}\big((G_{27}),t\big)|<(\hat{k}_{24})^{(4)}||(G_{27})-(G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}',t)$ and$(a_i'')^{(4)}(T_{25},t)$. (T_{25}',t) and (T_{25},t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)},(\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)}=1$ then the function $(a_i'')^{(4)}(T_{25},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$: 120

$(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}},\frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}}<1$$

Definition of $(\hat{P}_{24})^{(4)},(\hat{Q}_{24})^{(4)}$: 121

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)},(\hat{k}_{24})^{(4)},(\hat{A}_{24})^{(4)}and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)},(a_i')^{(4)},(b_i)^{(4)},(b_i')^{(4)},(p_i)^{(4)},\ (r_i)^{(4)},i=24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\,(a_i)^{(4)}+(a_i')^{(4)}+\ (\hat{A}_{24})^{(4)}+\ (\hat{P}_{24})^{(4)}\,(\hat{k}_{24})^{(4)}]<1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[\ (b_i)^{(4)}+(b_i')^{(4)}+\ (\hat{B}_{24})^{(4)}+\ (\hat{Q}_{24})^{(4)}\ (\hat{k}_{24})^{(4)}]<1$$

Where we suppose

$$(a_i)^{(5)},(a_i')^{(5)},(a_i'')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(b_i'')^{(5)}>0,\quad i,j=28,29,30 \qquad 122$$

The functions $(a_i'')^{(5)},(b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)},\ (r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t)\le(p_i)^{(5)}\le(\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}\big((G_{31}),t\big)\le\ (r_i)^{(5)}\le(b_i')^{(5)}\le(\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}\,(T_{29},t)=(p_i)^{(5)} \qquad 123$$

$$\lim_{G\to\infty}(b_i'')^{(5)}\,(G_{31},t)=\ (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)},(\hat{B}_{28})^{(5)},(p_i)^{(5)},\ (r_i)^{(5)}}$ are positive constants and $\boxed{i=28,29,30}$

They satisfy Lipschitz condition: 124

$$|(a_i'')^{(5)}(T_{29}',t)-(a_i'')^{(5)}(T_{29},t)|\le(\hat{k}_{28})^{(5)}|T_{29}-T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t)-(b_i'')^{(5)}\big((G_{31}),t\big)|<(\hat{k}_{28})^{(5)}||(G_{31})-(G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$

and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)},(\widehat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{28})^{(5)}=1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$: 125

$(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\ ,\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}<1$$

Definition of $(\hat{P}_{28})^{(5)},(\hat{Q}_{28})^{(5)}$: 126

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\widehat{M}_{28})^{(5)},(\hat{k}_{28})^{(5)},(\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)},(a_i')^{(5)},(b_i)^{(5)},(b_i')^{(5)},(p_i)^{(5)},\ (r_i)^{(5)},i=28,29,30,$ satisfy the inequalities

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\,(a_i)^{(5)}+(a_i')^{(5)}+\ (\hat{A}_{28})^{(5)}+\ (\hat{P}_{28})^{(5)}\,(\hat{k}_{28})^{(5)}]<1$$

$$\frac{1}{(\widehat{M}_{28})^{(5)}}[\ (b_i)^{(5)}+(b_i')^{(5)}+\ (\hat{B}_{28})^{(5)}+\ (\hat{Q}_{28})^{(5)}\ (\hat{k}_{28})^{(5)}]<1$$

Where we suppose

$$(a_i)^{(6)},(a_i')^{(6)},(a_i'')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(b_i'')^{(6)}>0,\quad i,j=32,33,34 \qquad 127$$

The functions $(a_i'')^{(6)},(b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)},\ (r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t)\le(p_i)^{(6)}\le(\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t)\le\ (r_i)^{(6)}\le(b_i')^{(6)}\le(\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\left((G_{35}),t\right)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\widehat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\left((G_{35}),t\right)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\widehat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\left[(\hat{k}_{32})^{(6)},(\widehat{M}_{32})^{(6)}\right]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to

the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}, \frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}} < 1$$

Definition of $(\hat{P}_{32})^{(6)}, (\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)}, (\hat{k}_{32})^{(6)}, (\hat{A}_{32})^{(6)} and (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)}, (a_i')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (p_i)^{(6)}, (r_i)^{(6)}, i = 32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(a_i)^{(6)} + (a_i')^{(6)} + (\hat{A}_{32})^{(6)} + (\hat{P}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[(b_i)^{(6)} + (b_i')^{(6)} + (\hat{B}_{32})^{(6)} + (\hat{Q}_{32})^{(6)} (\hat{k}_{32})^{(6)}] < 1$$

Where we suppose

) $(a_i)^{(7)}, (a_i')^{(7)}, (a_i'')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (b_i'')^{(7)} > 0, \quad i, j = 36,37,38$ 131

)The functions $(a_i'')^{(7)}, (b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)}$, $(r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37}, t) \le (p_i)^{(7)} \le (\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39}, t) \le (r_i)^{(7)} \le (b_i')^{(7)} \le (\hat{B}_{36})^{(7)}$$ 132

$\lim_{T_2 \to \infty} (a_i'')^{(7)}(T_{37}, t) = (p_i)^{(7)}$

$\lim_{G \to \infty} (b_i'')^{(7)}((G_{39}), t) = (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36,37,38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \le (\hat{k}_{36})^{(7)} |T_{37} - T_{37}'| e^{-(\hat{M}_{36})^{(7)} t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}((G_{39}), t)| < (\hat{k}_{36})^{(7)} ||(G_{39}) - (G_{39})'|| e^{-(\hat{M}_{36})^{(7)} t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and $(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if

$(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

) $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)},$ are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}} , \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, (r_i)^{(7)}, i = 36,37,38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(a_i)^{(7)} + (a_i')^{(7)} + (\hat{A}_{36})^{(7)} + (\hat{P}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[(b_i)^{(7)} + (b_i')^{(7)} + (\hat{B}_{36})^{(7)} + (\hat{Q}_{36})^{(7)} (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40,41,42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}$, $(r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$$\lim_{T_2 \to \infty} (a_i'')^{(8)}(T_{41}, t) = (p_i)^{(8)} \qquad 140$$

$$\lim_{G \to \infty} (b_i'')^{(8)}((G_{43}), t) = (r_i)^{(8)} \qquad 141$$

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(8)}(T_{41}', t) - (a_i'')^{(8)}(T_{41}, t)| \leq (\hat{k}_{40})^{(8)} |T_{41} - T_{41}'| e^{-(\hat{M}_{40})^{(8)} t} \qquad 142$$

$$|(b_i'')^{(8)}((G_{43})', t) - (b_i'')^{(8)}((G_{43}), t)| < (\hat{k}_{40})^{(8)} ||(G_{43}) - (G_{43})'|| e^{-(\hat{M}_{40})^{(8)} t} \qquad 143$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}', t)$ and $(a_i'')^{(8)}(T_{41}, t)$. (T_{41}', t) and (T_{41}, t) are points belonging to the interval $[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}]$. It

is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)}=1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},$ are positive constants

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\ ,\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}<1 \qquad 144$$

Definition of $(\hat{P}_{40})^{(8)},(\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)},(\hat{k}_{40})^{(8)},(\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)},(a_i')^{(8)},(b_i)^{(8)},(b_i')^{(8)},(p_i)^{(8)},\ (r_i)^{(8)},i=40,41,42,$
Satisfy the inequalities

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)}+(a_i')^{(8)}+(\hat{A}_{40})^{(8)}+(\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 145$$

$$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)}+(b_i')^{(8)}+(\hat{B}_{40})^{(8)}+(\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}]<1 \qquad 146$$

Where we suppose

$$(a_i)^{(9)},(a_i')^{(9)},(a_i'')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(b_i'')^{(9)}>0,\quad i,j=44,45,46 \qquad 146\,A$$

The functions $(a_i'')^{(9)},(b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)},\ (r_i)^{(9)}$:

$$(a_i'')^{(9)}(T_{45},t)\le(p_i)^{(9)}\le(\hat{A}_{44})^{(9)}$$

$$(b_i'')^{(9)}(G_{47},t)\le\ (r_i)^{(9)}\le(b_i')^{(9)}\le(\hat{B}_{44})^{(9)}$$

$$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$$

$$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44,45,46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$

and $(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\widehat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\widehat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$, are positive constants

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\widehat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)} and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44,45,46,$
satisfy the inequalities

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\widehat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,,T_i(0)$:

$$G_i(t)\le\ (\hat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le\ (\hat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,,T_i(0)$

$$G_i(t)\le\ (\hat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t)\le\ (\hat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad G_i(0)=G_i^0>0$$

$$T_i(t)\le\ (\hat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}\ ,\quad T_i(0)=T_i^0>0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,,T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 B

Definition of $G_i(0)$, $T_i(0)$:

$G_i(t) \leq \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{G_i(0) = G_i^0 > 0}$

$T_i(t) \leq \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t}$, $\boxed{T_i(0) = T_i^0 > 0}$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq \left(\hat{P}_{13}\right)^{(1)}$, $T_i^0 \leq \left(\hat{Q}_{13}\right)^{(1)}$, 155

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 156$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{13})^{(1)} e^{(\bar{M}_{13})^{(1)} t} \quad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)} (T_{14}(s_{(13)}), s_{(13)}) \right) G_{15}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{13}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{14}(s_{(13)}) \right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)} (G(s_{(13)}), s_{(13)}) \right) T_{15}(s_{(13)}) \right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{16})^{(2)}, T_i^0 \le (\hat{Q}_{16})^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{16})^{(2)} e^{(\bar{M}_{16})^{(2)} t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)} (T_{17}(s_{(16)}), s_{(17)}) \right) G_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)} (T_{17}(s_{(16)}), s_{(16)}) \right) G_{18}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{16}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{17}(s_{(16)}) \right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)} (G_{19}(s_{(16)}), s_{(16)}) \right) T_{18}(s_{(16)}) \right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i, $T_i: \mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$$G_i(0) = G_i^0\ ,\ T_i(0) = T_i^0\ ,\ G_i^0 \le (\hat{P}_{20})^{(3)}\ , T_i^0 \le (\hat{Q}_{20})^{(3)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)}t}$$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions $G_i\ ,\ T_i: \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\ ,\ T_i(0) = T_i^0\ ,\ G_i^0 \le (\hat{P}_{24})^{(4)}\ , T_i^0 \le (\hat{Q}_{24})^{(4)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)}t}$$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0,t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{28})^{(5)}$, $T_i^0 \leq (\hat{Q}_{28})^{(5)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)} (T_{29}(s_{(28)}), s_{(28)}) \right) G_{30}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{28}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{29}(s_{(28)}) \right] ds_{(28)}$$

$$\bar{T}_{30}(t) = T_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)} (G_{31}(s_{(28)}), s_{(28)}) \right) T_{30}(s_{(28)}) \right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0,t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{32})^{(6)}$, $T_i^0 \leq (\hat{Q}_{32})^{(6)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)} (T_{33}(s_{(32)}), s_{(32)}) \right) G_{34}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{32}(s_{(32)}) \right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)} (G_{35}(s_{(32)}), s_{(32)}) \right) T_{33}(s_{(32)}) \right] ds_{(32)}$$

$$\overline{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^{0} + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\bar{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\overline{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^{0} + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{40})^{(8)}$, $T_i^0 \le (\hat{Q}_{40})^{(8)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{40})^{(8)} e^{(\bar{M}_{40})^{(8)} t}$$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{\mathrm{T}}_{42}(\mathrm{t}) = \mathrm{T}_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i, T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$$G_i(0) = G_i^0,\ T_i(0) = T_i^0,\ G_i^0 \le (\hat{P}_{44})^{(9)},\ T_i^0 \le (\hat{Q}_{44})^{(9)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^0 + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47}(s_{(44)}), s_{(44)})\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} \left(G_{14}^0 + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^0 + \frac{(a_{13})^{(1)} (\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}} \left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t)-G_{13}^{0})e^{-(\hat{M}_{13})^{(1)}t}\leq\frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[((\hat{P}_{13})^{(1)}+G_{14}^{0})e^{\left(-\frac{(\hat{P}_{13})^{(1)}+G_{14}^{0}}{G_{14}^{0}}\right)}+(\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t)\leq G_{16}^{0}+\int_{0}^{t}\left[(a_{16})^{(2)}\left(G_{17}^{0}+(\hat{P}_{16})^{(6)}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\right)\right]ds_{(16)}=\left(1+(a_{16})^{(2)}t\right)G_{17}^{0}+\frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)}t}-1\right) \quad (169)$$

From which it follows that (170)

$$(G_{16}(t)-G_{16}^{0})e^{-(\hat{M}_{16})^{(2)}t}\leq\frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[((\hat{P}_{16})^{(2)}+G_{17}^{0})e^{\left(-\frac{(\hat{P}_{16})^{(2)}+G_{17}^{0}}{G_{17}^{0}}\right)}+(\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that (171)

$$G_{20}(t)\leq G_{20}^{0}+\int_{0}^{t}\left[(a_{20})^{(3)}\left(G_{21}^{0}+(\hat{P}_{20})^{(3)}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\right)\right]ds_{(20)}=$$

$$\left(1+(a_{20})^{(3)}t\right)G_{21}^{0}+\frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)}t}-1\right)$$

From which it follows that (172)

$$(G_{20}(t)-G_{20}^{0})e^{-(\hat{M}_{20})^{(3)}t}\leq\frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[((\hat{P}_{20})^{(3)}+G_{21}^{0})e^{\left(-\frac{(\hat{P}_{20})^{(3)}+G_{21}^{0}}{G_{21}^{0}}\right)}+(\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that (173)

$$G_{24}(t)\leq G_{24}^{0}+\int_{0}^{t}\left[(a_{24})^{(4)}\left(G_{25}^{0}+(\hat{P}_{24})^{(4)}e^{(\hat{M}_{24})^{(4)}s_{(24)}}\right)\right]ds_{(24)}=$$

$$\left(1+(a_{24})^{(4)}t\right)G_{25}^{0}+\frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)}t}-1\right)$$

From which it follows that (174)

$$(G_{24}(t)-G_{24}^{0})e^{-(\hat{M}_{24})^{(4)}t}\leq\frac{(a_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t)\leq G_{28}^{0}+\int_{0}^{t}\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\hat{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)}=$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left(e^{(\hat{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\hat{M}_{28})^{(5)}t}\leq\frac{(a_{28})^{(5)}}{(\hat{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t)\leq G_{32}^{0}+\int_{0}^{t}\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\hat{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)}=$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left(e^{(\hat{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\hat{M}_{32})^{(6)}t}\leq\frac{(a_{32})^{(6)}}{(\hat{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t)\leq G_{36}^{0}+\int_{0}^{t}\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\hat{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)}=$$
$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left(e^{(\hat{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\hat{M}_{36})^{(7)}t}\leq\frac{(a_{36})^{(7)}}{(\hat{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t)\leq G_{40}^{0}+\int_{0}^{t}\left[(a_{40})^{(8)}\left(G_{41}^{0}+(\hat{P}_{40})^{(8)}e^{(\hat{M}_{40})^{(8)}s_{(40)}}\right)\right]ds_{(40)}=$$ 180
$$\left(1+(a_{40})^{(8)}t\right)G_{41}^{0}+\frac{(a_{40})^{(8)}(\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left(e^{(\hat{M}_{40})^{(8)}t}-1\right)$$

From which it follows that 181

$$(G_{40}(t)-G_{40}^{0})e^{-(\hat{M}_{40})^{(8)}t}\leq\frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}}\left[((\hat{P}_{40})^{(8)}+G_{41}^{0})e^{\left(-\frac{(\hat{P}_{40})^{(8)}+G_{41}^{0}}{G_{41}^{0}}\right)}+(\hat{P}_{40})^{(8)}\right]$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \leq G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \leq \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose 182

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{13})^{(1)} \quad 183$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \leq (\hat{Q}_{13})^{(1)} \quad 184$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric 185

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\}$$

Indeed if we denote

Definition of $\tilde{G}, \tilde{T} : \left(\tilde{G}, \tilde{T}\right) = \mathcal{A}^{(1)}(G, T)$

It results

$$\left|\tilde{G}_{13}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{13})^{(1)} \left|G_{14}^{(1)} - G_{14}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} ds_{(13)} +$$

$$\int_0^t \{(a'_{13})^{(1)} \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{-(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$$(a''_{13})^{(1)} \left(T_{14}^{(1)}, s_{(13)}\right) \left|G_{13}^{(1)} - G_{13}^{(2)}\right| e^{-(\hat{M}_{13})^{(1)} s_{(13)}} e^{(\hat{M}_{13})^{(1)} s_{(13)}} +$$

$G_{13}^{(2)}|(a_{13}'')^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a_{13}'')^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad (186)$$
$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+(a_{13}')^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a_{13}'')^{(1)}$ and $(b_{13}'')^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(1)}$ and $(b_i'')^{(1)}, i=13,14,15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(1)}-(a_i'')^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{(-(b_i')^{(1)}t)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: (187)

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a_{14}')^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a_{14}')^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a_{15}')^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If $G_{13}\ \ is$ bounded, from below, the same property holds for $G_{14}\ and\ G_{15}$. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. (188)

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}\ (G(t),t))=(b_{14}')^{(1)}$ then $T_{14}\ \to\infty$. (189)

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t\ >t_1$

$$(b_{14})^{(1)}-(b_i'')^{(1)}(G(t),t)<\varepsilon_1, T_{13}\ (t)>(m)^{(1)}$$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t), t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[(\widehat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\widehat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)}, (T_{19})^{(1)}), ((G_{19})^{(2)}, (T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$\left|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{16})^{(2)}\left|G_{17}^{(1)} - G_{17}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$$

$$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right|\ e^{-(\widehat{M}_{16})^{(2)}s_{(16)}}e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\}ds_{(16)}$$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right|e^{-(\widehat{M}_{16})^{(2)}t} \leq$$

$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a'_{16})^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right)\mathrm{d}\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a''_{16})^{(2)}$ and $(b''_{16})^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)}e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a''_i)^{(2)}$ and $(b''_i)^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a'_i)^{(2)} - (a''_i)^{(2)}(T_{17}(s_{(16)}), s_{(16)})\}\mathrm{d}s_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b'_i)^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a'_{17})^{(2)}G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1/(a'_{17})^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2/(a'_{18})^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b''_i)^{(2)}((G_{19})(t), t)) = (b'_{17})^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b''_i)^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205

unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b''_{18})^{(2)}\left((G_{19})(t),t\right)=(b'_{18})^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}$, $\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}<1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)}+\left((\hat{P}_{20})^{(3)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{20})^{(3)}\right]\leq(\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself 210

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)},(T_{23})^{(1)}),((G_{23})^{(2)},(T_{23})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}},\widetilde{T_{23}}:(\widetilde{(G_{23})},\widetilde{(T_{23})})=\mathcal{A}^{(3)}((G_{23}),(T_{23}))$

It results 213

$$\left|\tilde{G}_{20}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{20})^{(3)}\left|G_{21}^{(1)}-G_{21}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)}+$$

$$\int_0^t\{(a'_{20})^{(3)}\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)\left|G_{20}^{(1)}-G_{20}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}+$$

$$G_{20}^{(2)}\left|(a''_{20})^{(3)}\left(T_{21}^{(1)},s_{(20)}\right)-(a''_{20})^{(3)}\left(T_{21}^{(2)},s_{(20)}\right)\right|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)}-G_{23}{}^{(2)}\right|e^{-(\hat{M}_{20})^{(3)}t}\leq \quad 214$$
$$\frac{1}{(\hat{M}_{20})^{(3)}}\left((a_{20})^{(3)}+(a'_{20})^{(3)}+(\hat{A}_{20})^{(3)}+(\hat{P}_{20})^{(3)}(\hat{k}_{20})^{(3)}\right)d\left(((G_{23})^{(1)},(T_{23})^{(1)};(G_{23})^{(2)},(T_{23})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 215

$(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i(t) = 0\ and\ T_i(t) = 0$ 216

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(3)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a_{21}')^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a_{21}')^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a_{22}')^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b_{21}')^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1 - e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}$, $\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)}+\left((\widehat{P}_{24})^{(4)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{24})^{(4)} \tag{222}$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{24})^{(4)}\right]\leq(\widehat{Q}_{24})^{(4)} \tag{223}$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)},(T_{27})^{(1)}),((G_{27})^{(2)},(T_{27})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{27})},\widetilde{(T_{27})}$: $\left(\widetilde{(G_{27})},\widetilde{(T_{27})}\right)=\mathcal{A}^{(4)}((G_{27}),(T_{27}))$

It results

$$|\tilde{G}_{24}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{24})^{(4)}|G_{25}^{(1)}-G_{25}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)}+$$

$$\int_0^t\{(a'_{24})^{(4)}|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)|G_{24}^{(1)}-G_{24}^{(2)}|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}+$$

$$G_{24}^{(2)}|(a''_{24})^{(4)}\left(T_{25}^{(1)},s_{(24)}\right)-(a''_{24})^{(4)}\left(T_{25}^{(2)},s_{(24)}\right)|\;e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on Equations it follows

$$|(G_{27})^{(1)}-(G_{27})^{(2)}|e^{-(\widehat{M}_{24})^{(4)}t}\leq \tag{226}$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)}+(a'_{24})^{(4)}+(\widehat{A}_{24})^{(4)}+(\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)},(T_{27})^{(1)};(G_{27})^{(2)},(T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i=24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2 \; and \; \left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also $G_{25} \; and \; G_{26}$. indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a_{25}')^{(4)} G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_1 / (a_{25}')^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)} \left((\widehat{M}_{24})^{(4)}\right)_2 / (a_{26}')^{(4)}$$

If $G_{25} \; or \; G_{26}$ is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If $G_{24} \; is$ bounded, from below, the same property holds for $G_{25} \; and \; G_{26}$. The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b_{25}')^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b_{26}'')^{(4)}\left((G_{27})(t), t\right) = (b_{26}')^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 42

Analogous inequalities hold also for G_{29} , $G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}$, $\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)}+\left((\hat{P}_{28})^{(5)}+G_j^0\right)e^{-\left(\frac{(\hat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right]\leq(\hat{P}_{28})^{(5)} \tag{234}$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\hat{Q}_{28})^{(5)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{28})^{(5)}\right]\leq(\hat{Q}_{28})^{(5)} \tag{235}$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric (236)

$$d\left(((G_{31})^{(1)},(T_{31})^{(1)}),((G_{31})^{(2)},(T_{31})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}),(\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}),(\widetilde{T_{31}})\right)=\mathcal{A}^{(5)}\left((G_{31}),(T_{31})\right)$

It results

$$\left|\tilde{G}_{28}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{28})^{(5)}\left|G_{29}^{(1)}-G_{29}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)}+$$

$$\int_0^t\{(a'_{28})^{(5)}\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)\left|G_{28}^{(1)}-G_{28}^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}+$$

$$G_{28}^{(2)}\left|(a''_{28})^{(5)}\left(T_{29}^{(1)},s_{(28)}\right)-(a''_{28})^{(5)}\left(T_{29}^{(2)},s_{(28)}\right)\right|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$\left|(G_{31})^{(1)}-(G_{31})^{(2)}\right|e^{-(\widehat{M}_{28})^{(5)}t}\leq \tag{237}$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)}+(a'_{28})^{(5)}+(\widehat{A}_{28})^{(5)}+(\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)},(T_{31})^{(1)};(G_{31})^{(2)},(T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$. (238)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i=28,29,30$ depend only on T_{29} and respectively on (G_{31})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (239)

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(5)} - (a_i'')^{(5)}(T_{29}(s_{(28)}), s_{(28)})\} ds_{(28)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(5)} t)} > 0 \quad \text{for t} > 0$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2$ *and* $\left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also G_{29} *and* G_{30} . indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)} G_{29}$ and by integrating

$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1 / (a_{29}')^{(5)}$

In the same way , one can obtain

$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2 / (a_{30}')^{(5)}$

If G_{29} *or* G_{30} is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If G_{28} *is* bounded, from below, the same property holds for G_{29} *and* G_{30} . The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}((G_{31})(t), t)) = (b_{29}')^{(5)}$ then $T_{29} \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t > t_5$

$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t), t) < \varepsilon_5, T_{28}(t) > (m)^{(5)}$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29} \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t), t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)} + \left((\widehat{P}_{32})^{(6)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)} + G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{32})^{(6)} \quad (245)$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\hat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{32})^{(6)}\right]\leq(\hat{Q}_{32})^{(6)} \quad (246)$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$\left|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}\right|\leq\int_0^t(a_{32})^{(6)}\left|G_{33}^{(1)}-G_{33}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)\left|G_{32}^{(1)}-G_{32}^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}\left|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)\right|\ e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses it follows

$$\left|(G_{35})^{(1)}-(G_{35})^{(2)}\right|e^{-(\widehat{M}_{32})^{(6)}t}\leq \quad (248)$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ (250)

it results

$$G_i(t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(6)}-(a_i'')^{(6)}(T_{33}(s_{(32)}),s_{(32)})\}ds_{(32)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(6)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2$ *and* $\left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also G_{33} *and* G_{34} . indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_1/(a_{33}')^{(6)}$

In the same way , one can obtain

$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)}\left((\widehat{M}_{32})^{(6)}\right)_2/(a_{34}')^{(6)}$

If G_{33} *or* G_{34} is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for G_{33} *and* G_{34} . The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}\,((G_{35})(t),t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t),t\right) < \varepsilon_6, T_{32}\,(t) > (m)^{(6)}$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t),t(t),t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

$$\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{36})^{(7)}\right] \leq (\hat{Q}_{36})^{(7)} \quad 257$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{39})},\widetilde{(T_{39})}:\left(\widetilde{(G_{39})},\widetilde{(T_{39})}\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$\left|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{36})^{(7)}\left|G_{37}^{(1)}-G_{37}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\,ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)\left|G_{36}^{(1)}-G_{36}^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}\left|(a''_{36})^{(7)}\left(T_{37}^{(1)},s_{(36)}\right)-(a''_{36})^{(7)}\left(T_{37}^{(2)},s_{(36)}\right)\right|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on it follows

$$\left|(G_{39})^{(1)}-(G_{39})^{(2)}\right|e^{-(\widehat{M}_{36})^{(7)}t}\le \quad 259$$
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ *and* $(\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on (G_{39})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 261

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{\left(-(b_i')^{(7)}t\right)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1,\left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a'_{37})^{(7)} G_{37}$ and by integrating

$$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a'_{37})^{(7)}$$

In the same way , one can obtain

$$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a'_{38})^{(7)}$$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b'_{37})^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}(t) > (m)^{(7)}$$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b'_{38})^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\widehat{P}_{40})^{(8)}$ and $(\widehat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[(\widehat{P}_{40})^{(8)} + \left((\widehat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\widehat{M}_{40})^{(8)}}\left[\left((\widehat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{40})^{(8)}\right] \leq (\widehat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right) = \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}|\le\int_0^t(a_{40})^{(8)}|G_{41}^{(1)}-G_{41}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})|G_{40}^{(1)}-G_{40}^{(2)}|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}|(a''_{40})^{(8)}(T_{41}^{(1)},s_{(40)})-(a''_{40})^{(8)}(T_{41}^{(2)},s_{(40)})|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,t]$ 272

From the hypotheses it follows

$$|(G_{43})^{(1)}-(G_{43})^{(2)}|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ *and* $(\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40,41,42$ depend only on T_{41} and respectively on (G_{43})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i(t)=0$ *and* $T_i(t)=0$ 275

it results

$$G_i(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i(t)\ge T_i^0e^{(-(b_i')^{(8)}t)}>0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2$ *and* $\left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also G_{41} *and* G_{42} . indeed if

$G_{40}<(\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt}\le\left((\widehat{M}_{40})^{(8)}\right)_1-(a'_{41})^{(8)}G_{41}$ and by integrating

$$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$$

In the same way , one can obtain

$$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$$

If G_{41} *or* G_{42} is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If G_{40} *is* bounded, from below, the same property holds for G_{41} *and* G_{42} . The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right)$, $t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

279 A

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}$, $\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right) =$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{44})^{(9)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $\widetilde{(G_{47})}, \widetilde{(T_{47})} : \left(\widetilde{(G_{47})}, \widetilde{(T_{47})} \right) = \mathcal{A}^{(9)}\left((G_{47}), (T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)} - \tilde{G}_{i}^{(2)}\right| \leq \int_0^t (a_{44})^{(9)} \left|G_{45}^{(1)} - G_{45}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\, ds_{(44)} +$$
$$\int_0^t \{(a'_{44})^{(9)} \left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right)\left|G_{44}^{(1)} - G_{44}^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}} +$$
$$G_{44}^{(2)} \left|(a''_{44})^{(9)}\left(T_{45}^{(1)}, s_{(44)}\right) - (a''_{44})^{(9)}\left(T_{45}^{(2)}, s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)} s_{(44)}} e^{(\widehat{M}_{44})^{(9)} s_{(44)}}\} ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)} - G^{(2)}\right| e^{-(\widehat{M}_{44})^{(9)} t} \leq$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)} + (a'_{44})^{(9)} + (\widehat{A}_{44})^{(9)} + (\widehat{P}_{44})^{(9)} (\widehat{k}_{44})^{(9)}\right) d\left(\left((G_{47})^{(1)}, (T_{47})^{(1)};\ (G_{47})^{(2)}, (T_{47})^{(2)}\right)\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ *and* $(\widehat{Q}_{44})^{(9)} e^{(\widehat{M}_{44})^{(9)} t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i = 44{,}45{,}46$ depend only on T_{45} and respectively on (G_{47})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$

From 99 to 44 it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(9)} - (a_i'')^{(9)}\left(T_{45}(s_{(44)}), s_{(44)}\right)\} ds_{(44)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{\left(-(b_i')^{(9)} t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1, \left((\widehat{M}_{44})^{(9)}\right)_2$ *and* $\left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also G_{45} *and* G_{46} . indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1 / (a'_{45})^{(9)}$$

In the same way , one can obtain

$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14}, t) + (a''_{14})^{(1)}(T_{14}, t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (v_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)} t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)} t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)} t}\right] + G_{15}^0 e^{-(S_2)^{(1)} t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)} t} - e^{-(a'_{15})^{(1)} t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)} t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)} t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)} t} - e^{-(b'_{15})^{(1)} t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)} t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)} t}\right] + T_{15}^0 e^{-(R_2)^{(1)} t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$

$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$

$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, $\boldsymbol{if}\ (\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}, \boldsymbol{if}\ (\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, $\boldsymbol{if}\ (\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, $\boldsymbol{if}\ (u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}, \boldsymbol{if}\ (u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)}$, $\boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$\mathrm{G}^0_{16} e^{((\mathrm{S}_1)^{(2)} - (p_{16})^{(2)})\mathrm{t}} \le G_{16}(t) \le \mathrm{G}^0_{16} e^{(\mathrm{S}_1)^{(2)}\mathrm{t}}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}}\mathrm{G}_{16}^{0}\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}} \quad 311$$

$$\left(\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_1)^{(2)}((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)}-(\mathrm{S}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{S}_1)^{(2)}-(p_{16})^{(2)})\mathrm{t}}-\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(\mathrm{S}_2)^{(2)}\mathrm{t}} \le \mathrm{G}_{18}(t) \le \right. \quad 312$$

$$\left.\frac{(a_{18})^{(2)}\mathrm{G}_{16}^{0}}{(m_2)^{(2)}((\mathrm{S}_1)^{(2)}-(a'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{S}_1)^{(2)}\mathrm{t}}-\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right]+\mathrm{G}_{18}^{0}\mathrm{e}^{-(a'_{18})^{(2)}\mathrm{t}}\right)$$

$$\boxed{\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{(\mathrm{R}_1)^{(2)}t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}}\mathrm{T}_{16}^{0}\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_1)^{(2)}((\mathrm{R}_1)^{(2)}-(b'_{18})^{(2)})}\left[\mathrm{e}^{(\mathrm{R}_1)^{(2)}t}-\mathrm{e}^{-(b'_{18})^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(b'_{18})^{(2)}t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)}\mathrm{T}_{16}^{0}}{(\mu_2)^{(2)}((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)}+(\mathrm{R}_2)^{(2)})}\left[\mathrm{e}^{((\mathrm{R}_1)^{(2)}+(r_{16})^{(2)})t}-\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}\right]+\mathrm{T}_{18}^{0}\mathrm{e}^{-(\mathrm{R}_2)^{(2)}t}$$

Definition of $(\mathrm{S}_1)^{(2)}, (\mathrm{S}_2)^{(2)}, (\mathrm{R}_1)^{(2)}, (\mathrm{R}_2)^{(2)}$:- 316

Where $(\mathrm{S}_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(\mathrm{S}_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(\mathrm{R}_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}, (m_2)^{(3)}, (\mu_1)^{(3)}, (\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (\nu_0)^{(3)}, (m_1)^{(3)} = (\nu_1)^{(3)}$, ***if*** $(\nu_0)^{(3)} < (\nu_1)^{(3)}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\bar{\nu}_1)^{(3)}$, ***if*** $(\nu_1)^{(3)} < (\nu_0)^{(3)} < (\bar{\nu}_1)^{(3)}$,

and $\boxed{(\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (\nu_1)^{(3)}, (m_1)^{(3)} = (\nu_0)^{(3)}$, ***if*** $(\bar{\nu}_1)^{(3)} < (\nu_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, ***if*** $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, ***if*** $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, ***if*** $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{20}(t) \leq G_{20}^0 e^{(S_1)^{(3)} t}$$

$(p_i)^{(3)}$ is defined by equation

$$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \leq G_{21}(t) \leq \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)} t} \quad 323$$

$$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)} t}\right] + G_{22}^0 e^{-(S_2)^{(3)} t} \leq G_{22}(t) \leq \right. \quad 324$$

$$\left. \frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)} t} - e^{-(a'_{22})^{(3)} t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)} t}\right)$$

$$\boxed{T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}} \quad 325$$

$$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)} t} \leq T_{20}(t) \leq \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t} \quad 326$$

$$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)} t} - e^{-(b'_{22})^{(3)} t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)} t} \leq T_{22}(t) \leq \quad 327$$

$$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)} t}\right] + T_{22}^0 e^{-(R_2)^{(3)} t}$$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$

$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$

$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously

331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)} \text{ where } (u_1)^{(4)}, (\bar{u}_1)^{(4)}$$

Then the solution of global equations satisfies the inequalities

332

$$G^0_{24} e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G^0_{24} e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)}-(p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)}t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)}-(p_{24})^{(4)}-(S_2)^{(4)})}\left[e^{((S_1)^{(4)}-(p_{24})^{(4)})t} - e^{-(S_2)^{(4)}t}\right] + G_{26}^0 e^{-(S_2)^{(4)}t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)}-(a'_{26})^{(4)})}\left[e^{(S_1)^{(4)}t} - e^{-(a'_{26})^{(4)}t}\right] + G_{26}^0 e^{-(a'_{26})^{(4)}t}\right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)}t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)}+(r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)}-(b'_{26})^{(4)})}\left[e^{(R_1)^{(4)}t} - e^{-(b'_{26})^{(4)}t}\right] + T_{26}^0 e^{-(b'_{26})^{(4)}t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)}+(r_{24})^{(4)}+(R_2)^{(4)})}\left[e^{((R_1)^{(4)}+(r_{24})^{(4)})t} - e^{-(R_2)^{(4)}t}\right] + T_{26}^0 e^{-(R_2)^{(4)}t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}\big((G_{31}), t\big) - (b''_{29})^{(5)}\big((G_{31}), t\big) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\big(v^{(5)}\big)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\big(u^{(5)}\big)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{\nu}_1)^{(5)},,(\bar{\nu}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{\nu}_1)^{(5)} > 0$, $(\bar{\nu}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)} = 0$ and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)},(\nu_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)},(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (\nu_0)^{(5)}, (m_1)^{(5)} = (\nu_1)^{(5)},$ ***if*** $(\nu_0)^{(5)} < (\nu_1)^{(5)}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\bar{\nu}_1)^{(5)}$, ***if*** $(\nu_1)^{(5)} < (\nu_0)^{(5)} < (\bar{\nu}_1)^{(5)},$

and $\boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (\nu_1)^{(5)}, (m_1)^{(5)} = (\nu_0)^{(5)},$ ***if*** $(\bar{\nu}_1)^{(5)} < (\nu_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)},$ ***if*** $(u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, ***if*** $(u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)},$

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)},$ ***if*** $(\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}\left((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)}\right)}\left[e^{\left((S_1)^{(5)} - (p_{28})^{(5)}\right)t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}\left((S_1)^{(5)} - (a'_{30})^{(5)}\right)}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{\left((R_1)^{(5)} + (r_{28})^{(5)}\right)t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}} T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}} T_{28}^0 e^{\left((R_1)^{(5)} + (r_{28})^{(5)}\right)t} \quad 346$$

$$\frac{(b_{30})^{(5)} T_{28}^0}{(\mu_1)^{(5)}\left((R_1)^{(5)} - (b'_{30})^{(5)}\right)}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^{0}}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t}-e^{-(R_2)^{(5)}t}\right]+T_{30}^{0}e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)}, , (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\left(v^{(6)}\right)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\left(u^{(6)}\right)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, ***if*** $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, ***if*** $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,
and $\boxed{(v_0)^{(6)} = \frac{G_{32}^{0}}{G_{33}^{0}}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)}$, ***if*** $(\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)}, \ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)} \text{ where } (u_1)^{(6)}, (\bar{u}_1)^{(6)}$$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$$

$$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$$

$$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(v_1)^{(7)}, (v_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, v^{(7)}, u^{(7)}$: 361

By $(v_1)^{(7)} > 0$, $(v_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0$, $(u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{v}_1)^{(7)}, , (\bar{v}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{v}_1)^{(7)} > 0$, $(\bar{v}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0$, $(\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(v^{(7)}\big)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}$, $(m_2)^{(7)}$, $(\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}$, $(m_2)^{(7)}$, $(\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$
and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$
and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)}$$ where $(u_1)^{(7)}, (\bar{u}_1)^{(7)}$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G^0_{36} e^{((S_1)^{(7)} - (p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G^0_{36} e^{(S_1)^{(7)}t}$$ 365

$$\left(\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t}-e^{-(S_2)^{(7)}t}\right]+G_{38}^{0}e^{-(S_2)^{(7)}t}\leq G_{38}(t)\leq\right. \quad 366$$

$$\left.\frac{(a_{38})^{(7)}G_{36}^{0}}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t}-e^{-(a'_{38})^{(7)}t}\right]+G_{38}^{0}e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}}T_{36}^{0}e^{(R_1)^{(7)}t}\leq T_{36}(t)\leq\frac{1}{(\mu_2)^{(7)}}T_{36}^{0}e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)}T_{36}^{0}}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t}-e^{-(b'_{38})^{(7)}t}\right]+T_{38}^{0}e^{-(b'_{38})^{(7)}t}\leq T_{38}(t)\leq \quad 369$$

$$\frac{(a_{38})^{(7)}T_{36}^{0}}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t}-e^{-(R_2)^{(7)}t}\right]+T_{38}^{0}e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)},(S_2)^{(7)},(R_1)^{(7)},(R_2)^{(7)}$:- 370

Where $(S_1)^{(7)}=(a_{36})^{(7)}(m_2)^{(7)}-(a'_{36})^{(7)}$

$(S_2)^{(7)}=(a_{38})^{(7)}-(p_{38})^{(7)}$

$(R_1)^{(7)}=(b_{36})^{(7)}(\mu_2)^{(7)}-(b'_{36})^{(7)}$

$(R_2)^{(7)}=(b'_{38})^{(7)}-(r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$:

$(\sigma_1)^{(8)},(\sigma_2)^{(8)},(\tau_1)^{(8)},(\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)}\leq-(a'_{40})^{(8)}+(a'_{41})^{(8)}-(a''_{40})^{(8)}(T_{41},t)+(a''_{41})^{(8)}(T_{41},t)\leq-(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)}\leq-(b'_{40})^{(8)}+(b'_{41})^{(8)}-(b''_{40})^{(8)}\big((G_{43}),t\big)-(b''_{41})^{(8)}\big((G_{43}),t\big)\leq-(\tau_1)^{(8)}$$

Definition of $(v_1)^{(8)},(v_2)^{(8)},(u_1)^{(8)},(u_2)^{(8)},v^{(8)},u^{(8)}$: 372

By $(v_1)^{(8)}>0$, $(v_2)^{(8)}<0$ and respectively $(u_1)^{(8)}>0$, $(u_2)^{(8)}<0$ the roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_1)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2+(\tau_1)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$ and

Definition of $(\bar{v}_1)^{(8)},,(\bar{v}_2)^{(8)},(\bar{u}_1)^{(8)},(\bar{u}_2)^{(8)}$:

By $(\bar{v}_1)^{(8)}>0$, $(\bar{v}_2)^{(8)}<0$ and respectively $(\bar{u}_1)^{(8)}>0$, $(\bar{u}_2)^{(8)}<0$ the

roots of the equations $(a_{41})^{(8)}\big(v^{(8)}\big)^2+(\sigma_2)^{(8)}v^{(8)}-(a_{40})^{(8)}=0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2+(\tau_2)^{(8)}u^{(8)}-(b_{40})^{(8)}=0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (\nu_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)}=(\nu_0)^{(8)}, (m_1)^{(8)}=(\nu_1)^{(8)},\ \boldsymbol{if}\ (\nu_0)^{(8)}<(\nu_1)^{(8)}$

$(m_2)^{(8)}=(\nu_1)^{(8)}, (m_1)^{(8)}=(\bar{\nu}_1)^{(8)}, \boldsymbol{if}\ (\nu_1)^{(8)}<(\nu_0)^{(8)}<(\bar{\nu}_1)^{(8)},$
and $\boxed{(\nu_0)^{(8)}=\frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)}=(\nu_1)^{(8)}, (m_1)^{(8)}=(\nu_0)^{(8)},\ \boldsymbol{if}\ (\bar{\nu}_1)^{(8)}<(\nu_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)}=(u_0)^{(8)}, (\mu_1)^{(8)}=(u_1)^{(8)},\ \boldsymbol{if}\ (u_0)^{(8)}<(u_1)^{(8)}$

$(\mu_2)^{(8)}=(u_1)^{(8)}, (\mu_1)^{(8)}=(\bar{u}_1)^{(8)}, \boldsymbol{if}\ (u_1)^{(8)}<(u_0)^{(8)}<(\bar{u}_1)^{(8)},$
and $\boxed{(u_0)^{(8)}=\frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)}=(u_1)^{(8)}, (\mu_1)^{(8)}=(u_0)^{(8)}, \boldsymbol{if}\ (\bar{u}_1)^{(8)}<(u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{\left((S_1)^{(8)}-(p_{40})^{(8)}\right)t}\le G_{40}(t)\le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}}G_{40}^0 e^{\left((S_1)^{(8)}-(p_{40})^{(8)}\right)t}\le G_{41}(t)\le\frac{1}{(m_2)^{(8)}}G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)}G_{40}^0}{(m_1)^{(8)}\left((S_1)^{(8)}-(p_{40})^{(8)}-(S_2)^{(8)}\right)}\left[e^{\left((S_1)^{(8)}-(p_{40})^{(8)}\right)t}-e^{-(S_2)^{(8)}t}\right]+G_{42}^0 e^{-(S_2)^{(8)}t}\le G_{42}(t)\le\right. \quad 377$$

$$\left.\frac{(a_{42})^{(8)}G_{40}^0}{(m_2)^{(8)}\left((S_1)^{(8)}-(a'_{42})^{(8)}\right)}\left[e^{(S_1)^{(8)}t}-e^{-(a'_{42})^{(8)}t}\right]+G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t}\le T_{40}(t)\le T_{40}^0 e^{\left((R_1)^{(8)}+(r_{40})^{(8)}\right)t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}}T_{40}^0 e^{(R_1)^{(8)}t}\le T_{40}(t)\le\frac{1}{(\mu_2)^{(8)}}T_{40}^0 e^{\left((R_1)^{(8)}+(r_{40})^{(8)}\right)t} \quad 379$$

$$\frac{(b_{42})^{(8)}T_{40}^0}{(\mu_1)^{(8)}\left((R_1)^{(8)}-(b'_{42})^{(8)}\right)}\left[e^{(R_1)^{(8)}t}-e^{-(b'_{42})^{(8)}t}\right]+T_{42}^0 e^{-(b'_{42})^{(8)}t}\le T_{42}(t)\le \quad 380$$

$$\frac{(a_{42})^{(8)}T_{40}^0}{(\mu_2)^{(8)}\left((R_1)^{(8)}+(r_{40})^{(8)}+(R_2)^{(8)}\right)}\left[e^{\left((R_1)^{(8)}+(r_{40})^{(8)}\right)t}-e^{-(R_2)^{(8)}t}\right]+T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$

$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$

$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$

$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}$, $\boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}$, $\boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)}$,
and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$

and analogously

$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}$, $\boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}\ (u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,
and $\boxed{(u_0)^{(9)} = \frac{T^0_{44}}{T^0_{45}}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}\ (\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{d\nu^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)\nu^{(1)} - (a_{14})^{(1)}\nu^{(1)}$$

Definition of $\nu^{(1)}$:- $\boxed{\nu^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}(\nu^{(1)})^2 + (\sigma_2)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right) \leq \frac{d\nu^{(1)}}{dt} \leq -\left((a_{14})^{(1)}(\nu^{(1)})^2 + (\sigma_1)^{(1)}\nu^{(1)} - (a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)}, (\nu_0)^{(1)}$:-

For $0 < \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}} < (\nu_1)^{(1)} < (\bar{\nu}_1)^{(1)}$

$$\nu^{(1)}(t) \geq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_0)^{(1)}\right)t\right]}} \quad , \quad \boxed{(C)^{(1)} = \frac{(\nu_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\nu_2)^{(1)}}}$$

it follows $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\nu_1)^{(1)}$

In the same manner , we get 384

$$\nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(1)} = \frac{(\bar{\nu}_1)^{(1)} - (\nu_0)^{(1)}}{(\nu_0)^{(1)} - (\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)} \leq \nu^{(1)}(t) \leq (\bar{\nu}_1)^{(1)}$

If $0 < (\nu_1)^{(1)} < (\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0} < (\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)} \leq \frac{(\nu_1)^{(1)} + (C)^{(1)}(\nu_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}}{1 + (C)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)} - (\nu_2)^{(1)}\right)t\right]}} \leq \nu^{(1)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\bar{\nu}_1)^{(1)}$$

If $0 < (\nu_1)^{(1)} \leq (\bar{\nu}_1)^{(1)} \leq \boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)} \leq \nu^{(1)}(t) \leq \frac{(\bar{\nu}_1)^{(1)} + (\bar{C})^{(1)}(\bar{\nu}_2)^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}}{1 + (\bar{C})^{(1)} e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)} - (\bar{\nu}_2)^{(1)}\right)t\right]}} \leq (\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(1)}(t)$:-

$$(m_2)^{(1)} \leq \nu^{(1)}(t) \leq (m_1)^{(1)}, \quad \boxed{\nu^{(1)}(t) = \frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)} \leq u^{(1)}(t) \leq (\mu_1)^{(1)}, \quad \boxed{u^{(1)}(t) = \frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also

defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)}T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \le \nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq v^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{v^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $v^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)} G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)} T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{dv^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)v^{(3)} - (a_{21})^{(3)}v^{(3)}$$

Definition of $v^{(3)}$:- $\boxed{v^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)}\right) \leq \frac{dv^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_1)^{(3)}v^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$v^{(3)}(t) \geq \frac{(\nu_1)^{(3)} + (C)^{(3)}(\nu_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}}{1 + (C)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)} - (\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq v^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$v^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)} + (\bar{C})^{(3)}(\bar{\nu}_2)^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}}{1 + (\bar{C})^{(3)} e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)} - (\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)} - (\nu_0)^{(3)}}{(\nu_0)^{(3)} - (\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{20}'')^{(3)} = (a_{21}'')^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b_{20}'')^{(3)} = (b_{21}'')^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a_{24}')^{(4)} - (a_{25}')^{(4)} + (a_{24}'')^{(4)}(T_{25}, t)\right) - (a_{25}'')^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\quad \boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_2)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)\le\frac{dv^{(4)}}{dt}\le-\left((a_{25})^{(4)}\left(v^{(4)}\right)^2+(\sigma_4)^{(4)}v^{(4)}-(a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(4)},(v_0)^{(4)}$:-

For $0<\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}<(v_1)^{(4)}<(\bar{v}_1)^{(4)}$

$$v^{(4)}(t)\ge\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}{4+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_0)^{(4)}\right)t\right]}}\quad,\quad\boxed{(C)^{(4)}=\frac{(v_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(v_2)^{(4)}}}$$

it follows $(v_0)^{(4)}\le v^{(4)}(t)\le(v_1)^{(4)}$

In the same manner , we get 405

$$v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{4+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(4)}=\frac{(\bar{v}_1)^{(4)}-(v_0)^{(4)}}{(v_0)^{(4)}-(\bar{v}_2)^{(4)}}}$$

From which we deduce $(v_0)^{(4)}\le v^{(4)}(t)\le(\bar{v}_1)^{(4)}$

If $0<(v_1)^{(4)}<(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}<(\bar{v}_1)^{(4)}$ we find like in the previous case, 406

$$(v_1)^{(4)}\le\frac{(v_1)^{(4)}+(C)^{(4)}(v_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}{1+(C)^{(4)}e^{\left[-(a_{25})^{(4)}\left((v_1)^{(4)}-(v_2)^{(4)}\right)t\right]}}\le v^{(4)}(t)\le$$

$$\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(\bar{v}_1)^{(4)}$$

If $0<(v_1)^{(4)}\le(\bar{v}_1)^{(4)}\le\boxed{(v_0)^{(4)}=\frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(v_1)^{(4)}\le v^{(4)}(t)\le\frac{(\bar{v}_1)^{(4)}+(\bar{C})^{(4)}(\bar{v}_2)^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}{1+(\bar{C})^{(4)}e^{\left[-(a_{25})^{(4)}\left((\bar{v}_1)^{(4)}-(\bar{v}_2)^{(4)}\right)t\right]}}\le(v_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(4)}(t)$:-

$$(m_2)^{(4)}\le v^{(4)}(t)\le(m_1)^{(4)},\quad\boxed{v^{(4)}(t)=\frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)}\le u^{(4)}(t)\le(\mu_1)^{(4)},\quad\boxed{u^{(4)}(t)=\frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \leq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}}{1 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_2)^{(5)}\right)t\right]}} \leq \nu^{(5)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \leq (\bar{\nu}_1)^{(5)} \leq \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain

$$(\nu_1)^{(5)} \leq \nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{1 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \leq (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$$(m_2)^{(5)} \leq \nu^{(5)}(t) \leq (m_1)^{(5)}, \quad \boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$$(\mu_2)^{(5)} \leq u^{(5)}(t) \leq (\mu_1)^{(5)}, \quad \boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)} G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$ if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)} T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\quad \boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \leq \frac{d\nu^{(6)}}{dt} \leq -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then $(u_1)^{(6)} = (\bar{u}_1)^{(6)}$ if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(\nu_1)^{(6)}$ and $(\bar{\nu}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right) - (a''_{37})^{(7)}(T_{37},t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \le \frac{dv^{(7)}}{dt} \le -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(7)}, (\nu_0)^{(7)}$:-

For $0 < \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (\nu_1)^{(7)} < (\bar{\nu}_1)^{(7)}$

$$v^{(7)}(t) \ge \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(\nu_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\nu_2)^{(7)}}}$$

it follows $(\nu_0)^{(7)} \le v^{(7)}(t) \le (\nu_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{\nu}_1)^{(7)} - (\nu_0)^{(7)}}{(\nu_0)^{(7)} - (\bar{\nu}_2)^{(7)}}}$$

From which we deduce $(\nu_0)^{(7)} \le v^{(7)}(t) \le (\bar{\nu}_1)^{(7)}$

If $0 < (\nu_1)^{(7)} < (\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{\nu}_1)^{(7)}$ we find like in the previous case, 418

$$(\nu_1)^{(7)} \le \frac{(\nu_1)^{(7)} + (C)^{(7)}(\nu_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\nu_1)^{(7)} - (\nu_2)^{(7)}\right)t\right]}} \le v^{(7)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\bar{\nu}_1)^{(7)}$$

If $0 < (\nu_1)^{(7)} \le (\bar{\nu}_1)^{(7)} \le \boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(\nu_1)^{(7)} \le v^{(7)}(t) \le \frac{(\bar{\nu}_1)^{(7)} + (\bar{C})^{(7)}(\bar{\nu}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{\nu}_1)^{(7)} - (\bar{\nu}_2)^{(7)}\right)t\right]}} \le (\nu_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$$(m_2)^{(7)} \le v^{(7)}(t) \le (m_1)^{(7)}, \quad \boxed{v^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}, \quad \boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\quad \boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

$$\text{For } 0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain

424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{40}'')^{(8)} = (a_{41}'')^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b_{40}'')^{(8)} = (b_{41}'')^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424
A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a_{44}')^{(9)} - (a_{45}')^{(9)} + (a_{44}'')^{(9)}(T_{45}, t)\right) - (a_{45}'')^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_2)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right) \leq \frac{d\nu^{(9)}}{dt} \leq -\left((a_{45})^{(9)}\left(\nu^{(9)}\right)^2 + (\sigma_1)^{(9)}\nu^{(9)} - (a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)}, (\nu_0)^{(9)}$:-

For $0 < \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}} < (\nu_1)^{(9)} < (\bar{\nu}_1)^{(9)}$

$$\nu^{(9)}(t) \geq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}} \quad , \quad \boxed{(C)^{(9)} = \frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\nu_9)^{(9)}$

In the same manner , we get

$$\nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(9)} = \frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)} \leq \nu^{(9)}(t) \leq (\bar{\nu}_1)^{(9)}$

If $0 < (\nu_1)^{(9)} < (\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0} < (\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)} \leq \frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}} \leq \nu^{(9)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\bar{\nu}_1)^{(9)}$$

If $0 < (\nu_1)^{(9)} \leq (\bar{\nu}_1)^{(9)} \leq \boxed{(\nu_0)^{(9)} = \frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)} \leq \nu^{(9)}(t) \leq \frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}} \leq (\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(9)}(t)$:-

$$(m_2)^{(9)} \leq \nu^{(9)}(t) \leq (m_1)^{(9)}, \quad \boxed{\nu^{(9)}(t) = \frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)} \leq u^{(9)}(t) \leq (\mu_1)^{(9)}, \quad \boxed{u^{(9)}(t) = \frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a_{44}'')^{(9)} = (a_{45}'')^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)}G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then
$(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 436 A
45,46,27,28)

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\,]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\,]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\,]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\,]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\,]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\,]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\,]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\,]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\,]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(o) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$$

Definition and uniqueness of T^*_{14} :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T^*_{14} for which $f(T^*_{14}) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}\mathrm{G}_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(\mathrm{T}^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}\mathrm{G}_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(\mathrm{T}^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T^*_{41} :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T^*_{41} for which $f(T^*_{41}) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T^*_{41})]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T^*_{41})]}$$

Definition and uniqueness of T^*_{45} :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T^*_{45} for which $f(T^*_{45}) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T^*_{45})]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T^*_{45})]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502
$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see 504
that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{14} such that $\varphi((G_{19})^*) = 0$

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{21} such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right]+(b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right]+(b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right]+(b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right]+(b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right]+(b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0\,, T^*_{14}$ given by $f(T^*_{14}) = 0$ and

$$G^*_{13} = \frac{(a_{13})^{(1)}G^*_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G^*_{15} = \frac{(a_{15})^{(1)}G^*_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

$$T^*_{13} = \frac{(b_{13})^{(1)}T^*_{14}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^*)]} \quad , \quad T^*_{15} = \frac{(b_{15})^{(1)}T^*_{14}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^*)]}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^* given by $\varphi((G_{19})^*) = 0$, T_{17}^* given by $f(T_{17}^*) = 0$ and 512

$$G_{16}^* = \frac{(a_{16})^{(2)} G_{17}^*}{[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17}^*)]} \quad , \quad G_{18}^* = \frac{(a_{18})^{(2)} G_{17}^*}{[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}^*)]} \tag{513}$$

$$T_{16}^* = \frac{(b_{16})^{(2)} T_{17}^*}{[(b'_{16})^{(2)} - (b''_{16})^{(2)}((G_{19})^*)]} \quad , \quad T_{18}^* = \frac{(b_{18})^{(2)} T_{17}^*}{[(b'_{18})^{(2)} - (b''_{18})^{(2)}((G_{19})^*)]} \tag{514}$$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^* given by $\varphi((G_{23})^*) = 0$, T_{21}^* given by $f(T_{21}^*) = 0$ and

$$G_{20}^* = \frac{(a_{20})^{(3)} G_{21}^*}{[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21}^*)]} \quad , \quad G_{22}^* = \frac{(a_{22})^{(3)} G_{21}^*}{[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}^*)]}$$

$$T_{20}^* = \frac{(b_{20})^{(3)} T_{21}^*}{[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}{}^*)]} \quad , \quad T_{22}^* = \frac{(b_{22})^{(3)} T_{21}^*}{[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}{}^*)]}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^* given by $\varphi(G_{27}) = 0$, T_{25}^* given by $f(T_{25}^*) = 0$ and

$$G_{24}^* = \frac{(a_{24})^{(4)} G_{25}^*}{[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25}^*)]} \quad , \quad G_{26}^* = \frac{(a_{26})^{(4)} G_{25}^*}{[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}^*)]}$$

$$T_{24}^* = \frac{(b_{24})^{(4)} T_{25}^*}{[(b'_{24})^{(4)} - (b''_{24})^{(4)}((G_{27})^*)]} \quad , \quad T_{26}^* = \frac{(b_{26})^{(4)} T_{25}^*}{[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27})^*)]} \tag{517}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^* given by $\varphi((G_{31})^*) = 0$, T_{29}^* given by $f(T_{29}^*) = 0$ and

$$G_{28}^* = \frac{(a_{28})^{(5)} G_{29}^*}{[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29}^*)]} \quad , \quad G_{30}^* = \frac{(a_{30})^{(5)} G_{29}^*}{[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}^*)]}$$

$$T_{28}^* = \frac{(b_{28})^{(5)} T_{29}^*}{[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31})^*)]} \quad , \quad T_{30}^* = \frac{(b_{30})^{(5)} T_{29}^*}{[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31})^*)]} \tag{519}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^* given by $\varphi((G_{35})^*) = 0$, T_{33}^* given by $f(T_{33}^*) = 0$ and

$$G_{32}^* = \frac{(a_{32})^{(6)} G_{33}^*}{[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33}^*)]} \quad , \quad G_{34}^* = \frac{(a_{34})^{(6)} G_{33}^*}{[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}^*)]}$$

$$T_{32}^* = \frac{(b_{32})^{(6)} T_{33}^*}{[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35})^*)]} \quad , \quad T_{34}^* = \frac{(b_{34})^{(6)} T_{33}^*}{[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35})^*)]} \tag{521}$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^* given by $\varphi((G_{39})^*) = 0$, T_{37}^* given by $f(T_{37}^*) = 0$ and

$$G_{36}^* = \frac{(a_{36})^{(7)} G_{37}^*}{[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37}^*)]} \quad , \quad G_{38}^* = \frac{(a_{38})^{(7)} G_{37}^*}{[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}^*)]}$$

$$T_{36}^* = \frac{(b_{36})^{(7)} T_{37}^*}{[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39})^*)]} \quad , \quad T_{38}^* = \frac{(b_{38})^{(7)} T_{37}^*}{[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39})^*)]}$$

Finally we obtain the unique solution 523

G_{41}^* given by $\varphi((G_{43})^*) = 0$, T_{41}^* given by $f(T_{41}^*) = 0$ and

$$G_{40}^* = \frac{(a_{40})^{(8)} G_{41}^*}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42}^* = \frac{(a_{42})^{(8)} G_{41}^*}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

$$T_{40}^* = \frac{(b_{40})^{(8)} T_{41}^*}{[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43})^*)]} \quad , \quad T_{42}^* = \frac{(b_{42})^{(8)} T_{41}^*}{[(b'_{42})^{(8)} - (b''_{42})^{(8)}((G_{43})^*)]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^* given by $\varphi((G_{47})^*) = 0$, T_{45}^* given by $f(T_{45}^*) = 0$ and

$$G_{44}^* = \frac{(a_{44})^{(9)} G_{45}^*}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46}^* = \frac{(a_{46})^{(9)} G_{45}^*}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

$$T_{44}^* = \frac{(b_{44})^{(9)} T_{45}^*}{[(b'_{44})^{(9)} - (b''_{44})^{(9)}((G_{47})^*)]} \quad , \quad T_{46}^* = \frac{(b_{46})^{(9)} T_{45}^*}{[(b'_{46})^{(9)} - (b''_{46})^{(9)}((G_{47})^*)]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^*) = (q_{14})^{(1)} \quad , \frac{\partial (b_i'')^{(1)}}{\partial G_j}(G^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\big((a'_{13})^{(1)} + (p_{13})^{(1)}\big)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)} G_{13}^* \mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\big((a'_{14})^{(1)} + (p_{14})^{(1)}\big)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)} G_{14}^* \mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\big((a'_{15})^{(1)} + (p_{15})^{(1)}\big)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)} G_{15}^* \mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}\left((G_{19})^*\right) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{21})^{(3)}}{\partial T_{21}}(T^*_{21}) = (q_{21})^{(3)} \quad , \frac{\partial(b''_i)^{(3)}}{\partial G_j}\left((G_{23})^*\right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a'_{20})^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G^*_{20}\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a'_{21})^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G^*_{21}\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a'_{22})^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G^*_{22}\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b'_{20})^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T^*_{20}\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b'_{21})^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T^*_{21}\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b'_{22})^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T^*_{22}\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}\ and\ (b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{25})^{(4)}}{\partial T_{25}}(T^*_{25}) = (q_{25})^{(4)} \ , \ \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a'_{24})^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G^*_{24}\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a'_{25})^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G^*_{25}\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a'_{26})^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G^*_{26}\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b'_{24})^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T^*_{24}\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b'_{25})^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T^*_{25}\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b'_{26})^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T^*_{26}\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}\ and\ (b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{29})^{(5)}}{\partial T_{29}}(T^*_{29}) = (q_{29})^{(5)} \ , \ \frac{\partial(b_i'')^{(5)}}{\partial G_j}(\ (G_{31})^*\ ) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a'_{28})^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G^*_{28}\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a'_{29})^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G^*_{29}\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a'_{30})^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G^*_{30}\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b'_{28})^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T^*_{28}\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b'_{29})^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T^*_{29}\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b'_{30})^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T^*_{30}\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(6)}$ *and* $(b''_i)^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a''_{33})^{(6)}}{\partial T_{33}}(T^*_{33}) = (q_{33})^{(6)} \quad , \frac{\partial(b''_i)^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a'_{32})^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G^*_{32}\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a'_{33})^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G^*_{33}\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a'_{34})^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G^*_{34}\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b'_{32})^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T^*_{32}\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b'_{33})^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T^*_{33}\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b'_{34})^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T^*_{34}\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(7)}$ *and* $(b''_i)^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G^*_i + \mathbb{G}_i \qquad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^{*}) = (q_{37})^{(7)} \quad , \frac{\partial(b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^{*}\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^{*}\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^{*}\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^{*}\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^{*}\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^{*}\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ and $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^{*} + \mathbb{G}_i \quad , T_i = T_i^{*} + \mathbb{T}_i$$

$$\frac{\partial(a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^{*}) = (q_{41})^{(8)} \quad , \frac{\partial(b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^{*}\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^{*}\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^{*}\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b_{40}')^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T_{40}^{*}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b_{41}')^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T_{41}^{*}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b_{42}')^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T_{42}^{*}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{45}'')^{(9)}}{\partial T_{45}}(T_{45}^*) = (q_{45})^{(9)} \quad , \frac{\partial (b_i'')^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a_{44}')^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G_{44}^*\mathbb{T}_{45} \quad (586\,B)$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a_{45}')^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G_{45}^*\mathbb{T}_{45} \quad (586\,C)$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a_{46}')^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G_{46}^*\mathbb{T}_{45} \quad (586\,D)$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b_{44}')^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T_{44}^*\mathbb{G}_j\right) \quad (586\,E)$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b_{45}')^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T_{45}^*\mathbb{G}_j\right) \quad (586\,F)$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b_{46}')^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T_{46}^*\mathbb{G}_j\right) \quad (586\,G)$$

The characteristic equation of this system is (587)

$$\left((\lambda)^{(1)} + (b_{15}')^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a_{15}')^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(q_{13})^{(1)}G_{13}^*\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T_{14}^* + (b_{14})^{(1)}s_{(13),(14)}T_{14}^*\right)$$

$$+\left(\left((\lambda)^{(1)} + (a_{14}')^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G_{13}^* + (a_{13})^{(1)}(q_{14})^{(1)}G_{14}^*\right)$$

$$\left(\left((\lambda)^{(1)} + (b_{13}')^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T_{14}^* + (b_{14})^{(1)}s_{(13),(13)}T_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((b_{13}')^{(1)} + (b_{14}')^{(1)} - (r_{13})^{(1)} + (r_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$+\left(\left((\lambda)^{(1)}\right)^2 + \left((a_{13}')^{(1)} + (a_{14}')^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)} + (a_{13}')^{(1)} + (p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G_{14}^* + (a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G_{13}^*\right)$$

$$\left(\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^{*}_{14}+(b_{14})^{(1)}s_{(13),(15)}T^{*}_{13}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\right)\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left(\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)\right]$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^{*}_{17}\right)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^{*}_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$\left(\left((\lambda)^{(2)}\right)^{2}+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\right)$$

$$+\left(\left((\lambda)^{(2)}\right)^{2}+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^{*}_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^{*}_{16}\right)$$

$$\left(\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^{*}_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^{*}_{16}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\right)\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left(\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)\right]$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(21)}T^{*}_{21}\right)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^{*}_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^{*}_{21}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(20)}T^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$\left(\left((\lambda)^{(3)}\right)^{2}+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\right)$$

$$+\left(\left((\lambda)^{(3)}\right)^{2}+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\right)\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\right)\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\Big)$$

$$+\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\Big)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\Big)$$

$$+\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\Big)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\Big)\}=0$$

$$+$$

$$(\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\Big)\right]$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\Big)$$

$$+\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\Big)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\Big)$$

$$+\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\Big)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\Big)\}=0$$

$$+$$

$$(\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\Big)\right]$$

$$\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\Big)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)})\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

SECTION SIXTEEN

Electroweak-Symmetry Breaking Mechanism (Popularly Known As The Brout-Englert-Higgs Mechanism), Boson's Spin-Parity, ZZ And Γγ (Two-Photon), Et Al. : Deus Et Natua Non Faciunt Frusta -Domino Optimo Maximo Model

INTRODUCTION—VARIABLES USED

(1) The data collected by CMS have now been used to show that the boson's spin-parity **is even less (e)consistent** with 0− (excluded with greater than 99.8% confidence) and is also not consistent with 2+ (excluded with greater than 98.5% or 99.9% confidence for the simplest 2+ model with production modes via gluons or quarks respectively), 1+ or 1− (each of which is excluded with greater than 99.9% confidence for exotic spin-1 models).

(2) Also, in July 2012, combining the ZZ and γγ (two-photon) channels gave CMS an observation with a significance of 5σ. Now, using the full dataset collected until the end of 2012, the ZZ channel alone gives the observation of the boson a significance of 6.7σ. The mass of the new particle has also been measured precisely in this channel, giving us a value of 125.8 GeV. The new results include the measurement of the production mechanism of this particle, and the rate of production with subsequent decay into two Z bosons divided by the expected rate for a SM Higgs gives a ratio (μ) of around 0.91, **suggesting (eb) that the** particle behaves like a SM Higgs. In addition, CMS has excluded the existence of the SM Higgs boson in the range 130–827 GeV at 95% confidence level.

Decays **to** two W bosons: HIG-13-003

(3) This channel studies the decays of the Higgs boson **candidate (eb) into** two W bosons, each of which subsequently decay into a lepton and a neutrino. As the neutrino cannot be detected directly by CMS, the resolution of this channel is not as sharp as that of the ZZ or γγ channels. From studies of the full collected dataset, CMS excludes the SM Higgs boson in the mass range of 128–600 GeV at 95% confidence level, while observing an excess at around 125 GeV with a significance of 4σ.

Decays to two τ leptons: HIG-13-004

(4) The electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism) was proposed **to (eb) explain** how the heavy bosons (Ws and Zs) got their masses.

(5) However, this mechanism (electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism)) **can also(eb) explain** how elementary fermions (quarks or leptons) acquire mass.

(6) An important test of whether the discovered boson is indeed the SM Higgs is to observe **its decay (eb) into** fermions. In other words observation of the decay of Boson in to fermions **determines (eb)** whether discovered boson is SM Higgs.

(7) CMS performed a search for the decay of the new boson **into (eb)** τ leptons. When the discovery announcement was made in July 2012, the observation in this channel was compatible with both the existence and the absence of a SM Higgs. Since then, with the additional data recorded by CMS, a clear signal in this channel has emerged for a boson with a mass of around 125 GeV, corresponding to a significance of 2.8σ, lending further credence to the expectation that the particle discovered **behaves like** a SM Higgs boson.

Searches for new physics

The Standard Model of particle physics has stood tall for decades, with successful prediction after successful prediction. However, we know that this is not the whole story and that new physics — that which is beyond the Standard Model — must exist out there. The LHC is a discovery machine, and should such new physics exist within the energies that can be probed here, we expect to find it.

Module One

G_{13} : Category one of 0− (excluded with greater than 99.8% confidence) and is also not consistent with 2+ (excluded with greater than 98.5% or 99.9% confidence for the simplest 2+ model with production modes via gluons or quarks respectively), 1+ or 1− (each of which is excluded with greater than 99.9% confidence for exotic spin-1 models).

G_{14} : Category two of 0− (excluded with greater than 99.8% confidence) and is also not consistent with 2+ (excluded with greater than 98.5% or 99.9% confidence for the simplest 2+ model with production modes via gluons or quarks respectively), 1+ or 1− (each of which is excluded with greater than 99.9% confidence for exotic spin-1 models).

G_{15} : Category three of 0− (excluded with greater than 99.8% confidence) and is also not consistent with 2+ (excluded with greater than 98.5% or 99.9% confidence for the simplest 2+ model with production modes via gluons or quarks respectively), 1+ or 1− (each of which is excluded with greater than 99.9% confidence for exotic spin-1 models).

T_{13} : Category one of data collected by CMS have now been used to show that the boson's spin-parity

T_{14} : Category two of data collected by CMS have now been used to show that the boson's spin-parity

T_{15} : Category three of data collected by CMS have now been used to show that the boson's spin-parity

Module Two

G_{16} : Category one of rate of production with subsequent decay into two Z bosons divided by the expected rate for a SM Higgs gives a ratio (μ) of around 0.91

G_{17} : Category two of rate of production with subsequent decay into two Z bosons divided by the expected rate for a SM Higgs gives a ratio (μ) of around 0.91

G_{18} : Category three of rate of production with subsequent decay into two Z bosons divided by the expected rate for a SM Higgs gives a ratio (μ) of around 0.91

T_{16} : Category one of particle behaves like a SM Higgs. In addition, CMS has excluded the existence of the SM Higgs boson in the range 130–827 GeV at 95% confidence level

T_{17} : Category two of particle behaves like a SM Higgs. In addition, CMS has excluded the existence of the SM Higgs boson in the range 130–827 GeV at 95% confidence level

T_{18} : Category three of particle behaves like a SM Higgs. In addition, CMS has excluded the existence of the SM Higgs boson in the range 130–827 GeV at 95% confidence level

Module three

G_{20} : Category one of Higgs boson candidate If the characterstics of the experiment and environment is same, then accentuation coefficient and Detrition coefficient is taken as zero. We talk of experiments that are held for the Higgs Boson candidate, and note that characterstics of those experiments are taken in to consideration in the classification.

G_{21} : Category two of Higgs boson candidate

G_{22} : Category three of Higgs boson candidate

T_{20} : Category one of two W bosons, each of which subsequently decays into a lepton and a neutrino. As the neutrino cannot be detected directly by CMS, the resolution of this channel is not as sharp as that of the ZZ or γγ channels. From studies of the full collected dataset, CMS excludes the SM Higgs boson in the mass range of 128–600 GeV at 95% confidence level, while observing an excess at around

125 GeV with a significance of 4σ

T_{21} : Category two of two W bosons, each of which subsequently decays into a lepton and a neutrino. As the neutrino cannot be detected directly by CMS, the resolution of this channel is not as sharp as that of the ZZ or γγ channels. From studies of the full collected dataset, CMS excludes the SM Higgs boson in the mass range of 128–600 GeV at 95% confidence level, while observing an excess at around 125 GeV with a significance of 4σ

T_{22} : Category three of two W bosons, each of which subsequently decays into a lepton and a neutrino. As the neutrino cannot be detected directly by CMS, the resolution of this channel is not as sharp as that of the ZZ or γγ channels. From studies of the full collected dataset, CMS excludes the SM Higgs boson in the mass range of 128–600 GeV at 95% confidence level, while observing an excess at around 125 GeV with a significance of 4σ

Module four

electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism) was proposed **to (eb) explain** how the heavy bosons (Ws and Zs) got their masses

G_{24} : Category one of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism)

G_{25} : Category two of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism)

G_{26} : Category three of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism)

T_{24} : Category one of heavy bosons (Ws and Zs) got their masses

T_{25} : Category two of heavy bosons (Ws and Zs) got their masses

T_{26} : Category three of heavy bosons (Ws and Zs) got their masses

Module five

G_{28} : Category one of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism

G_{29} : Category two of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism

G_{30} : Category three of electroweak-symmetry breaking mechanism (popularly known as the Brout-Englert-Higgs mechanism

T_{28} : Category one of elementary fermions (quarks or leptons) acquire mass

T_{29} : Category two of elementary fermions (quarks or leptons) acquire mass

T_{30} : Category three of elementary fermions (quarks or leptons) acquire mass

Module six

G_{32} : Category one of observation of decay in to fermions

G_{33} : Category two of observation of decay in to fermions

G_{34} : Category three of observation of decay in to fermions

T_{32} : Category one of test of whether the discovered boson is indeed the SM Higgs

T_{33} : Category two of test of whether the discovered boson is indeed the SM Higgs

T_{34} : Category three of test of whether the discovered boson is indeed the SM Higgs

Module seven

G_{36} : Category one of new boson There shall be new boson in various experiments' and they the experiments' would have characterstics which form the basis for classification theme.

G_{37} : Category two of new boson

G_{38} : Category three of new boson

T_{36} : Category one of τ leptons

T_{37} : Category two of τ leptons

T_{38} : Category three of τ leptons (It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

Module eight

Previous searches, utilizing 5 fb−1 of data collected at a centre-of-mass energy of 7 TeV in 2011, yielded no top or bottom squarks (the superpartners of the top and bottom quark, respectively, also known as stops and sbottoms). CMS has now studied the entire dataset from 2012 (~20 fb−1, 8 TeV) looking for these sparticles. If they are produced, we expect stops and sbottoms **to decay (eb) into** bottom quarks, producing particle jets (b-jets, in this case).

G_{40} : Category one of stops and sbottoms

G_{41} : Category two of stops and sbottoms

G_{42} : Category three of stops and sbottoms

T_{40} : Category one of bottom quarks, producing particle jets (b-jets, in this case).

T_{41} : Category two of bottom quarks, producing particle jets (b-jets, in this case).

T_{42} : Category three of bottom quarks, producing particle jets (b-jets, in this case).

Module Nine

These decays (stops and sbottoms **to decay (eb) into** bottom quarks, producing particle jets (b-jets, in this case)) are also expected **to (eb) produce** the lightest SUSY particle (LSP), which might leave no trace in the detector and will be characterised only by large missing energy

G_{44} : Category one of stops and sbottoms **to decay (eb) into** bottom quarks, producing particle jets (b-jets

G_{45} : Category two of stops and sbottoms **to decay (eb) into** bottom quarks, producing particle jets (b-

jets

G_{46} : Category three of stops and sbottoms **to decay (eb) into** bottom quarks, producing particle jets (b-jets

T_{44} : Category one of lightest SUSY particle (LSP), which might leave no trace in the detector and will be characterised only by large missing energy

T_{45} : Category two of lightest SUSY particle (LSP), which might leave no trace in the detector and will be characterised only by large missing energy

T_{46} : Category three of lightest SUSY particle (LSP), which might leave no trace in the detector and will be characterised only by large missing energy(It is to be noted in unmistakable terms for unequivocal recordial evidence that we are talking of the systems for which the given Theorem, inequality holds. For detailed scientific and philosophical explanation, reader is referred to the introduction).

.

The Coefficients:

$(a_{13})^{(1)}, (a_{14})^{(1)}, (a_{15})^{(1)}, (b_{13})^{(1)}, (b_{14})^{(1)}, (b_{15})^{(1)}\ (a_{16})^{(2)}, (a_{17})^{(2)}, (a_{18})^{(2)}$
$(b_{16})^{(2)}, (b_{17})^{(2)}, (b_{18})^{(2)}: (a_{20})^{(3)}, (a_{21})^{(3)}, (a_{22})^{(3)}\ ,\ (b_{20})^{(3)}, (b_{21})^{(3)}, (b_{22})^{(3)}$
$(a_{24})^{(4)}, (a_{25})^{(4)}, (a_{26})^{(4)}, (b_{24})^{(4)}, (b_{25})^{(4)}, (b_{26})^{(4)}, (b_{28})^{(5)}, (b_{29})^{(5)}, (b_{30})^{(5)},$
$(a_{28})^{(5)}, (a_{29})^{(5)}, (a_{30})^{(5)}, (a_{32})^{(6)}, (a_{33})^{(6)}, (a_{34})^{(6)}, (b_{32})^{(6)}, (b_{33})^{(6)}, (b_{34})^{(6)}$
$(a_{36})^{(7)}, (a_{37})^{(7)}, (a_{38})^{(7)}, (b_{36})^{(7)}, (b_{37})^{(7)}, (b_{38})^{(7)}$
$(a_{40})^{(8)}, (a_{41})^{(8)}, (a_{42})^{(8)}, (b_{40})^{(8)}, (b_{41})^{(8)}, (b_{42})^{(8)}$
$(a_{44})^{(9)}, (a_{45})^{(9)}, (a_{46})^{(9)}, (b_{44})^{(9)}, (b_{45})^{(9)}, (b_{46})^{(9)}$

are Accentuation coefficients
$(a'_{13})^{(1)}, (a'_{14})^{(1)}, (a'_{15})^{(1)}, (b'_{13})^{(1)}, (b'_{14})^{(1)}, (b'_{15})^{(1)}, (a'_{16})^{(2)}, (a'_{17})^{(2)}, (a'_{18})^{(2)},$
$(b'_{16})^{(2)}, (b'_{17})^{(2)}, (b'_{18})^{(2)}\ , (a'_{20})^{(3)}, (a'_{21})^{(3)}, (a'_{22})^{(3)}, (b'_{20})^{(3)}, (b'_{21})^{(3)}, (b'_{22})^{(3)}$
$(a'_{24})^{(4)}, (a'_{25})^{(4)}, (a'_{26})^{(4)}, (b'_{24})^{(4)}, (b'_{25})^{(4)}, (b'_{26})^{(4)}, (b'_{28})^{(5)}, (b'_{29})^{(5)}, (b'_{30})^{(5)}$
$(a'_{28})^{(5)}, (a'_{29})^{(5)}, (a'_{30})^{(5)}\ , (a'_{32})^{(6)}, (a'_{33})^{(6)}, (a'_{34})^{(6)}, (b'_{32})^{(6)}, (b'_{33})^{(6)}, (b'_{34})^{(6)}$
$(a'_{36})^{(7)}, (a'_{37})^{(7)}, (a'_{38})^{(7)}, (b'_{36})^{(7)}, (b'_{37})^{(7)}, (b'_{38})^{(7)},$
$(a'_{40})^{(8)}, (a'_{41})^{(8)}, (a'_{42})^{(8)}, (b'_{40})^{(8)}, (b'_{41})^{(8)}, (b'_{42})^{(8)},$
$(a'_{44})^{(9)}, (a'_{45})^{(9)}, (a'_{46})^{(9)}, (b'_{44})^{(9)}, (b'_{45})^{(9)}, (b'_{46})^{(9)},$
are Dissipation coefficients

Module Numbered One

The differential system of this model is now (Module Numbered one)

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14},t)\right]G_{13} \quad (1)$$
$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14},t)\right]G_{14} \quad (2)$$
$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14},t)\right]G_{15} \quad (3)$$
$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \left[(b'_{13})^{(1)} - (b''_{13})^{(1)}(G,t)\right]T_{13} \quad (4)$$
$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \left[(b'_{14})^{(1)} - (b''_{14})^{(1)}(G,t)\right]T_{14} \quad (5)$$
$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \left[(b'_{15})^{(1)} - (b''_{15})^{(1)}(G,t)\right]T_{15} \quad (6)$$

$+(a''_{13})^{(1)}(T_{14},t) =$ First augmentation factor

$-(b''_{13})^{(1)}(G,t) =$ First detritions factor

Module Numbered Two

The differential system of this model is now (Module numbered two)

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17},t)\right]G_{16} \quad (7)$$
$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17},t)\right]G_{17} \quad (8)$$
$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17},t)\right]G_{18} \quad (9)$$
$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \left[(b'_{16})^{(2)} - (b''_{16})^{(2)}\big((G_{19}),t\big)\right]T_{16} \quad (10)$$
$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \left[(b'_{17})^{(2)} - (b''_{17})^{(2)}\big((G_{19}),t\big)\right]T_{17} \quad (11)$$
$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \left[(b'_{18})^{(2)} - (b''_{18})^{(2)}\big((G_{19}),t\big)\right]T_{18} \quad (12)$$

$+(a''_{16})^{(2)}(T_{17},t) =$ First augmentation factor

$-(b''_{16})^{(2)}\big((G_{19}),t\big) =$ First detritions factor

Module Numbered Three

The differential system of this model is now (Module numbered three)

$$\frac{dG_{20}}{dt} = (a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21},t)\right]G_{20} \quad (13)$$
$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21},t)\right]G_{21} \quad (14)$$
$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21},t)\right]G_{22} \quad (15)$$
$$\frac{dT_{20}}{dt} = (b_{20})^{(3)}T_{21} - \left[(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23},t)\right]T_{20} \quad (16)$$
$$\frac{dT_{21}}{dt} = (b_{21})^{(3)}T_{20} - \left[(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23},t)\right]T_{21} \quad (17)$$
$$\frac{dT_{22}}{dt} = (b_{22})^{(3)}T_{21} - \left[(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23},t)\right]T_{22} \quad (18)$$

$+(a''_{20})^{(3)}(T_{21},t) =$ First augmentation factor

$-(b''_{20})^{(3)}(G_{23},t) =$ First detritions factor

Module Numbered Four

The differential system of this model is now (Module numbered Four)

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25},t)\right]G_{24} \quad (19)$$
$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25},t)\right]G_{25} \quad (20)$$
$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25},t)\right]G_{26} \quad (21)$$
$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \left[(b'_{24})^{(4)} - (b''_{24})^{(4)}\big((G_{27}),t\big)\right]T_{24} \quad (22)$$
$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \left[(b'_{25})^{(4)} - (b''_{25})^{(4)}\big((G_{27}),t\big)\right]T_{25} \quad (23)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \left[(b'_{26})^{(4)} - (b''_{26})^{(4)}((G_{27}),t)\right]T_{26} \quad 24$$

$+(a''_{24})^{(4)}(T_{25},t) =$ **First augmentation factor**

$-(b''_{24})^{(4)}((G_{27}),t) =$ **First detritions factor**

Module Numbered Five:

The differential system of this model is now (Module number five)

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29},t)\right]G_{28} \quad 25$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29},t)\right]G_{29} \quad 26$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29},t)\right]G_{30} \quad 27$$

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[(b'_{28})^{(5)} - (b''_{28})^{(5)}((G_{31}),t)\right]T_{28} \quad 28$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[(b'_{29})^{(5)} - (b''_{29})^{(5)}((G_{31}),t)\right]T_{29} \quad 29$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[(b'_{30})^{(5)} - (b''_{30})^{(5)}((G_{31}),t)\right]T_{30} \quad 30$$

$+(a''_{28})^{(5)}(T_{29},t) =$ **First augmentation factor**

$-(b''_{28})^{(5)}((G_{31}),t) =$ **First detritions factor**

Module Numbered Six

The differential system of this model is now (Module numbered Six)

$$\frac{dG_{32}}{dt} = (a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33},t)\right]G_{32} \quad 31$$

$$\frac{dG_{33}}{dt} = (a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33},t)\right]G_{33} \quad 32$$

$$\frac{dG_{34}}{dt} = (a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33},t)\right]G_{34} \quad 33$$

$$\frac{dT_{32}}{dt} = (b_{32})^{(6)}T_{33} - \left[(b'_{32})^{(6)} - (b''_{32})^{(6)}((G_{35}),t)\right]T_{32} \quad 34$$

$$\frac{dT_{33}}{dt} = (b_{33})^{(6)}T_{32} - \left[(b'_{33})^{(6)} - (b''_{33})^{(6)}((G_{35}),t)\right]T_{33} \quad 35$$

$$\frac{dT_{34}}{dt} = (b_{34})^{(6)}T_{33} - \left[(b'_{34})^{(6)} - (b''_{34})^{(6)}((G_{35}),t)\right]T_{34} \quad 36$$

$+(a''_{32})^{(6)}(T_{33},t) =$ **First augmentation factor**

Module Numbered Seven:

The differential system of this model is now (Seventh Module)

$$\frac{dG_{36}}{dt} = (a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37},t)\right]G_{36} \quad 37$$

$$\frac{dG_{37}}{dt} = (a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37},t)\right]G_{37} \quad 38$$

$$\frac{dG_{38}}{dt} = (a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37},t)\right]G_{38} \quad 39$$

$$\frac{dT_{36}}{dt} = (b_{36})^{(7)}T_{37} - \left[(b'_{36})^{(7)} - (b''_{36})^{(7)}((G_{39}),t)\right]T_{36} \quad 40$$

$$\frac{dT_{37}}{dt} = (b_{37})^{(7)}T_{36} - \left[(b'_{37})^{(7)} - (b''_{37})^{(7)}((G_{39}),t)\right]T_{37} \quad 41$$

$$\frac{dT_{38}}{dt} = (b_{38})^{(7)}T_{37} - \left[(b'_{38})^{(7)} - (b''_{38})^{(7)}((G_{39}),t)\right]T_{38} \quad 42$$

$+(a''_{36})^{(7)}(T_{37},t) =$ **First augmentation factor**

Module Numbered Eight

The differential system of this model is now

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41},t)\right]G_{40} \quad 43$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41},t)\right]G_{41} \quad 44$$

$$\frac{dG_{42}}{dt} = (a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41},t)\right]G_{42} \quad 45$$

$$\frac{dT_{40}}{dt} = (b_{40})^{(8)}T_{41} - \left[(b'_{40})^{(8)} - (b''_{40})^{(8)}((G_{43}),t)\right]T_{40} \quad 46$$

$$\frac{dT_{41}}{dt} = (b_{41})^{(8)}T_{40} - \left[(b'_{41})^{(8)} - (b''_{41})^{(8)}\left((G_{43}),t\right)\right]T_{41} \quad 47$$

$$\frac{dT_{42}}{dt} = (b_{42})^{(8)}T_{41} - \left[(b'_{42})^{(8)} - (b''_{42})^{(8)}\left((G_{43}),t\right)\right]T_{42} \quad 48$$

Module Numbered Nine

The differential system of this model is now

$$\frac{dG_{44}}{dt} = (a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45},t)\right]G_{44} \quad 49$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45},t)\right]G_{45} \quad 50$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45},t)\right]G_{46} \quad 51$$

$$\frac{dT_{44}}{dt} = (b_{44})^{(9)}T_{45} - \left[(b'_{44})^{(9)} - (b''_{44})^{(9)}\left((G_{47}),t\right)\right]T_{44} \quad 52$$

$$\frac{dT_{45}}{dt} = (b_{45})^{(9)}T_{44} - \left[(b'_{45})^{(9)} - (b''_{45})^{(9)}\left((G_{47}),t\right)\right]T_{45} \quad 53$$

$$\frac{dT_{46}}{dt} = (b_{46})^{(9)}T_{45} - \left[(b'_{46})^{(9)} - (b''_{46})^{(9)}\left((G_{47}),t\right)\right]T_{46} \quad 54$$

$+(a''_{44})^{(9)}(T_{45},t) =$ **First augmentation factor**

$-(b''_{44})^{(9)}\left((G_{47}),t\right) =$ **First detrition factor**

$$\frac{dG_{13}}{dt} = (a_{13})^{(1)}G_{14} - \begin{bmatrix} (a'_{13})^{(1)} & \boxed{+(a''_{13})^{(1)}(T_{14},t)} & \boxed{+(a''_{16})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13} \quad 55$$

$$\frac{dG_{14}}{dt} = (a_{14})^{(1)}G_{13} - \begin{bmatrix} (a'_{14})^{(1)} & \boxed{+(a''_{14})^{(1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14} \quad 56$$

$$\frac{dG_{15}}{dt} = (a_{15})^{(1)}G_{14} - \begin{bmatrix} (a'_{15})^{(1)} & \boxed{+(a''_{15})^{(1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15} \quad 57$$

Where $\boxed{(a''_{13})^{(1)}(T_{14},t)}$, $\boxed{(a''_{14})^{(1)}(T_{14},t)}$, $\boxed{(a''_{15})^{(1)}(T_{14},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{38})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{37})^{(7,7)}(T_{37},t)}$ $\boxed{+(a''_{36})^{(7,7)}(T_{37},t)}$ are seventh augmentation coefficient for 1,2,3

$\boxed{+(a''_{40})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{41})^{(8,8)}(T_{41},t)}$ $\boxed{+(a''_{42})^{(8,8)}(T_{41},t)}$ are eight augmentation coefficient for 1,2,3

$\boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{13}}{dt} = (b_{13})^{(1)}T_{14} - \begin{bmatrix} (b'_{13})^{(1)}\boxed{-(b''_{13})^{(1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13} \quad 58$$

$$\frac{dT_{14}}{dt} = (b_{14})^{(1)}T_{13} - \begin{bmatrix} (b'_{14})^{(1)}\boxed{-(b''_{14})^{(1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14} \quad 59$$

$$\frac{dT_{15}}{dt} = (b_{15})^{(1)}T_{14} - \begin{bmatrix} (b'_{15})^{(1)}\boxed{-(b''_{15})^{(1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)} \\ \boxed{-(b''_{38})^{(7,7,)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15} \quad 60$$

Where $\boxed{-(b''_{13})^{(1)}(G,t)}$, $\boxed{-(b''_{14})^{(1)}(G,t)}$, $\boxed{-(b''_{15})^{(1)}(G,t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{37})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{36})^{(7,7,)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{41})^{(8,8)}(G_{43},t)}$ $\boxed{-(b''_{42})^{(8,8)}(G_{43},t)}$ are eight detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{44})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{16}}{dt} = (a_{16})^{(2)}G_{17} - \begin{bmatrix} (a'_{16})^{(2)}\boxed{+(a''_{16})^{(2)}(T_{17},t)} & \boxed{+(a''_{13})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{20})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{16} \quad 61$$

$$\frac{dG_{17}}{dt} = (a_{17})^{(2)}G_{16} - \begin{bmatrix} (a'_{17})^{(2)}\boxed{+(a''_{17})^{(2)}(T_{17},t)} & \boxed{+(a''_{14})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{21})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{17} \quad 62$$

$$\frac{dG_{18}}{dt} = (a_{18})^{(2)}G_{17} - \begin{bmatrix} (a'_{18})^{(2)}\boxed{+(a''_{18})^{(2)}(T_{17},t)} & \boxed{+(a''_{15})^{(1,1,)}(T_{14},t)} & \boxed{+(a''_{22})^{(3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9)}(T_{45},t)} \end{bmatrix} G_{18} \quad 63$$

Where $\boxed{+(a''_{16})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2)}(T_{17},t)}$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{13})^{(1,1,)}(T_{14},t)$, $+(a''_{14})^{(1,1,)}(T_{14},t)$, $+(a''_{15})^{(1,1,)}(T_{14},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{36})^{(7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7)}(T_{37},t)$ are seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{40})^{(8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8)}(T_{41},t)$ are eight augmentation coefficient for category 1, 2 and 3

$+(a''_{44})^{(9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9)}(T_{45},t)$, $+(a''_{46})^{(9,9)}(T_{45},t)$ are ninth augmentation coefficient for category 1, 2 and 3

$$\frac{dT_{16}}{dt} = (b_{16})^{(2)}T_{17} - \begin{bmatrix} (b'_{16})^{(2)} & -(b''_{16})^{(2)}(G_{19},t) & -(b''_{13})^{(1,1,)}(G,t) & -(b''_{20})^{(3,3,3,)}(G_{23},t) \\ -(b''_{24})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{32})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{36})^{(7,7,7)}(G_{39},t) & -(b''_{40})^{(8,8,8)}(G_{43},t) & -(b''_{44})^{(9,9)}(G_{47},t) \end{bmatrix} T_{16} \quad 64$$

$$\frac{dT_{17}}{dt} = (b_{17})^{(2)}T_{16} - \begin{bmatrix} (b'_{17})^{(2)} & -(b''_{17})^{(2)}(G_{19},t) & -(b''_{14})^{(1,1,)}(G,t) & -(b''_{21})^{(3,3,3,)}(G_{23},t) \\ -(b''_{25})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{33})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{37})^{(7,7,7)}(G_{39},t) & -(b''_{41})^{(8,8,8)}(G_{43},t) & -(b''_{45})^{(9,9)}(G_{47},t) \end{bmatrix} T_{17} \quad 65$$

$$\frac{dT_{18}}{dt} = (b_{18})^{(2)}T_{17} - \begin{bmatrix} (b'_{18})^{(2)} & -(b''_{18})^{(2)}(G_{19},t) & -(b''_{15})^{(1,1,)}(G,t) & -(b''_{22})^{(3,3,3,)}(G_{23},t) \\ -(b''_{26})^{(4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6)}(G_{35},t) \\ -(b''_{38})^{(7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9)}(G_{47},t) \end{bmatrix} T_{18} \quad 66$$

where $-(b''_{16})^{(2)}(G_{19},t)$, $-(b''_{17})^{(2)}(G_{19},t)$, $-(b''_{18})^{(2)}(G_{19},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{13})^{(1,1,)}(G,t)$, $-(b''_{14})^{(1,1,)}(G,t)$, $-(b''_{15})^{(1,1,)}(G,t)$ are second detrition coefficients for category 1,2 and 3

$-(b''_{20})^{(3,3,3,)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,)}(G_{23},t)$ are third detrition coefficients for category 1,2 and 3

$-(b''_{24})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1,2 and 3

$-(b''_{28})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1,2 and 3

$-(b''_{32})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1,2 and 3

$-(b''_{36})^{(7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2 and 3

$-(b''_{40})^{(8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1,2 and 3

$\boxed{-(b''_{44})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{46})^{(9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1,2 and 3

$$\frac{dG_{20}}{dt} = \qquad (67)$$

$$(a_{20})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{20})^{(3)} & \boxed{+(a''_{20})^{(3)}(T_{21},t)} & \boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{20}$$

$$\frac{dG_{21}}{dt} = (a_{21})^{(3)}G_{20} - \left[\begin{array}{ccc} (a'_{21})^{(3)} & \boxed{+(a''_{21})^{(3)}(T_{21},t)} & \boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{21} \qquad (68)$$

$$\frac{dG_{22}}{dt} = (a_{22})^{(3)}G_{21} - \left[\begin{array}{ccc} (a'_{22})^{(3)} & \boxed{+(a''_{22})^{(3)}(T_{21},t)} & \boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)} \quad \boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)} \end{array}\right] G_{22} \qquad (69)$$

$\boxed{+(a''_{20})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3)}(T_{21},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2)}(T_{17},t)}$ are second augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,)}(T_{14},t)}$ are third augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{36})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{40})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8)}(T_{41},t)}$ are eight augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{44})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1, 2 and 3

$$\frac{dT_{20}}{dt} = \qquad (70)$$

$$(b_{20})^{(3)}T_{21} - \left[\begin{array}{ccc} (b'_{20})^{(3)} & \boxed{-(b''_{20})^{(3)}(G_{23},t)} & \boxed{-(b''_{16})^{(2,2,2)}(G_{19},t)} \quad \boxed{-(b''_{13})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{20}$$

$$\frac{dT_{21}}{dt} = \qquad (71)$$

$$(b_{21})^{(3)}T_{20} - \left[\begin{array}{ccc} (b'_{21})^{(3)} & \boxed{-(b''_{21})^{(3)}(G_{23},t)} & \boxed{-(b''_{17})^{(2,2,2)}(G_{19},t)} \quad \boxed{-(b''_{14})^{(1,1,1,)}(G,t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9)}(G_{47},t)} \end{array}\right] T_{21}$$

$$\frac{dT_{22}}{dt} = \qquad (72)$$

$$(b_{22})^{(3)}T_{21} - \begin{bmatrix} (b'_{22})^{(3)} & -(b''_{22})^{(3)}(G_{23},t) & -(b''_{18})^{(2,2,2)}(G_{19},t) & -(b''_{15})^{(1,1,1,)}(G,t) \\ -(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t) & -(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t) & \\ -(b''_{38})^{(7,7,7,7)}(G_{39},t) & -(b''_{42})^{(8,8,8,8)}(G_{43},t) & -(b''_{46})^{(9,9,9)}(G_{47},t) & \end{bmatrix} T_{22}$$

$-(b''_{20})^{(3)}(G_{23},t)$, $-(b''_{21})^{(3)}(G_{23},t)$, $-(b''_{22})^{(3)}(G_{23},t)$ *are first detrition coefficients for category 1, 2 and 3*

$-(b''_{16})^{(2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2)}(G_{19},t)$ *are second detrition coefficients for category 1, 2 and 3*

$-(b''_{13})^{(1,1,1,)}(G,t)$, $-(b''_{14})^{(1,1,1,)}(G,t)$, $-(b''_{15})^{(1,1,1,)}(G,t)$ *are third detrition coefficients for category 1,2 and 3*

$-(b''_{24})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4)}(G_{27},t)$ *are fourth detrition coefficients for category 1, 2 and 3*

$-(b''_{28})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5)}(G_{31},t)$ *are fifth detrition coefficients for category 1, 2 and 3*

$-(b''_{32})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6)}(G_{35},t)$ *are sixth detrition coefficients for category 1, 2 and 3*

$-(b''_{36})^{(7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7)}(G_{39},t)$ $-(b''_{38})^{(7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1, 2 and 3

$-(b''_{40})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8)}(G_{43},t)$ are eight detrition coefficients for category 1, 2 and 3

$-(b''_{46})^{(9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{24}}{dt} = (a_{24})^{(4)}G_{25} - \begin{bmatrix} (a'_{24})^{(4)} & +(a''_{24})^{(4)}(T_{25},t) & +(a''_{28})^{(5,5,)}(T_{29},t) & +(a''_{32})^{(6,6,)}(T_{33},t) \\ +(a''_{13})^{(1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{36})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{24} \quad 73$$

$$\frac{dG_{25}}{dt} = (a_{25})^{(4)}G_{24} - \begin{bmatrix} (a'_{25})^{(4)} & +(a''_{25})^{(4)}(T_{25},t) & +(a''_{29})^{(5,5,)}(T_{29},t) & +(a''_{33})^{(6,6)}(T_{33},t) \\ +(a''_{14})^{(1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{37})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{25} \quad 74$$

$$\frac{dG_{26}}{dt} = (a_{26})^{(4)}G_{25} - \begin{bmatrix} (a'_{26})^{(4)} & +(a''_{26})^{(4)}(T_{25},t) & +(a''_{30})^{(5,5,)}(T_{29},t) & +(a''_{34})^{(6,6,)}(T_{33},t) \\ +(a''_{15})^{(1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3)}(T_{21},t) & \\ +(a''_{38})^{(7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9)}(T_{45},t) & \end{bmatrix} G_{26} \quad 75$$

$(a''_{24})^{(4)}(T_{25},t)$, $(a''_{25})^{(4)}(T_{25},t)$, $(a''_{26})^{(4)}(T_{25},t)$ *are first augmentation coefficients category* 1, 2 3

$+(a''_{28})^{(5,5,)}(T_{29},t)$, $+(a''_{29})^{(5,5,)}(T_{29},t)$, $+(a''_{30})^{(5,5,)}(T_{29},t)$ *are second augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{32})^{(6,6,)}(T_{33},t)$, $+(a''_{33})^{(6,6,)}(T_{33},t)$, $+(a''_{34})^{(6,6,)}(T_{33},t)$ *are third augmentation coefficient for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1)}(T_{14},t)$
are fourth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{16})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2)}(T_{17},t)$
are fifth augmentation coefficients for category 1, 2 *and* 3

$+(a''_{20})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3)}(T_{21},t)$
are sixth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{36})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7)}(T_{37},t)$
are seventh augmentation coefficients for category 1, 2 *and* 3
$+(a''_{40})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8)}(T_{41},t)$
are eighth augmentation coefficients for category 1, 2 *and* 3
$+(a''_{46})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9)}(T_{45},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dT_{24}}{dt} = (b_{24})^{(4)}T_{25} - \begin{bmatrix} (b'_{24})^{(4)}\left[-(b''_{24})^{(4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{16})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{44})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{24} \quad (76)$$

$$\frac{dT_{25}}{dt} = (b_{25})^{(4)}T_{24} - \begin{bmatrix} (b'_{25})^{(4)}\left[-(b''_{25})^{(4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{17})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{45})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{25} \quad (77)$$

$$\frac{dT_{26}}{dt} = (b_{26})^{(4)}T_{25} - \begin{bmatrix} (b'_{26})^{(4)}\left[-(b''_{26})^{(4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1)}(G,t)\right] & \left[-(b''_{18})^{(2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)\right] & \left[-(b''_{46})^{(9,9,9,9)}(G_{47},t)\right] \end{bmatrix} T_{26} \quad (78)$$

Where $-(b''_{24})^{(4)}(G_{27},t)$, $-(b''_{25})^{(4)}(G_{27},t)$, $-(b''_{26})^{(4)}(G_{27},t)$ *are first detrition coefficients for category* 1, 2 *and* 3
$-(b''_{28})^{(5,5,)}(G_{31},t)$, $-(b''_{29})^{(5,5,)}(G_{31},t)$, $-(b''_{30})^{(5,5,)}(G_{31},t)$ *are second detrition coefficients for category* 1, 2 *and* 3
$-(b''_{32})^{(6,6,)}(G_{35},t)$, $-(b''_{33})^{(6,6,)}(G_{35},t)$, $-(b''_{34})^{(6,6,)}(G_{35},t)$ *are third detrition coefficients for category* 1, 2 *and* 3
$-(b''_{13})^{(1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1)}(G,t)$
are fourth detrition coefficients for category 1, 2 *and* 3
$-(b''_{16})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2)}(G_{19},t)$
are fifth detrition coefficients for category 1, 2 *and* 3
$-(b''_{20})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3)}(G_{23},t)$
are sixth detrition coefficients for category 1, 2 *and* 3
$-(b''_{36})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7)}(G_{39},t)$
are seventh detrition coefficients for category 1, 2 *and* 3
$-(b''_{40})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8)}(G_{43},t)$, $-(b''_{42})^{(8,8,8,8,8)}(G_{43},t)$
are eighth detrition coefficients for category 1, 2 *and* 3
$-(b''_{46})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1 2 3

$$\frac{dG_{28}}{dt} = (a_{28})^{(5)}G_{29} - \begin{bmatrix} (a'_{28})^{(5)}\left[+(a''_{28})^{(5)}(T_{29},t)\right] & \left[+(a''_{24})^{(4,4,)}(T_{25},t)\right] & \left[+(a''_{32})^{(6,6,6)}(T_{33},t)\right] \\ \left[+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)\right] & \left[+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)\right] & \left[+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)\right] \\ \left[+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)\right] & \left[+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)\right] & \left[+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)\right] \end{bmatrix} G_{28} \quad (79)$$

$$\frac{dG_{29}}{dt} = (a_{29})^{(5)}G_{28} - \left[\begin{array}{ccc} (a'_{29})^{(5)}\boxed{+(a''_{29})^{(5)}(T_{29},t)} & \boxed{+(a''_{25})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{29} \quad (80)$$

$$\frac{dG_{30}}{dt} = (a_{30})^{(5)}G_{29} - \left[\begin{array}{ccc} (a'_{30})^{(5)}\boxed{+(a''_{30})^{(5)}(T_{29},t)} & \boxed{+(a''_{26})^{(4,4,)}(T_{25},t)} & \boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)} \end{array}\right] G_{30} \quad (81)$$

Where $\boxed{+(a''_{28})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5)}(T_{29},t)}$ *are first augmentation coefficients for category* 1, 2 *and* 3

And $\boxed{+(a''_{24})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,)}(T_{25},t)}$ *are second augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{32})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6)}(T_{33},t)}$ *are third augmentation coefficient for category* 1, 2 *and* 3

$\boxed{+(a''_{13})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{14})^{(1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1)}(T_{14},t)}$ are fourth augmentation coefficients for category 1,2, and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2)}(T_{17},t)}$ are fifth augmentation coefficients for category 1,2,and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3)}(T_{21},t)}$ are sixth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{36})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{37})^{(7,7,7,7,7,7)}(T_{37},t)}$, $\boxed{+(a''_{38})^{(7,7,7,7,7,7)}(T_{37},t)}$ are seventh augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{40})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{42})^{(8,8,8,8,8,8)}(T_{41},t)}$ are eighth augmentation coefficients for category 1,2, 3

$\boxed{+(a''_{46})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficients for category 1,2, 3

$$\frac{dT_{28}}{dt} = (b_{28})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{28})^{(5)}\boxed{-(b''_{28})^{(5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{28} \quad (82)$$

$$\frac{dT_{29}}{dt} = (b_{29})^{(5)}T_{28} - \left[\begin{array}{ccc} (b'_{29})^{(5)}\boxed{-(b''_{29})^{(5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{29} \quad (83)$$

$$\frac{dT_{30}}{dt} = (b_{30})^{(5)}T_{29} - \left[\begin{array}{ccc} (b'_{30})^{(5)}\boxed{-(b''_{30})^{(5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,)}(G_{27},t)} & \boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)} \end{array}\right] T_{30} \quad (84)$$

where $\boxed{-(b''_{28})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5)}(G_{31},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,)}(G_{27},t)}$ *are second detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{32})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6)}(G_{35},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$-(b''_{13})^{(1,1,1,1,1)}(G,t)$, $-(b''_{14})^{(1,1,1,1,1)}(G,t)$, $-(b''_{15})^{(1,1,1,1,1,)}(G,t)$ are fourth detrition coefficients for category 1,2, and 3

$-(b''_{16})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2)}(G_{19},t)$ are fifth detrition coefficients for category 1,2, and 3

$-(b''_{20})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3)}(G_{23},t)$ are sixth detrition coefficients for category 1,2, and 3

$-(b''_{36})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{37})^{(7,7,7,7,7,7)}(G_{39},t)$, $-(b''_{38})^{(7,7,7,7,7,7)}(G_{39},t)$ are seventh detrition coefficients for category 1,2, and 3

$-(b''_{42})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{41})^{(8,8,8,8,8,8)}(G_{43},t)$, $-(b''_{40})^{(8,8,8,8,8,8)}(G_{43},t)$ are eighth detrition coefficients for category 1,2, and 3

$-(b''_{46})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{45})^{(9,9,9,9,9)}(G_{47},t)$, $-(b''_{44})^{(9,9,9,9,9)}(G_{47},t)$ are ninth detrition coefficients for category 1,2, and 3

$$\frac{dG_{32}}{dt} \qquad 85$$

$$= (a_{32})^{(6)}G_{33} - \begin{bmatrix} (a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33},t) & +(a''_{28})^{(5,5,5)}(T_{29},t) & +(a''_{24})^{(4,4,4,)}(T_{25},t) \\ +(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{32}$$

$$\frac{dG_{33}}{dt} \qquad 86$$

$$= (a_{33})^{(6)}G_{32} - \begin{bmatrix} (a'_{33})^{(6)}+(a''_{33})^{(6)}(T_{33},t) & +(a''_{29})^{(5,5,5)}(T_{29},t) & +(a''_{25})^{(4,4,4,)}(T_{25},t) \\ +(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{33}$$

$$\frac{dG_{34}}{dt} \qquad 87$$

$$= (a_{34})^{(6)}G_{33} - \begin{bmatrix} (a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33},t) & +(a''_{30})^{(5,5,5)}(T_{29},t) & +(a''_{26})^{(4,4,4,)}(T_{25},t) \\ +(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t) & +(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t) & +(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t) \\ +(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t) & +(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t) & +(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t) \end{bmatrix} G_{34}$$

$+(a''_{32})^{(6)}(T_{33},t)$, $+(a''_{33})^{(6)}(T_{33},t)$, $+(a''_{34})^{(6)}(T_{33},t)$ *are first augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{28})^{(5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5)}(T_{29},t)$ *are second augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{24})^{(4,4,4,)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,)}(T_{25},t)$ *are third augmentation coefficients for category* 1, 2 *and* 3

$+(a''_{13})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1)}(T_{14},t)$ - are fourth augmentation coefficients

$+(a''_{16})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2)}(T_{17},t)$ - fifth augmentation coefficients

$+(a''_{20})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3)}(T_{21},t)$ sixth augmentation coefficients

$+(a''_{36})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7)}(T_{37},t)$ seventh augmentation coefficients

$+(a''_{40})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{42})^{(8,8,8,8,8,8,8)}(T_{41},t)$

Eighth augmentation coefficients
$\boxed{+(a''_{44})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{46})^{(9,9,9,9,9,9)}(T_{45},t)}$ ninth augmentation coefficients

$$\frac{dT_{32}}{dt} \qquad 88$$

$$= (b_{32})^{(6)}T_{33} - \begin{bmatrix} (b'_{32})^{(6)}\boxed{-(b''_{32})^{(6)}(G_{35},t)} & \boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{32}$$

$$\frac{dT_{33}}{dt} \qquad 89$$

$$= (b_{33})^{(6)}T_{32} - \begin{bmatrix} (b'_{33})^{(6)}\boxed{-(b''_{33})^{(6)}(G_{35},t)} & \boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{33}$$

$$\frac{dT_{34}}{dt} \qquad 90$$

$$= (b_{34})^{(6)}T_{33} - \begin{bmatrix} (b'_{34})^{(6)}\boxed{-(b''_{34})^{(6)}(G_{35},t)} & \boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)} & \boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{34}$$

$\boxed{-(b''_{32})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6)}(G_{35},t)}$ *are first detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{28})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5)}(G_{31},t)}$ *are second detrition coefficients for category* 1, 2 *and* 3

$\boxed{-(b''_{24})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,)}(G_{27},t)}$ *are third detrition coefficients for category* 1,2 *and* 3

$\boxed{-(b''_{13})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{15})^{(1,1,1,1,1,1)}(G,t)}$ are fourth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2)}(G_{19},t)}$ are fifth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3)}(G_{23},t)}$ are sixth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{36})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7,7,7,7,7,7,7)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2, and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2, and 3

$$\frac{dG_{36}}{dt} \quad 91$$

$$= (a_{36})^{(7)} G_{37} - \begin{bmatrix} (a'_{36})^{(7)} \boxed{+(a''_{36})^{(7)}(T_{37},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{37}}{dt} \quad 92$$

$$= (a_{37})^{(7)} G_{36} - \begin{bmatrix} (a'_{37})^{(7)} \boxed{+(a''_{37})^{(7)}(T_{37},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{38}}{dt} \quad 93$$

$$= (a_{38})^{(7)} G_{37} - \begin{bmatrix} (a'_{38})^{(7)} \boxed{+(a''_{38})^{(7)}(T_{37},t)} & \boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)} & \boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{15}$$

Where $\boxed{(a''_{36})^{(7)}(T_{37},t)}$, $\boxed{(a''_{37})^{(7)}(T_{37},t)}$, $\boxed{(a''_{38})^{(7)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a''_{16})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{17})^{(2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a''_{18})^{(2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{20})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{21})^{(3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a''_{22})^{(3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{24})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{25})^{(4,4,4,4,4,4,4)}(T_{25},t)}$, $\boxed{+(a''_{26})^{(4,4,4,4,4,4,4)}(T_{25},t)}$ are fourth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{28})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{29})^{(5,5,5,5,5,5,5)}(T_{29},t)}$, $\boxed{+(a''_{30})^{(5,5,5,5,5,5,5)}(T_{29},t)}$ are fifth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{32})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{33})^{(6,6,6,6,6,6,6)}(T_{33},t)}$, $\boxed{+(a''_{34})^{(6,6,6,6,6,6,6)}(T_{33},t)}$ are sixth augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{13})^{(1,1,1,1,1,1,1)}(T_{14},t)}$, $\boxed{+(a''_{15})^{(1,1,1,1,1,1,1)}(T_{14},t)}$ are seventh augmentation coefficient for category 1, 2 and 3

$\boxed{+(a''_{42})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{41})^{(8,8,8,8,8,8,8,8)}(T_{41},t)}$, $\boxed{+(a''_{40})^{(8,8,8,8,8,8,8,8,)}(T_{41},t)}$ are eighth augmentation coefficient for 1,2,3

$\boxed{+(a''_{46})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{45})^{(9,9,9,9,9,9,9)}(T_{45},t)}$, $\boxed{+(a''_{44})^{(9,9,9,9,9,9,9)}(T_{45},t)}$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{36}}{dt} = \quad 94$$

$$(b_{36})^{(7)} T_{37} - \begin{bmatrix} (b'_{36})^{(7)} \boxed{-(b''_{36})^{(7)}(G_{39},t)} & \boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{13}$$

$$\frac{dT_{37}}{dt} =$$

$$(b_{37})^{(7)}T_{36} - \begin{bmatrix} (b'_{37})^{(7)}\boxed{-(b''_{37})^{(7)}(G_{39},t)} & \boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{14}$$

$$\frac{dT_{38}}{dt} =$$

$$(b_{38})^{(7)}T_{37} - \begin{bmatrix} (b'_{38})^{(7)}\boxed{-(b''_{38})^{(7)}(G_{39},t)} & \boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)} & \boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)} \\ \boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)} & \boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)} & \boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)} \\ \boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)} & \boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)} & \boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)} \end{bmatrix} T_{15}$$

Where $\boxed{-(b''_{36})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{37})^{(7)}(G_{39},t)}$, $\boxed{-(b''_{38})^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{16})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{17})^{(2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b''_{18})^{(2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{20})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{21})^{(3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b''_{22})^{(3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{24})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{25})^{(4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b''_{26})^{(4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{28})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{29})^{(5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b''_{30})^{(5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{32})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{33})^{(6,6,6,6,6,6,6)}(G_{35},t)}$, $\boxed{-(b''_{34})^{(6,6,6,6,6,6,6)}(G_{35},t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{15})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{14})^{(1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b''_{13})^{(1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{40})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{41})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b''_{42})^{(8,8,8,8,8,8,8,8)}(G_{43},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b''_{46})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{45})^{(9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b''_{44})^{(9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{40}}{dt} = (a_{40})^{(8)}G_{41} - \begin{bmatrix} (a'_{40})^{(8)}\boxed{+(a''_{40})^{(8)}(T_{41},t)} & \boxed{+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{41}}{dt} = (a_{41})^{(8)}G_{40} - \begin{bmatrix} (a'_{41})^{(8)}\boxed{+(a''_{41})^{(8)}(T_{41},t)} & \boxed{+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{42}}{dt}$$
$$= (a_{42})^{(8)} G_{41}$$
$$- \left[\begin{array}{ccc} (a'_{42})^{(8)} & +(a''_{42})^{(8)}(T_{41},t) & +(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t) \\ +(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t) & +(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t) & +(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t) \\ +(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t) & +(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t) & +(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t) \\ +(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t) & & \end{array} \right] G_{15}$$

Where $+(a''_{40})^{(8)}(T_{41},t)$, $+(a''_{41})^{(8)}(T_{41},t)$, $+(a''_{42})^{(8)}(T_{41},t)$ are first augmentation coefficients for category 1, 2 and 3

$+(a''_{16})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{17})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$, $+(a''_{18})^{(2,2,2,2,2,2,2,2)}(T_{17},t)$ are second augmentation coefficient for category 1, 2 and 3

$+(a''_{20})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{21})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$, $+(a''_{22})^{(3,3,3,3,3,3,3,3)}(T_{21},t)$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{14})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ $+(a''_{15})^{(1,1,1,1,1,1,1,1)}(T_{14},t)$ are seventh augmentation coefficient for 1,2,3

$+(a''_{36})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{37})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$, $+(a''_{38})^{(7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{46})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{45})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$, $+(a''_{44})^{(9,9,9,9,9,9,9,9)}(T_{45},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{40}}{dt} =$$
$$(b_{40})^{(8)} T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{40})^{(8)} & -(b''_{40})^{(8)}(G_{43},t) & -(b''_{16})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{20})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{24})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{28})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{32})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{13})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{36})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{44})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{13}$$

$$\frac{dT_{41}}{dt} =$$
$$(b_{41})^{(8)} T_{40} -$$
$$\left[\begin{array}{ccc} (b'_{41})^{(8)} & -(b''_{41})^{(8)}(G_{43},t) & -(b''_{17})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{21})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{25})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{29})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{33})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{14})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{37})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{45})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{14}$$

$$\frac{dT_{42}}{dt} =$$
$$(b_{42})^{(8)} T_{41} -$$
$$\left[\begin{array}{ccc} (b'_{42})^{(8)} & -(b''_{42})^{(8)}(G_{43},t) & -(b''_{18})^{(2,2,2,2,2,2,2,2)}(G_{19},t) \\ -(b''_{22})^{(3,3,3,3,3,3,3,3)}(G_{23},t) & -(b''_{26})^{(4,4,4,4,4,4,4,4)}(G_{27},t) & -(b''_{30})^{(5,5,5,5,5,5,5,5)}(G_{31},t) \\ -(b''_{34})^{(6,6,6,6,6,6,6,6)}(G_{35},t) & -(b''_{15})^{(1,1,1,1,1,1,1,1)}(G,t) & -(b''_{38})^{(7,7,7,7,7,7,7,7)}(G_{39},t) \\ -(b''_{46})^{(9,9,9,9,9,9,9,9)}(G_{47},t) & & \end{array} \right] T_{15}$$

Where $\boxed{-(b_{36}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7)}(G_{39},t)}$ are first detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{16}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{17}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$, $\boxed{-(b_{18}'')^{(2,2,2,2,2,2,2,2)}(G_{19},t)}$ are second detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{20}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{21}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$, $\boxed{-(b_{22}'')^{(3,3,3,3,3,3,3,3)}(G_{23},t)}$ are third detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{24}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{25}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$, $\boxed{-(b_{26}'')^{(4,4,4,4,4,4,4,4)}(G_{27},t)}$ are fourth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{28}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{29}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$, $\boxed{-(b_{30}'')^{(5,5,5,5,5,5,5,5)}(G_{31},t)}$ are fifth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{32}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{33}'')^{(6,6,6,6,)}(G_{35},t)}$, $\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{38}'')^{(7,7,)}(G_{39},t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{44}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{45}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$, $\boxed{-(b_{46}'')^{(9,9,9,9,9,9,9,9)}(G_{47},t)}$ are ninth detrition coefficients for category 1, 2 and 3

$$\frac{dG_{44}}{dt} \qquad 96$$

$$= (a_{44})^{(9)}G_{45} - \begin{bmatrix} (a_{44}')^{(9)}\boxed{+(a_{44}'')^{(9)}(T_{45},t)} & \boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{24}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{28}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{32}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{13}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{36}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{40}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{13}$$

$$\frac{dG_{45}}{dt} = (a_{45})^{(9)}G_{44} - \begin{bmatrix} (a_{45}')^{(9)}\boxed{+(a_{45}'')^{(9)}(T_{45},t)} & \boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{25}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{29}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{33}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{14}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{37}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{41}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{14}$$

$$\frac{dG_{46}}{dt} = (a_{46})^{(9)}G_{45} - \begin{bmatrix} (a_{46}')^{(9)}\boxed{+(a_{46}'')^{(9)}(T_{37},t)} & \boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)} & \boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)} \\ \boxed{+(a_{26}'')^{(4,4,4,4,4,4,4,4)}(T_{25},t)} & \boxed{+(a_{30}'')^{(5,5,5,5,5,5,5,5)}(T_{29},t)} & \boxed{+(a_{34}'')^{(6,6,6,6,6,6,6,6)}(T_{33},t)} \\ \boxed{+(a_{15}'')^{(1,1,1,1,1,1,1,1)}(T_{14},t)} & \boxed{+(a_{38}'')^{(7,7,7,7,7,7,7,7)}(T_{37},t)} & \boxed{+(a_{42}'')^{(8,8,8,8,8,8,8,8)}(T_{41},t)} \end{bmatrix} G_{15}$$

Where $\boxed{+(a_{44}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{45}'')^{(9)}(T_{45},t)}$, $\boxed{+(a_{46}'')^{(9)}(T_{37},t)}$ are first augmentation coefficients for category 1, 2 and 3

$\boxed{+(a_{16}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{17}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$, $\boxed{+(a_{18}'')^{(2,2,2,2,2,2,2,2)}(T_{17},t)}$ are second augmentation coefficient for category 1, 2 and 3

$\boxed{+(a_{20}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{21}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$, $\boxed{+(a_{22}'')^{(3,3,3,3,3,3,3,3)}(T_{21},t)}$ are third augmentation coefficient for category 1, 2 and 3

$+(a''_{24})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{25})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$, $+(a''_{26})^{(4,4,4,4,4,4,4,4,4)}(T_{25},t)$ are fourth augmentation coefficient for category 1, 2 and 3

$+(a''_{28})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{29})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$, $+(a''_{30})^{(5,5,5,5,5,5,5,5,5)}(T_{29},t)$ are fifth augmentation coefficient for category 1, 2 and 3

$+(a''_{32})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{33})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$, $+(a''_{34})^{(6,6,6,6,6,6,6,6,6)}(T_{33},t)$ are sixth augmentation coefficient for category 1, 2 and 3

$+(a''_{13})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{14})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$, $+(a''_{15})^{(1,1,1,1,1,1,1,1,1)}(T_{14},t)$ are Seventh augmentation coefficient for category 1, 2 and 3

$+(a''_{38})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{37})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ $+(a''_{36})^{(7,7,7,7,7,7,7,7,7)}(T_{37},t)$ are eighth augmentation coefficient for 1,2,3

$+(a''_{40})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$,$+(a''_{42})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$, $+(a''_{41})^{(8,8,8,8,8,8,8,8,8)}(T_{41},t)$ are ninth augmentation coefficient for 1,2,3

$$\frac{dT_{44}}{dt} =$$
$$(b_{44})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{44})^{(9)}\left[-(b''_{44})^{(9)}(G_{47},t)\right] & \left[-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{13})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{36})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{40})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{13}$$

$$\frac{dT_{45}}{dt} =$$
$$(b_{45})^{(9)}T_{44} -$$
$$\begin{bmatrix} (b'_{45})^{(9)}\left[-(b''_{45})^{(9)}(G_{47},t)\right] & \left[-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{14})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{37})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{41})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{14}$$

$$\frac{dT_{46}}{dt} =$$
$$(b_{46})^{(9)}T_{45} -$$
$$\begin{bmatrix} (b'_{46})^{(9)}\left[-(b''_{46})^{(9)}(G_{47},t)\right] & \left[-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)\right] & \left[-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)\right] \\ \left[-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)\right] & \left[-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)\right] & \left[-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)\right] \\ \left[-(b''_{15})^{(1,1,1,1,1,1,1,1,1)}(G,t)\right] & \left[-(b''_{38})^{(7,7,7,7,7,7,7,7,7)}(G_{39},t)\right] & \left[-(b''_{42})^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)\right] \end{bmatrix} T_{15}$$

Where $-(b''_{44})^{(9)}(G_{47},t)$, $-(b''_{45})^{(9)}(G_{47},t)$, $-(b''_{46})^{(9)}(G_{47},t)$ are first detrition coefficients for category 1, 2 and 3

$-(b''_{16})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{17})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$, $-(b''_{18})^{(2,2,2,2,2,2,2,2,2)}(G_{19},t)$ are second detrition coefficients for category 1, 2 and 3

$-(b''_{20})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{21})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$, $-(b''_{22})^{(3,3,3,3,3,3,3,3,3)}(G_{23},t)$ are third detrition coefficients for category 1, 2 and 3

$-(b''_{24})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{25})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$, $-(b''_{26})^{(4,4,4,4,4,4,4,4,4)}(G_{27},t)$ are fourth detrition coefficients for category 1, 2 and 3

$-(b''_{28})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{29})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$, $-(b''_{30})^{(5,5,5,5,5,5,5,5,5)}(G_{31},t)$ are fifth detrition coefficients for category 1, 2 and 3

$-(b''_{32})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{33})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$, $-(b''_{34})^{(6,6,6,6,6,6,6,6,6)}(G_{35},t)$ are sixth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{15}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{14}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$, $\boxed{-(b_{13}'')^{(1,1,1,1,1,1,1,1)}(G,t)}$ are seventh detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{37}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{36}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$, $\boxed{-(b_{38}'')^{(7,7,7,7,7,7,7,7)}(G_{39},t)}$ are eighth detrition coefficients for category 1, 2 and 3

$\boxed{-(b_{42}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{41}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$, $\boxed{-(b_{40}'')^{(8,8,8,8,8,8,8,8,8)}(G_{43},t)}$ are ninth detrition coefficients for category 1, 2 and 3

Where we suppose

$$(a_i)^{(1)}, (a_i')^{(1)}, (a_i'')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (b_i'')^{(1)} > 0, \quad i,j = 13,14,15 \qquad 97$$

The functions $(a_i'')^{(1)}, (b_i'')^{(1)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(1)}$, $(r_i)^{(1)}$:

$$(a_i'')^{(1)}(T_{14},t) \le (p_i)^{(1)} \le (\hat{A}_{13})^{(1)}$$

$$(b_i'')^{(1)}(G,t) \le (r_i)^{(1)} \le (b_i')^{(1)} \le (\hat{B}_{13})^{(1)}$$

$$lim_{T_2\to\infty}(a_i'')^{(1)}(T_{14},t) = (p_i)^{(1)} \qquad 98$$

$$\lim_{G\to\infty}(b_i'')^{(1)}(G,t) = (r_i)^{(1)}$$

Definition of $(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}$:

Where $\boxed{(\hat{A}_{13})^{(1)}, (\hat{B}_{13})^{(1)}, (p_i)^{(1)}, (r_i)^{(1)}}$ are positive constants and $\boxed{i = 13,14,15}$

They satisfy Lipschitz condition: 99

$$|(a_i'')^{(1)}(T_{14}',t) - (a_i'')^{(1)}(T_{14},t)| \le (\hat{k}_{13})^{(1)}|T_{14} - T_{14}'|e^{-(\hat{M}_{13})^{(1)}t}$$

$$|(b_i'')^{(1)}(G',t) - (b_i'')^{(1)}(G,t)| < (\hat{k}_{13})^{(1)}||G - G'||e^{-(\hat{M}_{13})^{(1)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(1)}(T_{14}',t)$ and$(a_i'')^{(1)}(T_{14},t)$. (T_{14}',t) and (T_{14},t) are points belonging to the interval $[(\hat{k}_{13})^{(1)}, (\hat{M}_{13})^{(1)}]$. It is to be noted that $(a_i'')^{(1)}(T_{14},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{13})^{(1)} = 1$ then the function $(a_i'')^{(1)}(T_{14},t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$: 100

$(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}$, are positive constants

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$$

Definition of $(\hat{P}_{13})^{(1)}, (\hat{Q}_{13})^{(1)}$: 101

There exists two constants $(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$which together
With $(\hat{M}_{13})^{(1)}, (\hat{k}_{13})^{(1)}, (\hat{A}_{13})^{(1)}$ and $(\hat{B}_{13})^{(1)}$ and the constants
$(a_i)^{(1)}, (a_i')^{(1)}, (b_i)^{(1)}, (b_i')^{(1)}, (p_i)^{(1)}$, $(r_i)^{(1)}, i = 13,14,15,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(a_i)^{(1)} + (a_i')^{(1)} + (\hat{A}_{13})^{(1)} + (\hat{P}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

$$\frac{1}{(\hat{M}_{13})^{(1)}}[(b_i)^{(1)} + (b_i')^{(1)} + (\hat{B}_{13})^{(1)} + (\hat{Q}_{13})^{(1)}(\hat{k}_{13})^{(1)}] < 1$$

Where we suppose

$$(a_i)^{(2)}, (a_i')^{(2)}, (a_i'')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (b_i'')^{(2)} > 0, \quad i,j = 16,17,18$$

The functions $(a_i'')^{(2)}, (b_i'')^{(2)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(2)}$, $(r_i)^{(2)}$:

$$(a_i'')^{(2)}(T_{17}, t) \leq (p_i)^{(2)} \leq (\hat{A}_{16})^{(2)} \quad 102$$

$$(b_i'')^{(2)}(G_{19}, t) \leq (r_i)^{(2)} \leq (b_i')^{(2)} \leq (\hat{B}_{16})^{(2)} \quad 103$$

$$\lim_{T_2 \to \infty} (a_i'')^{(2)}(T_{17}, t) = (p_i)^{(2)} \quad 104$$

$$\lim_{G \to \infty} (b_i'')^{(2)}((G_{19}), t) = (r_i)^{(2)} \quad 105$$

Definition of $(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}$: 106

Where $\boxed{(\hat{A}_{16})^{(2)}, (\hat{B}_{16})^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}}$ are positive constants and $\boxed{i = 16,17,18}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(2)}(T_{17}', t) - (a_i'')^{(2)}(T_{17}, t)| \leq (\hat{k}_{16})^{(2)}|T_{17} - T_{17}'|e^{-(\hat{M}_{16})^{(2)}t} \quad 107$$

$$|(b_i'')^{(2)}((G_{19})', t) - (b_i'')^{(2)}((G_{19}), t)| < (\hat{k}_{16})^{(2)}||(G_{19}) - (G_{19})'||e^{-(\hat{M}_{16})^{(2)}t} \quad 108$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(2)}(T_{17}', t)$ and $(a_i'')^{(2)}(T_{17}, t)$. (T_{17}', t) and (T_{17}, t) are points belonging to the interval $[(\hat{k}_{16})^{(2)}, (\hat{M}_{16})^{(2)}]$. It is to be noted that $(a_i'')^{(2)}(T_{17}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{16})^{(2)} = 1$ then the function $(a_i'')^{(2)}(T_{17}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$:

$(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}$, are positive constants 109

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$$

Definition of $(\hat{P}_{13})^{(2)}, (\hat{Q}_{13})^{(2)}$:

There exists two constants $(\hat{P}_{16})^{(2)}$ and $(\hat{Q}_{16})^{(2)}$ which together with $(\hat{M}_{16})^{(2)}, (\hat{k}_{16})^{(2)}, (\hat{A}_{16})^{(2)} and (\hat{B}_{16})^{(2)}$ and the constants $(a_i)^{(2)}, (a_i')^{(2)}, (b_i)^{(2)}, (b_i')^{(2)}, (p_i)^{(2)}, (r_i)^{(2)}, i = 16,17,18,$

satisfy the inequalities

$$\frac{1}{(\hat{M}_{16})^{(2)}}[(a_i)^{(2)} + (a_i')^{(2)} + (\hat{A}_{16})^{(2)} + (\hat{P}_{16})^{(2)}(\hat{k}_{16})^{(2)}] < 1 \quad 110$$

$$\frac{1}{(\hat{M}_{16})^{(2)}}[\ (b_i)^{(2)} + (b_i')^{(2)} + \ (\hat{B}_{16})^{(2)} + (\hat{Q}_{16})^{(2)}\ (\hat{k}_{16})^{(2)}] < 1 \quad (111)$$

Where we suppose

$$(a_i)^{(3)}, (a_i')^{(3)}, (a_i'')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (b_i'')^{(3)} > 0, \quad i,j = 20,21,22 \quad (112)$$

The functions $(a_i'')^{(3)}, (b_i'')^{(3)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(3)}$, $(\mathrm{r_i})^{(3)}$:

$$(a_i'')^{(3)}(T_{21}, t) \leq (p_i)^{(3)} \leq (\hat{A}_{20})^{(3)}$$

$$(b_i'')^{(3)}(G_{23}, t) \leq \ (r_i)^{(3)} \leq (b_i')^{(3)} \leq (\hat{B}_{20})^{(3)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(3)}\ (T_{21}, t) = (p_i)^{(3)} \quad (113)$$

$$\lim_{G \to \infty}(b_i'')^{(3)}\ (G_{23}, t) = \ (r_i)^{(3)}$$

Definition of $(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}$:

Where $\boxed{(\hat{A}_{20})^{(3)}, (\hat{B}_{20})^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}}$ are positive constants and $\boxed{i = 20,21,22}$

They satisfy Lipschitz condition: (114)

$$|(a_i'')^{(3)}(T_{21}', t) - (a_i'')^{(3)}(T_{21}, t)| \leq (\hat{k}_{20})^{(3)}|T_{21} - T_{21}'|e^{-(\hat{M}_{20})^{(3)}t}$$

$$|(b_i'')^{(3)}(G_{23}', t) - (b_i'')^{(3)}(G_{23}, t)| < (\hat{k}_{20})^{(3)}||G_{23} - G_{23}'||e^{-(\hat{M}_{20})^{(3)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(3)}(T_{21}', t)$ and$(a_i'')^{(3)}(T_{21}, t)$. (T_{21}', t) And (T_{21}, t) are points belonging to the interval $\left[(\hat{k}_{20})^{(3)}, (\hat{M}_{20})^{(3)}\right]$. It is to be noted that $(a_i'')^{(3)}(T_{21}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{20})^{(3)} = 1$ then the function $(a_i'')^{(3)}(T_{21}, t)$, the first augmentation coefficient attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}$: (115)

$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)},$ are positive constants

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\ , \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$$

There exists two constants There exists two constants $(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ which together with (116)
$(\hat{M}_{20})^{(3)}, (\hat{k}_{20})^{(3)}, (\hat{A}_{20})^{(3)} and\ (\hat{B}_{20})^{(3)}$ and the constants
$(a_i)^{(3)}, (a_i')^{(3)}, (b_i)^{(3)}, (b_i')^{(3)}, (p_i)^{(3)}, \ (r_i)^{(3)}, i = 20,21,22,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (a_i)^{(3)} + (a_i')^{(3)} + \ (\hat{A}_{20})^{(3)} + (\hat{P}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

$$\frac{1}{(\hat{M}_{20})^{(3)}}[\ (b_i)^{(3)} + (b_i')^{(3)} + \ (\hat{B}_{20})^{(3)} + (\hat{Q}_{20})^{(3)}\ (\hat{k}_{20})^{(3)}] < 1$$

Where we suppose

$$(a_i)^{(4)}, (a_i')^{(4)}, (a_i'')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (b_i'')^{(4)} > 0, \quad i,j = 24,25,26 \quad (117)$$

The functions $(a_i'')^{(4)}, (b_i'')^{(4)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(4)}$, $(r_i)^{(4)}$:

$$(a_i'')^{(4)}(T_{25}, t) \leq (p_i)^{(4)} \leq (\hat{A}_{24})^{(4)}$$

$$(b_i'')^{(4)}\big((G_{27}), t\big) \leq (r_i)^{(4)} \leq (b_i')^{(4)} \leq (\hat{B}_{24})^{(4)}$$

$$lim_{T_2 \to \infty}(a_i'')^{(4)}(T_{25}, t) = (p_i)^{(4)} \quad (118)$$

$$\lim_{G \to \infty}(b_i'')^{(4)}\big((G_{27}), t\big) = (r_i)^{(4)}$$

Definition of $(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}$:

Where $\boxed{(\hat{A}_{24})^{(4)}, (\hat{B}_{24})^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}}$ are positive constants and $\boxed{i = 24,25,26}$

They satisfy Lipschitz condition: (119)

$$|(a_i'')^{(4)}(T_{25}', t) - (a_i'')^{(4)}(T_{25}, t)| \leq (\hat{k}_{24})^{(4)}|T_{25} - T_{25}'|e^{-(\hat{M}_{24})^{(4)}t}$$

$$|(b_i'')^{(4)}((G_{27})', t) - (b_i'')^{(4)}\big((G_{27}), t\big)| < (\hat{k}_{24})^{(4)}||(G_{27}) - (G_{27})'||e^{-(\hat{M}_{24})^{(4)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(4)}(T_{25}', t)$ and$(a_i'')^{(4)}(T_{25}, t)$. (T_{25}', t) and (T_{25}, t) are points belonging to the interval $\left[(\hat{k}_{24})^{(4)}, (\hat{M}_{24})^{(4)}\right]$. It is to be noted that $(a_i'')^{(4)}(T_{25}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{24})^{(4)} = 1$ then the function $(a_i'')^{(4)}(T_{25}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$: (120)

$(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}$, are positive constants

$$\frac{(a_i)^{(4)}}{(\hat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\hat{M}_{24})^{(4)}} < 1$$

Definition of $(\hat{P}_{24})^{(4)}, (\hat{Q}_{24})^{(4)}$: (121)

There exists two constants $(\hat{P}_{24})^{(4)}$ and $(\hat{Q}_{24})^{(4)}$ which together with $(\hat{M}_{24})^{(4)}, (\hat{k}_{24})^{(4)}, (\hat{A}_{24})^{(4)} and\ (\hat{B}_{24})^{(4)}$ and the constants $(a_i)^{(4)}, (a_i')^{(4)}, (b_i)^{(4)}, (b_i')^{(4)}, (p_i)^{(4)}, (r_i)^{(4)}, i = 24,25,26$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(a_i)^{(4)} + (a_i')^{(4)} + (\hat{A}_{24})^{(4)} + (\hat{P}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

$$\frac{1}{(\hat{M}_{24})^{(4)}}[(b_i)^{(4)} + (b_i')^{(4)} + (\hat{B}_{24})^{(4)} + (\hat{Q}_{24})^{(4)}(\hat{k}_{24})^{(4)}] < 1$$

Where we suppose

$$(a_i)^{(5)}, (a_i')^{(5)}, (a_i'')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (b_i'')^{(5)} > 0, \quad i, j = 28,29,30 \quad (122)$$

The functions $(a_i'')^{(5)}, (b_i'')^{(5)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(5)}$, $(r_i)^{(5)}$:

$$(a_i'')^{(5)}(T_{29},t) \le (p_i)^{(5)} \le (\hat{A}_{28})^{(5)}$$

$$(b_i'')^{(5)}((G_{31}),t) \le (r_i)^{(5)} \le (b_i')^{(5)} \le (\hat{B}_{28})^{(5)}$$

$$lim_{T_2\to\infty}(a_i'')^{(5)}(T_{29},t) = (p_i)^{(5)} \quad (123)$$

$$\lim_{G\to\infty}(b_i'')^{(5)}(G_{31},t) = (r_i)^{(5)}$$

Definition of $(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}$:

Where $\boxed{(\hat{A}_{28})^{(5)}, (\hat{B}_{28})^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}}$ are positive constants and $\boxed{i = 28,29,30}$

They satisfy Lipschitz condition: (124)

$$|(a_i'')^{(5)}(T_{29}',t) - (a_i'')^{(5)}(T_{29},t)| \le (\hat{k}_{28})^{(5)}|T_{29} - T_{29}'|e^{-(\hat{M}_{28})^{(5)}t}$$

$$|(b_i'')^{(5)}((G_{31})',t) - (b_i'')^{(5)}((G_{31}),t)| < (\hat{k}_{28})^{(5)}||(G_{31}) - (G_{31})'||e^{-(\hat{M}_{28})^{(5)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(5)}(T_{29}',t)$ and$(a_i'')^{(5)}(T_{29},t)$. (T_{29}',t) and (T_{29},t) are points belonging to the interval $\left[(\hat{k}_{28})^{(5)}, (\hat{M}_{28})^{(5)}\right]$. It is to be noted that $(a_i'')^{(5)}(T_{29},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{28})^{(5)} = 1$ then the function $(a_i'')^{(5)}(T_{29},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$: (125)

$(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}$, are positive constants

$$\frac{(a_i)^{(5)}}{(\hat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\hat{M}_{28})^{(5)}} < 1$$

Definition of $(\hat{P}_{28})^{(5)}, (\hat{Q}_{28})^{(5)}$: (126)

There exists two constants $(\hat{P}_{28})^{(5)}$ and $(\hat{Q}_{28})^{(5)}$ which together with $(\hat{M}_{28})^{(5)}, (\hat{k}_{28})^{(5)}, (\hat{A}_{28})^{(5)} and\ (\hat{B}_{28})^{(5)}$ and the constants $(a_i)^{(5)}, (a_i')^{(5)}, (b_i)^{(5)}, (b_i')^{(5)}, (p_i)^{(5)}, (r_i)^{(5)}, i = 28,29,30,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(a_i)^{(5)} + (a_i')^{(5)} + (\hat{A}_{28})^{(5)} + (\hat{P}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

$$\frac{1}{(\hat{M}_{28})^{(5)}}[(b_i)^{(5)} + (b_i')^{(5)} + (\hat{B}_{28})^{(5)} + (\hat{Q}_{28})^{(5)}(\hat{k}_{28})^{(5)}] < 1$$

Where we suppose

$$(a_i)^{(6)}, (a_i')^{(6)}, (a_i'')^{(6)}, (b_i)^{(6)}, (b_i')^{(6)}, (b_i'')^{(6)} > 0, \quad i,j = 32,33,34 \quad (127)$$

The functions $(a_i'')^{(6)}, (b_i'')^{(6)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(6)}$, $(r_i)^{(6)}$:

$$(a_i'')^{(6)}(T_{33},t) \le (p_i)^{(6)} \le (\hat{A}_{32})^{(6)}$$

$$(b_i'')^{(6)}((G_{35}),t) \le (r_i)^{(6)} \le (b_i')^{(6)} \le (\hat{B}_{32})^{(6)}$$

$$lim_{T_2\to\infty}(a_i'')^{(6)}\,(T_{33},t)=(p_i)^{(6)} \qquad 128$$

$$\lim_{G\to\infty}(b_i'')^{(6)}\,\big((G_{35}),t\big)=\ (r_i)^{(6)}$$

Definition of $(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)}$:

Where $\boxed{(\hat{A}_{32})^{(6)},(\hat{B}_{32})^{(6)},(p_i)^{(6)},\ (r_i)^{(6)}}$ are positive constants and $\boxed{i=32,33,34}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(6)}(T_{33}',t)-(a_i'')^{(6)}(T_{33},t)|\le(\hat{k}_{32})^{(6)}|T_{33}-T_{33}'|e^{-(\hat{M}_{32})^{(6)}t}$$

$$|(b_i'')^{(6)}((G_{35})',t)-(b_i'')^{(6)}\big((G_{35}),t\big)|<(\hat{k}_{32})^{(6)}||(G_{35})-(G_{35})'||e^{-(\hat{M}_{32})^{(6)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(6)}(T_{33}',t)$ and$(a_i'')^{(6)}(T_{33},t)$. (T_{33}',t) and (T_{33},t) are points belonging to the interval $\big[(\hat{k}_{32})^{(6)},(\hat{M}_{32})^{(6)}\big]$. It is to be noted that $(a_i'')^{(6)}(T_{33},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{32})^{(6)}=1$ then the function $(a_i'')^{(6)}(T_{33},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$: 129

$(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)}$, are positive constants

$$\frac{(a_i)^{(6)}}{(\hat{M}_{32})^{(6)}}\ ,\frac{(b_i)^{(6)}}{(\hat{M}_{32})^{(6)}}<1$$

Definition of $(\hat{P}_{32})^{(6)},(\hat{Q}_{32})^{(6)}$: 130

There exists two constants $(\hat{P}_{32})^{(6)}$ and $(\hat{Q}_{32})^{(6)}$ which together with $(\hat{M}_{32})^{(6)},(\hat{k}_{32})^{(6)},(\hat{A}_{32})^{(6)}and\ (\hat{B}_{32})^{(6)}$ and the constants $(a_i)^{(6)},(a_i')^{(6)},(b_i)^{(6)},(b_i')^{(6)},(p_i)^{(6)},\ (r_i)^{(6)},i=32,33,34,$ satisfy the inequalities

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\,(a_i)^{(6)}+(a_i')^{(6)}+\ (\hat{A}_{32})^{(6)}+\ (\hat{P}_{32})^{(6)}\,(\hat{k}_{32})^{(6)}]<1$$

$$\frac{1}{(\hat{M}_{32})^{(6)}}[\ (b_i)^{(6)}+(b_i')^{(6)}+\ (\hat{B}_{32})^{(6)}+\ (\hat{Q}_{32})^{(6)}\ (\hat{k}_{32})^{(6)}]<1$$

Where we suppose

M) $(a_i)^{(7)},(a_i')^{(7)},(a_i'')^{(7)},(b_i)^{(7)},(b_i')^{(7)},(b_i'')^{(7)}>0,\quad i,j=36,37,38$ 131

)The functions $(a_i'')^{(7)},(b_i'')^{(7)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(7)},\ (r_i)^{(7)}$:

$$(a_i'')^{(7)}(T_{37},t)\le(p_i)^{(7)}\le(\hat{A}_{36})^{(7)}$$

$$(b_i'')^{(7)}(G_{39},t)\le\ (r_i)^{(7)}\le(b_i')^{(7)}\le(\hat{B}_{36})^{(7)}$$

132

)$\lim_{T_2\to\infty}\ (a_i'')^{(7)}\,(T_{37},t)=(p_i)^{(7)}$

$\lim_{G\to\infty}(b_i'')^{(7)}\left((G_{39}),t\right) = \ (r_i)^{(7)}$

Definition of $(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}$:

Where $\boxed{(\hat{A}_{36})^{(7)}, (\hat{B}_{36})^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}}$ are positive constants and $\boxed{i = 36{,}37{,}38}$

They satisfy Lipschitz condition: 133

$$|(a_i'')^{(7)}(T_{37}', t) - (a_i'')^{(7)}(T_{37}, t)| \leq (\hat{k}_{36})^{(7)}|T_{37} - T_{37}'|e^{-(\hat{M}_{36})^{(7)}t}$$

$$|(b_i'')^{(7)}((G_{39})', t) - (b_i'')^{(7)}\left((G_{39}),t\right)| < (\hat{k}_{36})^{(7)}||(G_{39}) - (G_{39})'||e^{-(\hat{M}_{36})^{(7)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(7)}(T_{37}', t)$ and$(a_i'')^{(7)}(T_{37}, t)$. (T_{37}', t) and (T_{37}, t) are points belonging to the interval $\left[(\hat{k}_{36})^{(7)}, (\hat{M}_{36})^{(7)}\right]$. It is to be noted that $(a_i'')^{(7)}(T_{37}, t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{36})^{(7)} = 1$ then the function $(a_i'')^{(7)}(T_{37}, t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$: 134

$(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}$, are positive constants

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}, \frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$$

Definition of $(\hat{P}_{36})^{(7)}, (\hat{Q}_{36})^{(7)}$: 135

There exists two constants $(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ which together with $(\hat{M}_{36})^{(7)}, (\hat{k}_{36})^{(7)}, (\hat{A}_{36})^{(7)} and\ (\hat{B}_{36})^{(7)}$ and the constants $(a_i)^{(7)}, (a_i')^{(7)}, (b_i)^{(7)}, (b_i')^{(7)}, (p_i)^{(7)}, \ (r_i)^{(7)}, i = 36{,}37{,}38$, satisfy the inequalities

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\,(a_i)^{(7)} + (a_i')^{(7)} + \ (\hat{A}_{36})^{(7)} + \ (\hat{P}_{36})^{(7)}\,(\hat{k}_{36})^{(7)}] < 1$$

$$\frac{1}{(\hat{M}_{36})^{(7)}}[\ (b_i)^{(7)} + (b_i')^{(7)} + \ (\hat{B}_{36})^{(7)} + \ (\hat{Q}_{36})^{(7)}\ (\hat{k}_{36})^{(7)}] < 1$$

Where we suppose

$$(a_i)^{(8)}, (a_i')^{(8)}, (a_i'')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (b_i'')^{(8)} > 0, \quad i, j = 40{,}41{,}42 \qquad 136$$

The functions $(a_i'')^{(8)}, (b_i'')^{(8)}$ are positive continuous increasing and bounded

Definition of $(p_i)^{(8)}, \ (r_i)^{(8)}$: 137

$$(a_i'')^{(8)}(T_{41}, t) \leq (p_i)^{(8)} \leq (\hat{A}_{40})^{(8)} \qquad 138$$

$$(b_i'')^{(8)}((G_{43}), t) \leq \ (r_i)^{(8)} \leq (b_i')^{(8)} \leq (\hat{B}_{40})^{(8)} \qquad 139$$

$\lim_{T_2\to\infty}(a_i'')^{(8)}(T_{41},t) = (p_i)^{(8)}$ 140

$\lim_{G\to\infty}(b_i'')^{(8)}\left((G_{43}),t\right) = (r_i)^{(8)}$ 141

Definition of $(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}$:

Where $\boxed{(\hat{A}_{40})^{(8)}, (\hat{B}_{40})^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}}$ are positive constants and $\boxed{i = 40,41,42}$

They satisfy Lipschitz condition:

$|(a_i'')^{(8)}(T_{41}',t) - (a_i'')^{(8)}(T_{41},t)| \leq (\hat{k}_{40})^{(8)}|T_{41} - T_{41}'|e^{-(\hat{M}_{40})^{(8)}t}$ 142

$|(b_i'')^{(8)}((G_{43})',t) - (b_i'')^{(8)}\left((G_{43}),t\right)| < (\hat{k}_{40})^{(8)}||(G_{43}) - (G_{43})'||e^{-(\hat{M}_{40})^{(8)}t}$ 143

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(8)}(T_{41}',t)$ and $(a_i'')^{(8)}(T_{41},t)$. (T_{41}',t) and (T_{41},t) are points belonging to the interval $\left[(\hat{k}_{40})^{(8)}, (\hat{M}_{40})^{(8)}\right]$. It is to be noted that $(a_i'')^{(8)}(T_{41},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{40})^{(8)} = 1$ then the function $(a_i'')^{(8)}(T_{41},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$:

$(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}$, are positive constants

$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ 144

Definition of $(\hat{P}_{40})^{(8)}, (\hat{Q}_{40})^{(8)}$:

There exists two constants $(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ which together with $(\hat{M}_{40})^{(8)}, (\hat{k}_{40})^{(8)}, (\hat{A}_{40})^{(8)}$ $(\hat{B}_{40})^{(8)}$ and the constants $(a_i)^{(8)}, (a_i')^{(8)}, (b_i)^{(8)}, (b_i')^{(8)}, (p_i)^{(8)}, (r_i)^{(8)}, i = 40,41,42,$
Satisfy the inequalities

$\frac{1}{(\hat{M}_{40})^{(8)}}[(a_i)^{(8)} + (a_i')^{(8)} + (\hat{A}_{40})^{(8)} + (\hat{P}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 145

$\frac{1}{(\hat{M}_{40})^{(8)}}[(b_i)^{(8)} + (b_i')^{(8)} + (\hat{B}_{40})^{(8)} + (\hat{Q}_{40})^{(8)}(\hat{k}_{40})^{(8)}] < 1$ 146

Where we suppose

$(a_i)^{(9)}, (a_i')^{(9)}, (a_i'')^{(9)}, (b_i)^{(9)}, (b_i')^{(9)}, (b_i'')^{(9)} > 0, \quad i,j = 44,45,46$ 146 A

The functions $(a_i'')^{(9)}, (b_i'')^{(9)}$ are positive continuous increasing and bounded.

Definition of $(p_i)^{(9)}$, $(r_i)^{(9)}$:

$(a_i'')^{(9)}(T_{45},t) \leq (p_i)^{(9)} \leq (\hat{A}_{44})^{(9)}$

$(b_i'')^{(9)}(G_{47},t) \leq (r_i)^{(9)} \leq (b_i')^{(9)} \leq (\hat{B}_{44})^{(9)}$

$\lim_{T_2\to\infty}(a_i'')^{(9)}(T_{45},t)=(p_i)^{(9)}$

$\lim_{G\to\infty}(b_i'')^{(9)}(G_{47},t)=\ (r_i)^{(9)}$

Definition of $(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)}$:

Where $\boxed{(\hat{A}_{44})^{(9)},(\hat{B}_{44})^{(9)},(p_i)^{(9)},\ (r_i)^{(9)}}$ are positive constants and $\boxed{i=44{,}45{,}46}$

They satisfy Lipschitz condition:

$$|(a_i'')^{(9)}(T_{45}',t)-(a_i'')^{(9)}(T_{45},t)|\le(\hat{k}_{44})^{(9)}|T_{45}-T_{45}'|e^{-(\hat{M}_{44})^{(9)}t}$$

$$|(b_i'')^{(9)}((G_{47})',t)-(b_i'')^{(9)}\big((G_{47}),t\big)|<(\hat{k}_{44})^{(9)}||(G_{47})-(G_{47})'||e^{-(\hat{M}_{44})^{(9)}t}$$

With the Lipschitz condition, we place a restriction on the behavior of functions $(a_i'')^{(9)}(T_{45}',t)$ and$(a_i'')^{(9)}(T_{45},t)$. (T_{45}',t) and (T_{45},t) are points belonging to the interval $\left[(\hat{k}_{44})^{(9)},(\hat{M}_{44})^{(9)}\right]$. It is to be noted that $(a_i'')^{(9)}(T_{45},t)$ is uniformly continuous. In the eventuality of the fact, that if $(\hat{M}_{44})^{(9)}=1$ then the function $(a_i'')^{(9)}(T_{45},t)$, the **first augmentation coefficient** attributable to the system, would be absolutely continuous.

Definition of $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)}$:

$(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},$ are positive constants

$$\frac{(a_i)^{(9)}}{(\hat{M}_{44})^{(9)}}\ ,\frac{(b_i)^{(9)}}{(\hat{M}_{44})^{(9)}}<1$$

Definition of $(\hat{P}_{44})^{(9)},(\hat{Q}_{44})^{(9)}$:

There exists two constants $(\hat{P}_{44})^{(9)}$ and $(\hat{Q}_{44})^{(9)}$ which together with $(\hat{M}_{44})^{(9)},(\hat{k}_{44})^{(9)},(\hat{A}_{44})^{(9)}and\ (\hat{B}_{44})^{(9)}$ and the constants $(a_i)^{(9)},(a_i')^{(9)},(b_i)^{(9)},(b_i')^{(9)},(p_i)^{(9)},\ (r_i)^{(9)},i=44{,}45{,}46,$
satisfy the inequalities

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\,(a_i)^{(9)}+(a_i')^{(9)}+\ (\hat{A}_{44})^{(9)}+(\hat{P}_{44})^{(9)}\,(\hat{k}_{44})^{(9)}]<1$$

$$\frac{1}{(\hat{M}_{44})^{(9)}}[\ (b_i)^{(9)}+(b_i')^{(9)}+\ (\hat{B}_{44})^{(9)}+(\hat{Q}_{44})^{(9)}\ (\hat{k}_{44})^{(9)}]<1$$

Theorem 1: if the conditions above are fulfilled, there exists a solution satisfying the conditions 147

Definition of $G_i(0)\,,T_i(0)$:

$$G_i(t)\le\left(\hat{P}_{13}\right)^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{G_i(0)=G_i^0>0}$$

$$T_i(t)\le(\hat{Q}_{13})^{(1)}e^{(\hat{M}_{13})^{(1)}t}\ ,\ \boxed{T_i(0)=T_i^0>0}$$

Theorem 2 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 148

Definition of $G_i(0)\,, T_i(0)$

$$G_i(t) \le (\hat{P}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{16})^{(2)} e^{(\hat{M}_{16})^{(2)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 3 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 149

$$G_i(t) \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad G_i(0) = G_i^0 > 0$$

$$T_i(t) \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t} \quad , \quad T_i(0) = T_i^0 > 0$$

Theorem 4 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 150

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{24}\right)^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 5 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 151

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{28}\right)^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 6 : if the conditions above are fulfilled, there exists a solution satisfying the conditions 152

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{32}\right)^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 7: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{36}\right)^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 8: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153 A

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{40}\right)^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Theorem 9: if the conditions above are fulfilled, there exists a solution satisfying the conditions 153

B

Definition of $G_i(0)\,, T_i(0)$:

$$G_i(t) \le \left(\hat{P}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{G_i(0) = G_i^0 > 0}$$

$$T_i(t) \le \left(\hat{Q}_{44}\right)^{(9)} e^{(\hat{M}_{44})^{(9)}t} \quad , \quad \boxed{T_i(0) = T_i^0 > 0}$$

Proof: Consider operator $\mathcal{A}^{(1)}$ defined on the space of sextuples of continuous functions 154
$G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{13}\right)^{(1)}, T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)}, \qquad 155$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 156$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{13}\right)^{(1)} e^{(\hat{M}_{13})^{(1)}t} \qquad 157$$

By 158

$$\bar{G}_{13}(t) = G_{13}^0 + \int_0^t \left[(a_{13})^{(1)} G_{14}(s_{(13)}) - \left((a'_{13})^{(1)} + a''_{13})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{14}(t) = G_{14}^0 + \int_0^t \left[(a_{14})^{(1)} G_{13}(s_{(13)}) - \left((a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{G}_{15}(t) = G_{15}^0 + \int_0^t \left[(a_{15})^{(1)} G_{14}(s_{(13)}) - \left((a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14}(s_{(13)}), s_{(13)})\right) G_{15}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{13}(t) = T_{13}^0 + \int_0^t \left[(b_{13})^{(1)} T_{14}(s_{(13)}) - \left((b'_{13})^{(1)} - (b''_{13})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{13}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{14}(t) = T_{14}^0 + \int_0^t \left[(b_{14})^{(1)} T_{13}(s_{(13)}) - \left((b'_{14})^{(1)} - (b''_{14})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{14}(s_{(13)})\right] ds_{(13)}$$

$$\bar{T}_{15}(t) = T_{15}^0 + \int_0^t \left[(b_{15})^{(1)} T_{14}(s_{(13)}) - \left((b'_{15})^{(1)} - (b''_{15})^{(1)}(G(s_{(13)}), s_{(13)})\right) T_{15}(s_{(13)})\right] ds_{(13)}$$

Where $s_{(13)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 159

Consider operator $\mathcal{A}^{(2)}$ defined on the space of sextuples of continuous functions $G_i\,,\ T_i\colon \mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0\,,\ T_i(0) = T_i^0\,,\ G_i^0 \le \left(\hat{P}_{16}\right)^{(2)}, T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)},$$

$$0 \le G_i(t) - G_i^0 \le \left(\hat{P}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

$$0 \le T_i(t) - T_i^0 \le \left(\hat{Q}_{16}\right)^{(2)} e^{(\hat{M}_{16})^{(2)}t}$$

By 160

$$\bar{G}_{16}(t) = G_{16}^0 + \int_0^t \left[(a_{16})^{(2)} G_{17}(s_{(16)}) - \left((a'_{16})^{(2)} + a''_{16})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{17}(t) = G_{17}^0 + \int_0^t \left[(a_{17})^{(2)} G_{16}(s_{(16)}) - \left((a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17}(s_{(16)}), s_{(17)})\right) G_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{G}_{18}(t) = G_{18}^0 + \int_0^t \left[(a_{18})^{(2)} G_{17}(s_{(16)}) - \left((a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17}(s_{(16)}), s_{(16)})\right) G_{18}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{16}(t) = T_{16}^0 + \int_0^t \left[(b_{16})^{(2)} T_{17}(s_{(16)}) - \left((b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{16}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{17}(t) = T_{17}^0 + \int_0^t \left[(b_{17})^{(2)} T_{16}(s_{(16)}) - \left((b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{17}(s_{(16)})\right] ds_{(16)}$$

$$\bar{T}_{18}(t) = T_{18}^0 + \int_0^t \left[(b_{18})^{(2)} T_{17}(s_{(16)}) - \left((b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19}(s_{(16)}), s_{(16)})\right) T_{18}(s_{(16)})\right] ds_{(16)}$$

Where $s_{(16)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(3)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{20})^{(3)}$, $T_i^0 \le (\hat{Q}_{20})^{(3)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} t}$

By 161

$$\bar{G}_{20}(t) = G_{20}^0 + \int_0^t \left[(a_{20})^{(3)} G_{21}(s_{(20)}) - \left((a'_{20})^{(3)} + a''_{20})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{21}(t) = G_{21}^0 + \int_0^t \left[(a_{21})^{(3)} G_{20}(s_{(20)}) - \left((a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{G}_{22}(t) = G_{22}^0 + \int_0^t \left[(a_{22})^{(3)} G_{21}(s_{(20)}) - \left((a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21}(s_{(20)}), s_{(20)})\right) G_{22}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{20}(t) = T_{20}^0 + \int_0^t \left[(b_{20})^{(3)} T_{21}(s_{(20)}) - \left((b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{20}(s_{(20)})\right] ds_{(20)}$$

$$\bar{T}_{21}(t) = T_{21}^0 + \int_0^t \left[(b_{21})^{(3)} T_{20}(s_{(20)}) - \left((b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{21}(s_{(20)})\right] ds_{(20)}$$

$$\bar{\mathrm{T}}_{22}(\mathrm{t}) = \mathrm{T}_{22}^0 + \int_0^t \left[(b_{22})^{(3)} T_{21}(s_{(20)}) - \left((b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23}(s_{(20)}), s_{(20)})\right) T_{22}(s_{(20)})\right] ds_{(20)}$$

Where $s_{(20)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(4)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{24})^{(4)}$, $T_i^0 \le (\hat{Q}_{24})^{(4)}$,

$0 \le G_i(t) - G_i^0 \le (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} t}$

By 162

$$\bar{G}_{24}(t) = G_{24}^0 + \int_0^t \left[(a_{24})^{(4)} G_{25}(s_{(24)}) - \left((a'_{24})^{(4)} + a''_{24})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{25}(t) = G_{25}^0 + \int_0^t \left[(a_{25})^{(4)} G_{24}(s_{(24)}) - \left((a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{G}_{26}(t) = G_{26}^0 + \int_0^t \left[(a_{26})^{(4)} G_{25}(s_{(24)}) - \left((a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25}(s_{(24)}), s_{(24)})\right) G_{26}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{24}(t) = T_{24}^0 + \int_0^t \left[(b_{24})^{(4)} T_{25}(s_{(24)}) - \left((b'_{24})^{(4)} - (b''_{24})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{24}(s_{(24)})\right] ds_{(24)}$$

$$\bar{T}_{25}(t) = T_{25}^0 + \int_0^t \left[(b_{25})^{(4)} T_{24}(s_{(24)}) - \left((b'_{25})^{(4)} - (b''_{25})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{25}(s_{(24)})\right] ds_{(24)}$$

$$\bar{\mathrm{T}}_{26}(\mathrm{t}) = \mathrm{T}_{26}^0 + \int_0^t \left[(b_{26})^{(4)} T_{25}(s_{(24)}) - \left((b'_{26})^{(4)} - (b''_{26})^{(4)}(G_{27}(s_{(24)}), s_{(24)})\right) T_{26}(s_{(24)})\right] ds_{(24)}$$

Where $s_{(24)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: Consider operator $\mathcal{A}^{(5)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{28})^{(5)} \text{ , } T_i^0 \le (\hat{Q}_{28})^{(5)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{28})^{(5)} e^{(\hat{M}_{28})^{(5)} t}$$

By 163

$$\bar{G}_{28}(t) = G_{28}^0 + \int_0^t \left[(a_{28})^{(5)} G_{29}(s_{(28)}) - \left((a'_{28})^{(5)} + a''_{28})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{29}(t) = G_{29}^0 + \int_0^t \left[(a_{29})^{(5)} G_{28}(s_{(28)}) - \left((a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{G}_{30}(t) = G_{30}^0 + \int_0^t \left[(a_{30})^{(5)} G_{29}(s_{(28)}) - \left((a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29}(s_{(28)}), s_{(28)})\right) G_{30}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{28}(t) = T_{28}^0 + \int_0^t \left[(b_{28})^{(5)} T_{29}(s_{(28)}) - \left((b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{28}(s_{(28)})\right] ds_{(28)}$$

$$\bar{T}_{29}(t) = T_{29}^0 + \int_0^t \left[(b_{29})^{(5)} T_{28}(s_{(28)}) - \left((b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{29}(s_{(28)})\right] ds_{(28)}$$

$$\bar{\mathrm{T}}_{30}(\mathrm{t}) = \mathrm{T}_{30}^0 + \int_0^t \left[(b_{30})^{(5)} T_{29}(s_{(28)}) - \left((b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31}(s_{(28)}), s_{(28)})\right) T_{30}(s_{(28)})\right] ds_{(28)}$$

Where $s_{(28)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(6)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$$G_i(0) = G_i^0 \text{ , } T_i(0) = T_i^0 \text{ , } G_i^0 \le (\hat{P}_{32})^{(6)} \text{ , } T_i^0 \le (\hat{Q}_{32})^{(6)},$$

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{32})^{(6)} e^{(\hat{M}_{32})^{(6)} t}$$

By 164

$$\bar{G}_{32}(t) = G_{32}^0 + \int_0^t \left[(a_{32})^{(6)} G_{33}(s_{(32)}) - \left((a'_{32})^{(6)} + a''_{32})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{33}(t) = G_{33}^0 + \int_0^t \left[(a_{33})^{(6)} G_{32}(s_{(32)}) - \left((a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{G}_{34}(t) = G_{34}^0 + \int_0^t \left[(a_{34})^{(6)} G_{33}(s_{(32)}) - \left((a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33}(s_{(32)}), s_{(32)})\right) G_{34}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{32}(t) = T_{32}^0 + \int_0^t \left[(b_{32})^{(6)} T_{33}(s_{(32)}) - \left((b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{32}(s_{(32)})\right] ds_{(32)}$$

$$\bar{T}_{33}(t) = T_{33}^0 + \int_0^t \left[(b_{33})^{(6)} T_{32}(s_{(32)}) - \left((b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{33}(s_{(32)})\right] ds_{(32)}$$

$$\bar{\mathrm{T}}_{34}(\mathrm{t}) = \mathrm{T}_{34}^0 + \int_0^t \left[(b_{34})^{(6)} T_{33}(s_{(32)}) - \left((b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35}(s_{(32)}), s_{(32)})\right) T_{34}(s_{(32)})\right] ds_{(32)}$$

Where $s_{(32)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(7)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$ which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \le (\hat{P}_{36})^{(7)}$, $T_i^0 \le (\hat{Q}_{36})^{(7)}$,

$$0 \le G_i(t) - G_i^0 \le (\hat{P}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

$$0 \le T_i(t) - T_i^0 \le (\hat{Q}_{36})^{(7)} e^{(\hat{M}_{36})^{(7)} t}$$

By 165

$$\bar{G}_{36}(t) = G_{36}^0 + \int_0^t \left[(a_{36})^{(7)} G_{37}(s_{(36)}) - \left((a'_{36})^{(7)} + a''_{36})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{37}(t) = G_{37}^0 + \int_0^t \left[(a_{37})^{(7)} G_{36}(s_{(36)}) - \left((a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{G}_{38}(t) = G_{38}^0 + \int_0^t \left[(a_{38})^{(7)} G_{37}(s_{(36)}) - \left((a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37}(s_{(36)}), s_{(36)})\right) G_{38}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{36}(t) = T_{36}^0 + \int_0^t \left[(b_{36})^{(7)} T_{37}(s_{(36)}) - \left((b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{36}(s_{(36)})\right] ds_{(36)}$$

$$\bar{T}_{37}(t) = T_{37}^0 + \int_0^t \left[(b_{37})^{(7)} T_{36}(s_{(36)}) - \left((b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{37}(s_{(36)})\right] ds_{(36)}$$

$$\bar{\mathrm{T}}_{38}(\mathrm{t}) = \mathrm{T}_{38}^0 + \int_0^t \left[(b_{38})^{(7)} T_{37}(s_{(36)}) - \left((b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39}(s_{(36)}), s_{(36)})\right) T_{38}(s_{(36)})\right] ds_{(36)}$$

Where $s_{(36)}$ is the integrand that is integrated over an interval $(0, t)$

Proof:

Consider operator $\mathcal{A}^{(8)}$ defined on the space of sextuples of continuous functions G_i , T_i: $\mathbb{R}_+ \to \mathbb{R}_+$

which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{40})^{(8)}$, $T_i^0 \leq (\hat{Q}_{40})^{(8)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} t}$

By 166

$$\bar{G}_{40}(t) = G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} G_{41}(s_{(40)}) - \left((a'_{40})^{(8)} + a''_{40})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{41}(t) = G_{41}^0 + \int_0^t \left[(a_{41})^{(8)} G_{40}(s_{(40)}) - \left((a'_{41})^{(8)} + (a''_{41})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{G}_{42}(t) = G_{42}^0 + \int_0^t \left[(a_{42})^{(8)} G_{41}(s_{(40)}) - \left((a'_{42})^{(8)} + (a''_{42})^{(8)} (T_{41}(s_{(40)}), s_{(40)})\right) G_{42}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{40}(t) = T_{40}^0 + \int_0^t \left[(b_{40})^{(8)} T_{41}(s_{(40)}) - \left((b'_{40})^{(8)} - (b''_{40})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{40}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{41}(t) = T_{41}^0 + \int_0^t \left[(b_{41})^{(8)} T_{40}(s_{(40)}) - \left((b'_{41})^{(8)} - (b''_{41})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{41}(s_{(40)})\right] ds_{(40)}$$

$$\bar{T}_{42}(t) = T_{42}^0 + \int_0^t \left[(b_{42})^{(8)} T_{41}(s_{(40)}) - \left((b'_{42})^{(8)} - (b''_{42})^{(8)} (G_{43}(s_{(40)}), s_{(40)})\right) T_{42}(s_{(40)})\right] ds_{(40)}$$

Where $s_{(40)}$ is the integrand that is integrated over an interval $(0, t)$

Proof: 166

Consider operator $\mathcal{A}^{(9)}$ defined on the space of sextuples of continuous functions G_i , $T_i: \mathbb{R}_+ \to \mathbb{R}_+$ A which satisfy

$G_i(0) = G_i^0$, $T_i(0) = T_i^0$, $G_i^0 \leq (\hat{P}_{44})^{(9)}$, $T_i^0 \leq (\hat{Q}_{44})^{(9)}$,

$0 \leq G_i(t) - G_i^0 \leq (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

$0 \leq T_i(t) - T_i^0 \leq (\hat{Q}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} t}$

By

$$\bar{G}_{44}(t) = G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} G_{45}(s_{(44)}) - \left((a'_{44})^{(9)} + a''_{44})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{45}(t) = G_{45}^0 + \int_0^t \left[(a_{45})^{(9)} G_{44}(s_{(44)}) - \left((a'_{45})^{(9)} + (a''_{45})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{45}(s_{(44)})\right] ds_{(44)}$$

$$\bar{G}_{46}(t) = G_{46}^0 + \int_0^t \left[(a_{46})^{(9)} G_{45}(s_{(44)}) - \left((a'_{46})^{(9)} + (a''_{46})^{(9)} (T_{45}(s_{(44)}), s_{(44)})\right) G_{46}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{44}(t) = T_{44}^0 + \int_0^t \left[(b_{44})^{(9)} T_{45}(s_{(44)}) - \left((b'_{44})^{(9)} - (b''_{44})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{44}(s_{(44)})\right] ds_{(44)}$$

$$\bar{T}_{45}(t) = T_{45}^0 + \int_0^t \left[(b_{45})^{(9)} T_{44}(s_{(44)}) - \left((b'_{45})^{(9)} - (b''_{45})^{(9)} (G_{47}(s_{(44)}), s_{(44)})\right) T_{45}(s_{(44)})\right] ds_{(44)}$$

$$\overline{\mathrm{T}}_{46}(\mathrm{t}) = \mathrm{T}_{46}^{0} + \int_0^t \left[(b_{46})^{(9)} T_{45}(s_{(44)}) - \left((b'_{46})^{(9)} - (b''_{46})^{(9)}\left(G_{47}(s_{(44)}), s_{(44)}\right)\right) T_{46}(s_{(44)})\right] ds_{(44)}$$

Where $s_{(44)}$ is the integrand that is integrated over an interval $(0, t)$

The operator $\mathcal{A}^{(1)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 167

$$G_{13}(t) \le G_{13}^{0} + \int_0^t \left[(a_{13})^{(1)}\left(G_{14}^{0} + (\hat{P}_{13})^{(1)} e^{(\hat{M}_{13})^{(1)} s_{(13)}}\right)\right] ds_{(13)} =$$

$$\left(1 + (a_{13})^{(1)} t\right) G_{14}^{0} + \frac{(a_{13})^{(1)}(\hat{P}_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left(e^{(\hat{M}_{13})^{(1)} t} - 1\right)$$

From which it follows that 168

$$(G_{13}(t) - G_{13}^{0}) e^{-(\hat{M}_{13})^{(1)} t} \le \frac{(a_{13})^{(1)}}{(\hat{M}_{13})^{(1)}}\left[\left((\hat{P}_{13})^{(1)} + G_{14}^{0}\right) e^{\left(-\frac{(\hat{P}_{13})^{(1)} + G_{14}^{0}}{G_{14}^{0}}\right)} + (\hat{P}_{13})^{(1)}\right]$$

(G_i^0) is as defined in the statement of theorem 1

Analogous inequalities hold also for $G_{14}, G_{15}, T_{13}, T_{14}, T_{15}$

The operator $\mathcal{A}^{(2)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{16}(t) \le G_{16}^{0} + \int_0^t \left[(a_{16})^{(2)}\left(G_{17}^{0} + (\hat{P}_{16})^{(6)} e^{(\hat{M}_{16})^{(2)} s_{(16)}}\right)\right] ds_{(16)} = \left(1 + (a_{16})^{(2)} t\right) G_{17}^{0} + \frac{(a_{16})^{(2)}(\hat{P}_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left(e^{(\hat{M}_{16})^{(2)} t} - 1\right)$$ 169

From which it follows that 170

$$(G_{16}(t) - G_{16}^{0}) e^{-(\hat{M}_{16})^{(2)} t} \le \frac{(a_{16})^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{P}_{16})^{(2)} + G_{17}^{0}\right) e^{\left(-\frac{(\hat{P}_{16})^{(2)} + G_{17}^{0}}{G_{17}^{0}}\right)} + (\hat{P}_{16})^{(2)}\right]$$

Analogous inequalities hold also for $G_{17}, G_{18}, T_{16}, T_{17}, T_{18}$

The operator $\mathcal{A}^{(3)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 171

$$G_{20}(t) \le G_{20}^{0} + \int_0^t \left[(a_{20})^{(3)}\left(G_{21}^{0} + (\hat{P}_{20})^{(3)} e^{(\hat{M}_{20})^{(3)} s_{(20)}}\right)\right] ds_{(20)} =$$

$$\left(1 + (a_{20})^{(3)} t\right) G_{21}^{0} + \frac{(a_{20})^{(3)}(\hat{P}_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left(e^{(\hat{M}_{20})^{(3)} t} - 1\right)$$

From which it follows that 172

$$(G_{20}(t) - G_{20}^{0}) e^{-(\hat{M}_{20})^{(3)} t} \le \frac{(a_{20})^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{P}_{20})^{(3)} + G_{21}^{0}\right) e^{\left(-\frac{(\hat{P}_{20})^{(3)} + G_{21}^{0}}{G_{21}^{0}}\right)} + (\hat{P}_{20})^{(3)}\right]$$

Analogous inequalities hold also for $G_{21}, G_{22}, T_{20}, T_{21}, T_{22}$

The operator $\mathcal{A}^{(4)}$ maps the space of functions satisfying into itself .Indeed it is obvious that 173

$$G_{24}(t) \le G_{24}^{0} + \int_0^t \left[(a_{24})^{(4)}\left(G_{25}^{0} + (\hat{P}_{24})^{(4)} e^{(\hat{M}_{24})^{(4)} s_{(24)}}\right)\right] ds_{(24)} =$$

$$\left(1 + (a_{24})^{(4)} t\right) G_{25}^{0} + \frac{(a_{24})^{(4)}(\hat{P}_{24})^{(4)}}{(\hat{M}_{24})^{(4)}}\left(e^{(\hat{M}_{24})^{(4)} t} - 1\right)$$

From which it follows that 174

$$(G_{24}(t)-G_{24}^{0})e^{-(\bar{M}_{24})^{(4)}t} \leq \frac{(a_{24})^{(4)}}{(\bar{M}_{24})^{(4)}}\left[((\hat{P}_{24})^{(4)}+G_{25}^{0})e^{\left(-\frac{(\hat{P}_{24})^{(4)}+G_{25}^{0}}{G_{25}^{0}}\right)}+(\hat{P}_{24})^{(4)}\right]$$

(G_i^0) is as defined in the statement of theorem 4

The operator $\mathcal{A}^{(5)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{28}(t) \leq G_{28}^{0}+\int_0^t\left[(a_{28})^{(5)}\left(G_{29}^{0}+(\hat{P}_{28})^{(5)}e^{(\bar{M}_{28})^{(5)}s_{(28)}}\right)\right]ds_{(28)} =$$

$$\left(1+(a_{28})^{(5)}t\right)G_{29}^{0}+\frac{(a_{28})^{(5)}(\hat{P}_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left(e^{(\bar{M}_{28})^{(5)}t}-1\right)$$

From which it follows that 175

$$(G_{28}(t)-G_{28}^{0})e^{-(\bar{M}_{28})^{(5)}t} \leq \frac{(a_{28})^{(5)}}{(\bar{M}_{28})^{(5)}}\left[((\hat{P}_{28})^{(5)}+G_{29}^{0})e^{\left(-\frac{(\hat{P}_{28})^{(5)}+G_{29}^{0}}{G_{29}^{0}}\right)}+(\hat{P}_{28})^{(5)}\right]$$

(G_i^0) is as defined in the statement of theorem 5

The operator $\mathcal{A}^{(6)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 176

$$G_{32}(t) \leq G_{32}^{0}+\int_0^t\left[(a_{32})^{(6)}\left(G_{33}^{0}+(\hat{P}_{32})^{(6)}e^{(\bar{M}_{32})^{(6)}s_{(32)}}\right)\right]ds_{(32)} =$$

$$\left(1+(a_{32})^{(6)}t\right)G_{33}^{0}+\frac{(a_{32})^{(6)}(\hat{P}_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left(e^{(\bar{M}_{32})^{(6)}t}-1\right)$$

From which it follows that 177

$$(G_{32}(t)-G_{32}^{0})e^{-(\bar{M}_{32})^{(6)}t} \leq \frac{(a_{32})^{(6)}}{(\bar{M}_{32})^{(6)}}\left[((\hat{P}_{32})^{(6)}+G_{33}^{0})e^{\left(-\frac{(\hat{P}_{32})^{(6)}+G_{33}^{0}}{G_{33}^{0}}\right)}+(\hat{P}_{32})^{(6)}\right]$$

(G_i^0) is as defined in the statement of theorem 6

Analogous inequalities hold also for $G_{25}, G_{26}, T_{24}, T_{25}, T_{26}$

The operator $\mathcal{A}^{(7)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that 178

$$G_{36}(t) \leq G_{36}^{0}+\int_0^t\left[(a_{36})^{(7)}\left(G_{37}^{0}+(\hat{P}_{36})^{(7)}e^{(\bar{M}_{36})^{(7)}s_{(36)}}\right)\right]ds_{(36)} =$$

$$\left(1+(a_{36})^{(7)}t\right)G_{37}^{0}+\frac{(a_{36})^{(7)}(\hat{P}_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left(e^{(\bar{M}_{36})^{(7)}t}-1\right)$$

From which it follows that

$$(G_{36}(t)-G_{36}^{0})e^{-(\bar{M}_{36})^{(7)}t} \leq \frac{(a_{36})^{(7)}}{(\bar{M}_{36})^{(7)}}\left[((\hat{P}_{36})^{(7)}+G_{37}^{0})e^{\left(-\frac{(\hat{P}_{36})^{(7)}+G_{37}^{0}}{G_{37}^{0}}\right)}+(\hat{P}_{36})^{(7)}\right]$$

(G_i^0) is as defined in the statement of theorem 7

The operator $\mathcal{A}^{(8)}$ maps the space of functions satisfying Equations into itself .Indeed it is obvious that

$$G_{40}(t) \le G_{40}^0 + \int_0^t \left[(a_{40})^{(8)} \left(G_{41}^0 + (\hat{P}_{40})^{(8)} e^{(\hat{M}_{40})^{(8)} s_{(40)}}\right)\right] ds_{(40)} = \tag{180}$$

$$\left(1 + (a_{40})^{(8)} t\right) G_{41}^0 + \frac{(a_{40})^{(8)} (\hat{P}_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left(e^{(\hat{M}_{40})^{(8)} t} - 1\right)$$

From which it follows that

$$(G_{40}(t) - G_{40}^0) e^{-(\hat{M}_{40})^{(8)} t} \le \frac{(a_{40})^{(8)}}{(\hat{M}_{40})^{(8)}} \left[\left((\hat{P}_{40})^{(8)} + G_{41}^0\right) e^{\left(-\frac{(\hat{P}_{40})^{(8)} + G_{41}^0}{G_{41}^0}\right)} + (\hat{P}_{40})^{(8)}\right] \tag{181}$$

(G_i^0) is as defined in the statement of theorem 8

Analogous inequalities hold also for $G_{41}, G_{42}, T_{40}, T_{41}, T_{42}$

The operator $\mathcal{A}^{(9)}$ maps the space of functions satisfying 34,35,36 into itself .Indeed it is obvious that

$$G_{44}(t) \le G_{44}^0 + \int_0^t \left[(a_{44})^{(9)} \left(G_{45}^0 + (\hat{P}_{44})^{(9)} e^{(\hat{M}_{44})^{(9)} s_{(44)}}\right)\right] ds_{(44)} =$$

$$\left(1 + (a_{44})^{(9)} t\right) G_{45}^0 + \frac{(a_{44})^{(9)} (\hat{P}_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left(e^{(\hat{M}_{44})^{(9)} t} - 1\right)$$

From which it follows that

$$(G_{44}(t) - G_{44}^0) e^{-(\hat{M}_{44})^{(9)} t} \le \frac{(a_{44})^{(9)}}{(\hat{M}_{44})^{(9)}} \left[\left((\hat{P}_{44})^{(9)} + G_{45}^0\right) e^{\left(-\frac{(\hat{P}_{44})^{(9)} + G_{45}^0}{G_{45}^0}\right)} + (\hat{P}_{44})^{(9)}\right]$$

(G_i^0) is as defined in the statement of theorem 9

Analogous inequalities hold also for $G_{45}, G_{46}, T_{44}, T_{45}, T_{46}$

It is now sufficient to take $\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}}, \frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} < 1$ and to choose $\tag{182}$

$(\hat{P}_{13})^{(1)}$ and $(\hat{Q}_{13})^{(1)}$ large to have

$$\frac{(a_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[(\hat{P}_{13})^{(1)} + \left((\hat{P}_{13})^{(1)} + G_j^0\right) e^{-\left(\frac{(\hat{P}_{13})^{(1)} + G_j^0}{G_j^0}\right)}\right] \le (\hat{P}_{13})^{(1)} \tag{183}$$

$$\frac{(b_i)^{(1)}}{(\hat{M}_{13})^{(1)}} \left[\left((\hat{Q}_{13})^{(1)} + T_j^0\right) e^{-\left(\frac{(\hat{Q}_{13})^{(1)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{13})^{(1)}\right] \le (\hat{Q}_{13})^{(1)} \tag{184}$$

In order that the operator $\mathcal{A}^{(1)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(1)}$ is a contraction with respect to the metric $\tag{185}$

$$d\left(\left(G^{(1)}, T^{(1)}\right), \left(G^{(2)}, T^{(2)}\right)\right) =$$

$$\sup_i \left\{\max_{t \in \mathbb{R}_+} \left|G_i^{(1)}(t) - G_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}, \max_{t \in \mathbb{R}_+} \left|T_i^{(1)}(t) - T_i^{(2)}(t)\right| e^{-(\hat{M}_{13})^{(1)} t}\right\}$$

Indeed if we denote

Definition of $\tilde{G},\tilde{T}:\left(\tilde{G},\tilde{T}\right)=\mathcal{A}^{(1)}(G,T)$

It results

$$\left|\tilde{G}_{13}^{(1)}-\tilde{G}_{i}^{(2)}\right|\leq\int_{0}^{t}(a_{13})^{(1)}\left|G_{14}^{(1)}-G_{14}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\,ds_{(13)}+$$

$$\int_{0}^{t}\{(a'_{13})^{(1)}\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)\left|G_{13}^{(1)}-G_{13}^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}+$$

$$G_{13}^{(2)}\left|(a''_{13})^{(1)}\left(T_{14}^{(1)},s_{(13)}\right)-(a''_{13})^{(1)}\left(T_{14}^{(2)},s_{(13)}\right)\right|\ e^{-(\widehat{M}_{13})^{(1)}s_{(13)}}e^{(\widehat{M}_{13})^{(1)}s_{(13)}}\}ds_{(13)}$$

Where $s_{(13)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses it follows

$$\left|G^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{13})^{(1)}t}\leq \qquad 186$$

$$\frac{1}{(\widehat{M}_{13})^{(1)}}\left((a_{13})^{(1)}+\ (a'_{13})^{(1)}+(\widehat{A}_{13})^{(1)}+(\widehat{P}_{13})^{(1)}(\widehat{k}_{13})^{(1)}\right)d\left(\left(G^{(1)},T^{(1)};\ G^{(2)},T^{(2)}\right)\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 1: The fact that we supposed $(a''_{13})^{(1)}$ and $(b''_{13})^{(1)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}\ and\ (\widehat{Q}_{13})^{(1)}e^{(\widehat{M}_{13})^{(1)}t}$ respectively of $\mathbb{R}_{+}$.

If instead of proving the existence of the solution on $\mathbb{R}_{+}$, we have to prove it only on a compact then it suffices to consider that $(a''_{i})^{(1)}$ and $(b''_{i})^{(1)},i=13{,}14{,}15$ depend only on T_{14} and respectively on $G(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 2: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$

From 19 to 24 it results

$$G_i\ (t)\geq G_i^0 e^{\left[-\int_0^t\{(a'_i)^{(1)}-(a''_i)^{(1)}(T_{14}(s_{(13)}),s_{(13)})\}ds_{(13)}\right]}\geq 0$$

$$T_i\ (t)\geq T_i^0 e^{\left(-(b'_i)^{(1)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{13})^{(1)}\right)_1,\left((\widehat{M}_{13})^{(1)}\right)_2\ and\ \left((\widehat{M}_{13})^{(1)}\right)_3$: 187

Remark 3: if G_{13} is bounded, the same property have also $G_{14}\ and\ G_{15}$. indeed if

$G_{13}<(\widehat{M}_{13})^{(1)}$ it follows $\frac{dG_{14}}{dt}\leq\left((\widehat{M}_{13})^{(1)}\right)_1-(a'_{14})^{(1)}G_{14}$ and by integrating

$$G_{14}\leq\left((\widehat{M}_{13})^{(1)}\right)_2=G_{14}^0+2(a_{14})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_1/(a'_{14})^{(1)}$$

In the same way , one can obtain

$$G_{15}\leq\left((\widehat{M}_{13})^{(1)}\right)_3=G_{15}^0+2(a_{15})^{(1)}\left((\widehat{M}_{13})^{(1)}\right)_2/(a'_{15})^{(1)}$$

If $G_{14}\ or\ G_{15}$ is bounded, the same property follows for G_{13} , G_{15} and G_{13} , G_{14} respectively.

Remark 4: If G_{13} *is* bounded, from below, the same property holds for G_{14} *and* G_{15}. The proof is analogous with the preceding one. An analogous property is true if G_{14} is bounded from below. 188

Remark 5: If T_{13} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(1)}(G(t),t)) = (b_{14}')^{(1)}$ then $T_{14} \to \infty$. 189

Definition of $(m)^{(1)}$ and ε_1 :

Indeed let t_1 be so that for $t > t_1$

$(b_{14})^{(1)} - (b_i'')^{(1)}(G(t),t) < \varepsilon_1, T_{13}(t) > (m)^{(1)}$

Then $\frac{dT_{14}}{dt} \geq (a_{14})^{(1)}(m)^{(1)} - \varepsilon_1 T_{14}$ which leads to

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{\varepsilon_1}\right)(1 - e^{-\varepsilon_1 t}) + T_{14}^0 e^{-\varepsilon_1 t}$ If we take t such that $e^{-\varepsilon_1 t} = \frac{1}{2}$ it results

$T_{14} \geq \left(\frac{(a_{14})^{(1)}(m)^{(1)}}{2}\right), \quad t = log\frac{2}{\varepsilon_1}$ By taking now ε_1 sufficiently small one sees that T_{14} is unbounded. The same property holds for T_{15} if $\lim_{t\to\infty}(b_{15}'')^{(1)}(G(t),t) = (b_{15}')^{(1)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}, \frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}} < 1$ and to choose 190

$(\hat{P}_{16})^{(2)}$ *and* $(\hat{Q}_{16})^{(2)}$ large to have

$$\frac{(a_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[(\hat{P}_{16})^{(2)} + \left((\hat{P}_{16})^{(2)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{16})^{(2)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{16})^{(2)} \quad 191$$

$$\frac{(b_i)^{(2)}}{(\hat{M}_{16})^{(2)}}\left[\left((\hat{Q}_{16})^{(2)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{16})^{(2)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{16})^{(2)}\right] \leq (\hat{Q}_{16})^{(2)} \quad 192$$

In order that the operator $\mathcal{A}^{(2)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 193

The operator $\mathcal{A}^{(2)}$ is a contraction with respect to the metric 194

$$d\left(((G_{19})^{(1)},(T_{19})^{(1)}),((G_{19})^{(2)},(T_{19})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{16})^{(2)}t}\}$$

Indeed if we denote 195

Definition of $\widetilde{G_{19}}, \widetilde{T_{19}}$: $(\widetilde{G_{19}}, \widetilde{T_{19}}) = \mathcal{A}^{(2)}(G_{19}, T_{19})$

It results 196

$$|\tilde{G}_{16}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{16})^{(2)}|G_{17}^{(1)} - G_{17}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{(\hat{M}_{16})^{(2)}s_{(16)}}\,ds_{(16)} +$$

$$\int_0^t\{(a_{16}')^{(2)}|G_{16}^{(1)} - G_{16}^{(2)}|e^{-(\hat{M}_{16})^{(2)}s_{(16)}}e^{-(\hat{M}_{16})^{(2)}s_{(16)}} +$$

$(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right)\left|G_{16}^{(1)} - G_{16}^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}} +$

$G_{16}^{(2)}\left|(a_{16}'')^{(2)}\left(T_{17}^{(1)}, s_{(16)}\right) - (a_{16}'')^{(2)}\left(T_{17}^{(2)}, s_{(16)}\right)\right| \; e^{-(\widehat{M}_{16})^{(2)}s_{(16)}} e^{(\widehat{M}_{16})^{(2)}s_{(16)}}\} ds_{(16)}$

Where $s_{(16)}$ represents integrand that is integrated over the interval $[0, t]$ 197

From the hypotheses it follows

$\left|(G_{19})^{(1)} - (G_{19})^{(2)}\right| e^{-(\widehat{M}_{16})^{(2)}t} \leq$
$\frac{1}{(\widehat{M}_{16})^{(2)}}\left((a_{16})^{(2)} + (a_{16}')^{(2)} + (\widehat{A}_{16})^{(2)} + (\widehat{P}_{16})^{(2)}(\widehat{k}_{16})^{(2)}\right) d\left(((G_{19})^{(1)}, (T_{19})^{(1)}; (G_{19})^{(2)}, (T_{19})^{(2)})\right)$

And analogous inequalities for G_i and T_i. Taking into account the hypothesis the result follows 198

Remark 6: The fact that we supposed $(a_{16}'')^{(2)}$ and $(b_{16}'')^{(2)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ and $(\widehat{Q}_{16})^{(2)} e^{(\widehat{M}_{16})^{(2)}t}$ respectively of $\mathbb{R}_+$. 199

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(2)}$ and $(b_i'')^{(2)}, i = 16,17,18$ depend only on T_{17} and respectively on (G_{19})(and not on t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 7: There does not exist any t where $G_i(t) = 0$ and $T_i(t) = 0$ 200

it results

$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(2)} - (a_i'')^{(2)}(T_{17}(s_{(16)}), s_{(16)})\} ds_{(16)}\right]} \geq 0$

$T_i(t) \geq T_i^0 e^{(-(b_i')^{(2)}t)} > 0 \quad \text{for } t > 0$

Definition of $\left((\widehat{M}_{16})^{(2)}\right)_1, \left((\widehat{M}_{16})^{(2)}\right)_2$ and $\left((\widehat{M}_{16})^{(2)}\right)_3$: 201

Remark 8: if G_{16} is bounded, the same property have also G_{17} and G_{18} . indeed if

$G_{16} < (\widehat{M}_{16})^{(2)}$ it follows $\frac{dG_{17}}{dt} \leq \left((\widehat{M}_{16})^{(2)}\right)_1 - (a_{17}')^{(2)} G_{17}$ and by integrating

$G_{17} \leq \left((\widehat{M}_{16})^{(2)}\right)_2 = G_{17}^0 + 2(a_{17})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_1 / (a_{17}')^{(2)}$

In the same way , one can obtain

$G_{18} \leq \left((\widehat{M}_{16})^{(2)}\right)_3 = G_{18}^0 + 2(a_{18})^{(2)}\left((\widehat{M}_{16})^{(2)}\right)_2 / (a_{18}')^{(2)}$

If G_{17} or G_{18} is bounded, the same property follows for G_{16} , G_{18} and G_{16} , G_{17} respectively.

Remark 9: If G_{16} is bounded, from below, the same property holds for G_{17} and G_{18} . The proof is analogous with the preceding one. An analogous property is true if G_{17} is bounded from below. 202

Remark 10: If T_{16} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(2)}((G_{19})(t), t)) = (b_{17}')^{(2)}$ then $T_{17} \to \infty$. 203

Definition of $(m)^{(2)}$ and ε_2 :

Indeed let t_2 be so that for $t > t_2$

$(b_{17})^{(2)} - (b_i'')^{(2)}((G_{19})(t), t) < \varepsilon_2, T_{16}(t) > (m)^{(2)}$

Then $\frac{dT_{17}}{dt} \geq (a_{17})^{(2)}(m)^{(2)} - \varepsilon_2 T_{17}$ which leads to 204

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{\varepsilon_2}\right)(1 - e^{-\varepsilon_2 t}) + T_{17}^0 e^{-\varepsilon_2 t}$ If we take t such that $e^{-\varepsilon_2 t} = \frac{1}{2}$ it results

$T_{17} \geq \left(\frac{(a_{17})^{(2)}(m)^{(2)}}{2}\right), \quad t = \log\frac{2}{\varepsilon_2}$ By taking now ε_2 sufficiently small one sees that T_{17} is 205
unbounded. The same property holds for T_{18} if $\lim_{t\to\infty}(b_{18}'')^{(2)}\left((G_{19})(t), t\right) = (b_{18}')^{(2)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}, \frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}} < 1$ and to choose 207

$(\hat{P}_{20})^{(3)}$ and $(\hat{Q}_{20})^{(3)}$ large to have

$$\frac{(a_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[(\hat{P}_{20})^{(3)} + \left((\hat{P}_{20})^{(3)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{20})^{(3)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{20})^{(3)} \quad 208$$

$$\frac{(b_i)^{(3)}}{(\hat{M}_{20})^{(3)}}\left[\left((\hat{Q}_{20})^{(3)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{20})^{(3)} + T_j^0}{T_j^0}\right)} + (\hat{Q}_{20})^{(3)}\right] \leq (\hat{Q}_{20})^{(3)} \quad 209$$

In order that the operator $\mathcal{A}^{(3)}$ transforms the space of sextuples of functions G_i, T_i satisfying 210
Equations into itself

The operator $\mathcal{A}^{(3)}$ is a contraction with respect to the metric 211

$$d\left(((G_{23})^{(1)}, (T_{23})^{(1)}), ((G_{23})^{(2)}, (T_{23})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\hat{M}_{20})^{(3)}t}\}$$

Indeed if we denote 212

Definition of $\widetilde{G_{23}}, \widetilde{T_{23}} :\left(\widetilde{(G_{23})}, \widetilde{(T_{23})}\right) = \mathcal{A}^{(3)}\left((G_{23}), (T_{23})\right)$

It results 213

$$|\tilde{G}_{20}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{20})^{(3)}|G_{21}^{(1)} - G_{21}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\,ds_{(20)} +$$

$$\int_0^t\{(a_{20}')^{(3)}|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{-(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)})|G_{20}^{(1)} - G_{20}^{(2)}|e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}} +$$

$$G_{20}^{(2)}|(a_{20}'')^{(3)}(T_{21}^{(1)}, s_{(20)}) - (a_{20}'')^{(3)}(T_{21}^{(2)}, s_{(20)})|\ e^{-(\hat{M}_{20})^{(3)}s_{(20)}}e^{(\hat{M}_{20})^{(3)}s_{(20)}}\}ds_{(20)}$$

Where $s_{(20)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses it follows

$$\left|G_{23}{}^{(1)} - G_{23}{}^{(2)}\right| e^{-(\widehat{M}_{20})^{(3)}t} \leq \frac{1}{(\widehat{M}_{20})^{(3)}}\left((a_{20})^{(3)} + (a'_{20})^{(3)} + (\widehat{A}_{20})^{(3)} + (\widehat{P}_{20})^{(3)}(\widehat{k}_{20})^{(3)}\right) d\left(((G_{23})^{(1)}, (T_{23})^{(1)};\ (G_{23})^{(2)}, (T_{23})^{(2)})\right) \quad 214$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 11: The fact that we supposed $(a''_{20})^{(3)}$ and $(b''_{20})^{(3)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ *and* $(\widehat{Q}_{20})^{(3)}e^{(\widehat{M}_{20})^{(3)}t}$ respectively of $\mathbb{R}_+$. 215

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(3)}$ and $(b_i'')^{(3)}, i = 20,21,22$ depend only on T_{21} and respectively on $(G_{23})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 12: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 216

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(3)} - (a_i'')^{(3)}(T_{21}(s_{(20)}), s_{(20)})\} ds_{(20)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{\left(-(b_i')^{(3)}t\right)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{20})^{(3)}\right)_1, \left((\widehat{M}_{20})^{(3)}\right)_2\ and\ \left((\widehat{M}_{20})^{(3)}\right)_3$: 217

Remark 13: if G_{20} is bounded, the same property have also $G_{21}\ and\ G_{22}$. indeed if

$G_{20} < (\widehat{M}_{20})^{(3)}$ it follows $\frac{dG_{21}}{dt} \leq \left((\widehat{M}_{20})^{(3)}\right)_1 - (a'_{21})^{(3)}G_{21}$ and by integrating

$$G_{21} \leq \left((\widehat{M}_{20})^{(3)}\right)_2 = G_{21}^0 + 2(a_{21})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_1/(a'_{21})^{(3)}$$

In the same way , one can obtain

$$G_{22} \leq \left((\widehat{M}_{20})^{(3)}\right)_3 = G_{22}^0 + 2(a_{22})^{(3)}\left((\widehat{M}_{20})^{(3)}\right)_2/(a'_{22})^{(3)}$$

If $G_{21}\ or\ G_{22}$ is bounded, the same property follows for G_{20} , G_{22} and G_{20} , G_{21} respectively.

Remark 14: If G_{20} *is* bounded, from below, the same property holds for $G_{21}\ and\ G_{22}$. The proof is analogous with the preceding one. An analogous property is true if G_{21} is bounded from below. 218

Remark 15: If T_{20} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(3)}\left((G_{23})(t), t\right)) = (b'_{21})^{(3)}$ then $T_{21} \to \infty$. 219

Definition of $(m)^{(3)}$ and ε_3 :

Indeed let t_3 be so that for $t > t_3$

$$(b_{21})^{(3)} - (b_i'')^{(3)}\left((G_{23})(t), t\right) < \varepsilon_3, T_{20}\,(t) > (m)^{(3)}$$

Then $\frac{dT_{21}}{dt} \geq (a_{21})^{(3)}(m)^{(3)} - \varepsilon_3 T_{21}$ which leads to 220

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{\varepsilon_3}\right)(1-e^{-\varepsilon_3 t}) + T_{21}^0 e^{-\varepsilon_3 t}$ If we take t such that $e^{-\varepsilon_3 t} = \frac{1}{2}$ it results

$T_{21} \geq \left(\frac{(a_{21})^{(3)}(m)^{(3)}}{2}\right), \quad t = log\,\frac{2}{\varepsilon_3}$ By taking now ε_3 sufficiently small one sees that T_{21} is unbounded. The same property holds for T_{22} if $\lim_{t\to\infty}(b_{22}'')^{(3)}\left((G_{23})(t), t\right) = (b_{22}')^{(3)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}, \frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}} < 1$ and to choose 221

$(\widehat{P}_{24})^{(4)}$ and $(\widehat{Q}_{24})^{(4)}$ large to have

$$\frac{(a_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[(\widehat{P}_{24})^{(4)} + \left((\widehat{P}_{24})^{(4)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{24})^{(4)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{24})^{(4)} \quad 222$$

$$\frac{(b_i)^{(4)}}{(\widehat{M}_{24})^{(4)}}\left[\left((\widehat{Q}_{24})^{(4)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{24})^{(4)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{24})^{(4)}\right] \leq (\widehat{Q}_{24})^{(4)} \quad 223$$

In order that the operator $\mathcal{A}^{(4)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself 224

The operator $\mathcal{A}^{(4)}$ is a contraction with respect to the metric 225

$$d\left(((G_{27})^{(1)}, (T_{27})^{(1)}), ((G_{27})^{(2)}, (T_{27})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t) - G_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}, \max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t) - T_i^{(2)}(t)\right|e^{-(\widehat{M}_{24})^{(4)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{27}}), (\widetilde{T_{27}})$: $\left((\widetilde{G_{27}}), (\widetilde{T_{27}})\right) = \mathcal{A}^{(4)}((G_{27}), (T_{27}))$

It results

$$\left|\tilde{G}_{24}^{(1)} - \tilde{G}_i^{(2)}\right| \leq \int_0^t (a_{24})^{(4)}\left|G_{25}^{(1)} - G_{25}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\,ds_{(24)} +$$

$$\int_0^t\{(a_{24}')^{(4)}\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{-(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right)\left|G_{24}^{(1)} - G_{24}^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}} +$$

$$G_{24}^{(2)}\left|(a_{24}'')^{(4)}\left(T_{25}^{(1)}, s_{(24)}\right) - (a_{24}'')^{(4)}\left(T_{25}^{(2)}, s_{(24)}\right)\right|\ e^{-(\widehat{M}_{24})^{(4)}s_{(24)}}e^{(\widehat{M}_{24})^{(4)}s_{(24)}}\}ds_{(24)}$$

Where $s_{(24)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses on Equations it follows

$$\left|(G_{27})^{(1)} - (G_{27})^{(2)}\right|e^{-(\widehat{M}_{24})^{(4)}t} \leq \quad 226$$
$$\frac{1}{(\widehat{M}_{24})^{(4)}}\left((a_{24})^{(4)} + (a_{24}')^{(4)} + (\widehat{A}_{24})^{(4)} + (\widehat{P}_{24})^{(4)}(\widehat{k}_{24})^{(4)}\right)d\left(((G_{27})^{(1)}, (T_{27})^{(1)};\ (G_{27})^{(2)}, (T_{27})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 16: The fact that we supposed $(a''_{24})^{(4)}$ and $(b''_{24})^{(4)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ *and* $(\widehat{Q}_{24})^{(4)}e^{(\widehat{M}_{24})^{(4)}t}$ respectively of $\mathbb{R}_+$. 227

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(4)}$ and $(b_i'')^{(4)}, i = 24,25,26$ depend only on T_{25} and respectively on (G_{27})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 17: There does not exist any t where $G_i(t) = 0$ *and* $T_i(t) = 0$ 228

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(4)} - (a_i'')^{(4)}(T_{25}(s_{(24)}), s_{(24)})\} ds_{(24)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(4)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{24})^{(4)}\right)_1, \left((\widehat{M}_{24})^{(4)}\right)_2$ *and* $\left((\widehat{M}_{24})^{(4)}\right)_3$: 229

Remark 18: if G_{24} is bounded, the same property have also G_{25} *and* G_{26} . indeed if

$G_{24} < (\widehat{M}_{24})^{(4)}$ it follows $\frac{dG_{25}}{dt} \leq \left((\widehat{M}_{24})^{(4)}\right)_1 - (a'_{25})^{(4)}G_{25}$ and by integrating

$$G_{25} \leq \left((\widehat{M}_{24})^{(4)}\right)_2 = G_{25}^0 + 2(a_{25})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_1/(a'_{25})^{(4)}$$

In the same way , one can obtain

$$G_{26} \leq \left((\widehat{M}_{24})^{(4)}\right)_3 = G_{26}^0 + 2(a_{26})^{(4)}\left((\widehat{M}_{24})^{(4)}\right)_2/(a'_{26})^{(4)}$$

If G_{25} *or* G_{26} is bounded, the same property follows for G_{24} , G_{26} and G_{24} , G_{25} respectively.

Remark 19: If G_{24} *is* bounded, from below, the same property holds for G_{25} *and* G_{26} . The proof is analogous with the preceding one. An analogous property is true if G_{25} is bounded from below. 230

Remark 20: If T_{24} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(4)}((G_{27})(t), t)) = (b'_{25})^{(4)}$ then $T_{25} \to \infty$. 231

Definition of $(m)^{(4)}$ and ε_4 :

Indeed let t_4 be so that for $t > t_4$

$$(b_{25})^{(4)} - (b_i'')^{(4)}((G_{27})(t), t) < \varepsilon_4, T_{24}(t) > (m)^{(4)}$$

Then $\frac{dT_{25}}{dt} \geq (a_{25})^{(4)}(m)^{(4)} - \varepsilon_4 T_{25}$ which leads to 232

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{\varepsilon_4}\right)(1 - e^{-\varepsilon_4 t}) + T_{25}^0 e^{-\varepsilon_4 t}$ If we take t such that $e^{-\varepsilon_4 t} = \frac{1}{2}$ it results

$T_{25} \geq \left(\frac{(a_{25})^{(4)}(m)^{(4)}}{2}\right), \quad t = log\frac{2}{\varepsilon_4}$ By taking now ε_4 sufficiently small one sees that T_{25} is unbounded. The same property holds for T_{26} if $\lim_{t\to\infty}(b''_{26})^{(4)}\left((G_{27})(t), t\right) = (b'_{26})^{(4)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for $G_{29}, G_{30}, T_{28}, T_{29}, T_{30}$

It is now sufficient to take $\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}, \frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}} < 1$ and to choose 233

$(\widehat{P}_{28})^{(5)}$ and $(\widehat{Q}_{28})^{(5)}$ large to have

$$\frac{(a_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[(\widehat{P}_{28})^{(5)} + \left((\widehat{P}_{28})^{(5)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{28})^{(5)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{28})^{(5)} \quad 234$$

$$\frac{(b_i)^{(5)}}{(\widehat{M}_{28})^{(5)}}\left[\left((\widehat{Q}_{28})^{(5)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{28})^{(5)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{28})^{(5)}\right] \leq (\widehat{Q}_{28})^{(5)} \quad 235$$

In order that the operator $\mathcal{A}^{(5)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(5)}$ is a contraction with respect to the metric 236

$$d\left(((G_{31})^{(1)}, (T_{31})^{(1)}), ((G_{31})^{(2)}, (T_{31})^{(2)})\right) =$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t) - G_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}, \max_{t\in\mathbb{R}_+}|T_i^{(1)}(t) - T_i^{(2)}(t)|e^{-(\widehat{M}_{28})^{(5)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{31}}), (\widetilde{T_{31}})$: $\left((\widetilde{G_{31}}), (\widetilde{T_{31}})\right) = \mathcal{A}^{(5)}((G_{31}), (T_{31}))$

It results

$$|\tilde{G}_{28}^{(1)} - \tilde{G}_i^{(2)}| \leq \int_0^t (a_{28})^{(5)}|G_{29}^{(1)} - G_{29}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\,ds_{(28)} +$$

$$\int_0^t\{(a'_{28})^{(5)}|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{-(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)})|G_{28}^{(1)} - G_{28}^{(2)}|e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}} +$$

$$G_{28}^{(2)}|(a''_{28})^{(5)}(T_{29}^{(1)}, s_{(28)}) - (a''_{28})^{(5)}(T_{29}^{(2)}, s_{(28)})|\ e^{-(\widehat{M}_{28})^{(5)}s_{(28)}}e^{(\widehat{M}_{28})^{(5)}s_{(28)}}\}ds_{(28)}$$

Where $s_{(28)}$ represents integrand that is integrated over the interval $[0, t]$

From the hypotheses on it follows

$$|(G_{31})^{(1)} - (G_{31})^{(2)}|e^{-(\widehat{M}_{28})^{(5)}t} \leq \quad 237$$
$$\frac{1}{(\widehat{M}_{28})^{(5)}}\left((a_{28})^{(5)} + (a'_{28})^{(5)} + (\widehat{A}_{28})^{(5)} + (\widehat{P}_{28})^{(5)}(\widehat{k}_{28})^{(5)}\right)d\left(((G_{31})^{(1)}, (T_{31})^{(1)}; (G_{31})^{(2)}, (T_{31})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 21: The fact that we supposed $(a''_{28})^{(5)}$ and $(b''_{28})^{(5)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by 238

$(\widehat{P}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ *and* $(\widehat{Q}_{28})^{(5)}e^{(\widehat{M}_{28})^{(5)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(5)}$ and $(b_i'')^{(5)}, i = 28,29,30$ depend only on T_{29} and respectively on $(G_{31})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 22: There does not exist any t where $G_i\,(t) = 0\ and\ T_i\,(t) = 0$ 239

it results

$$G_i\,(t) \geq G_i^0 e^{\left[-\int_0^t\{(a_i')^{(5)}-(a_i'')^{(5)}(T_{29}(s_{(28)}),s_{(28)})\}ds_{(28)}\right]} \geq 0$$

$$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(5)}t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{28})^{(5)}\right)_1, \left((\widehat{M}_{28})^{(5)}\right)_2\ and\ \left((\widehat{M}_{28})^{(5)}\right)_3$: 240

Remark 23: if G_{28} is bounded, the same property have also $G_{29}\ and\ G_{30}$. indeed if

$G_{28} < (\widehat{M}_{28})^{(5)}$ it follows $\frac{dG_{29}}{dt} \leq \left((\widehat{M}_{28})^{(5)}\right)_1 - (a_{29}')^{(5)}G_{29}$ and by integrating

$$G_{29} \leq \left((\widehat{M}_{28})^{(5)}\right)_2 = G_{29}^0 + 2(a_{29})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_1/(a_{29}')^{(5)}$$

In the same way , one can obtain

$$G_{30} \leq \left((\widehat{M}_{28})^{(5)}\right)_3 = G_{30}^0 + 2(a_{30})^{(5)}\left((\widehat{M}_{28})^{(5)}\right)_2/(a_{30}')^{(5)}$$

If $G_{29}\ or\ G_{30}$ is bounded, the same property follows for G_{28} , G_{30} and G_{28} , G_{29} respectively.

Remark 24: If $G_{28}\ \ is$ bounded, from below, the same property holds for $G_{29}\ and\ G_{30}$. The proof is analogous with the preceding one. An analogous property is true if G_{29} is bounded from below. 241

Remark 25: If T_{28} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(5)}\ ((G_{31})(t),t)) = (b_{29}')^{(5)}$ then $T_{29}\ \to \infty$. 242

Definition of $(m)^{(5)}$ and ε_5 :

Indeed let t_5 be so that for $t\ > t_5$

$$(b_{29})^{(5)} - (b_i'')^{(5)}((G_{31})(t),t) < \varepsilon_5, T_{28}\,(t) > (m)^{(5)}$$

Then $\frac{dT_{29}}{dt} \geq (a_{29})^{(5)}(m)^{(5)} - \varepsilon_5 T_{29}$ which leads to 243

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{\varepsilon_5}\right)(1 - e^{-\varepsilon_5 t}) + T_{29}^0 e^{-\varepsilon_5 t}$ If we take t such that $e^{-\varepsilon_5 t} = \frac{1}{2}$ it results

$T_{29}\ \geq \left(\frac{(a_{29})^{(5)}(m)^{(5)}}{2}\right), \quad t = log\frac{2}{\varepsilon_5}$ By taking now ε_5 sufficiently small one sees that T_{29} is unbounded. The same property holds for T_{30} if $\lim_{t\to\infty}(b_{30}'')^{(5)}\left((G_{31})(t),t\right) = (b_{30}')^{(5)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{33} , $G_{34}, T_{32}, T_{33}, T_{34}$

It is now sufficient to take $\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}$, $\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}} < 1$ and to choose 244

$(\widehat{P}_{32})^{(6)}$ and $(\widehat{Q}_{32})^{(6)}$ large to have

$$\frac{(a_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[(\widehat{P}_{32})^{(6)}+\left((\widehat{P}_{32})^{(6)}+G_j^0\right)e^{-\left(\frac{(\widehat{P}_{32})^{(6)}+G_j^0}{G_j^0}\right)}\right]\leq(\widehat{P}_{32})^{(6)} \tag{245}$$

$$\frac{(b_i)^{(6)}}{(\widehat{M}_{32})^{(6)}}\left[\left((\widehat{Q}_{32})^{(6)}+T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{32})^{(6)}+T_j^0}{T_j^0}\right)}+(\widehat{Q}_{32})^{(6)}\right]\leq(\widehat{Q}_{32})^{(6)} \tag{246}$$

In order that the operator $\mathcal{A}^{(6)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(6)}$ is a contraction with respect to the metric (247)

$$d\left(((G_{35})^{(1)},(T_{35})^{(1)}),((G_{35})^{(2)},(T_{35})^{(2)})\right)=$$

$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{32})^{(6)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{35}}),(\widetilde{T_{35}})$: $\left((\widetilde{G_{35}}),(\widetilde{T_{35}})\right)=\mathcal{A}^{(6)}\left((G_{35}),(T_{35})\right)$

It results

$$|\tilde{G}_{32}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{32})^{(6)}|G_{33}^{(1)}-G_{33}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\,ds_{(32)}+$$

$$\int_0^t\{(a'_{32})^{(6)}|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)|G_{32}^{(1)}-G_{32}^{(2)}|e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}+$$

$$G_{32}^{(2)}|(a''_{32})^{(6)}\left(T_{33}^{(1)},s_{(32)}\right)-(a''_{32})^{(6)}\left(T_{33}^{(2)},s_{(32)}\right)|\;e^{-(\widehat{M}_{32})^{(6)}s_{(32)}}e^{(\widehat{M}_{32})^{(6)}s_{(32)}}\}ds_{(32)}$$

Where $s_{(32)}$ represents integrand that is integrated over the interval [0, t]

From the hypotheses it follows

$$|(G_{35})^{(1)}-(G_{35})^{(2)}|e^{-(\widehat{M}_{32})^{(6)}t}\leq \tag{248}$$
$$\frac{1}{(\widehat{M}_{32})^{(6)}}\left((a_{32})^{(6)}+(a'_{32})^{(6)}+(\widehat{A}_{32})^{(6)}+(\widehat{P}_{32})^{(6)}(\widehat{k}_{32})^{(6)}\right)d\left(((G_{35})^{(1)},(T_{35})^{(1)};(G_{35})^{(2)},(T_{35})^{(2)})\right)$$

And analogous inequalities for G_i *and* T_i. Taking into account the hypothesis the result follows

Remark 26: The fact that we supposed $(a''_{32})^{(6)}$ and $(b''_{32})^{(6)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ *and* $(\widehat{Q}_{32})^{(6)}e^{(\widehat{M}_{32})^{(6)}t}$ respectively of $\mathbb{R}_+$. (249)

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(6)}$ and $(b_i'')^{(6)}, i=32,33,34$ depend only on T_{33} and respectively on (G_{35})(*and not on* t) and hypothesis can replaced by a usual Lipschitz condition.

Remark 27: There does not exist any t where $G_i(t) = 0 \; and \; T_i(t) = 0$ 250

it results

$$G_i(t) \geq G_i^0 e^{\left[-\int_0^t \{(a_i')^{(6)} - (a_i'')^{(6)}(T_{33}(s_{(32)}), s_{(32)})\} ds_{(32)}\right]} \geq 0$$

$$T_i(t) \geq T_i^0 e^{(-(b_i')^{(6)} t)} > 0 \quad \text{for t} > 0$$

Definition of $\left((\widehat{M}_{32})^{(6)}\right)_1, \left((\widehat{M}_{32})^{(6)}\right)_2 \; and \; \left((\widehat{M}_{32})^{(6)}\right)_3$: 251

Remark 28: if G_{32} is bounded, the same property have also $G_{33} \; and \; G_{34}$. indeed if

$G_{32} < (\widehat{M}_{32})^{(6)}$ it follows $\frac{dG_{33}}{dt} \leq \left((\widehat{M}_{32})^{(6)}\right)_1 - (a_{33}')^{(6)} G_{33}$ and by integrating

$$G_{33} \leq \left((\widehat{M}_{32})^{(6)}\right)_2 = G_{33}^0 + 2(a_{33})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_1 / (a_{33}')^{(6)}$$

In the same way , one can obtain

$$G_{34} \leq \left((\widehat{M}_{32})^{(6)}\right)_3 = G_{34}^0 + 2(a_{34})^{(6)} \left((\widehat{M}_{32})^{(6)}\right)_2 / (a_{34}')^{(6)}$$

If $G_{33} \; or \; G_{34}$ is bounded, the same property follows for G_{32} , G_{34} and G_{32} , G_{33} respectively.

Remark 29: If G_{32} *is* bounded, from below, the same property holds for $G_{33} \; and \; G_{34}$. The proof is analogous with the preceding one. An analogous property is true if G_{33} is bounded from below. 252

Remark 30: If T_{32} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(6)}((G_{35})(t), t)) = (b_{33}')^{(6)}$ then $T_{33} \to \infty$. 253

Definition of $(m)^{(6)}$ and ε_6 :

Indeed let t_6 be so that for $t > t_6$

$$(b_{33})^{(6)} - (b_i'')^{(6)}\left((G_{35})(t), t\right) < \varepsilon_6, T_{32}(t) > (m)^{(6)}$$

Then $\frac{dT_{33}}{dt} \geq (a_{33})^{(6)}(m)^{(6)} - \varepsilon_6 T_{33}$ which leads to 254

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{\varepsilon_6}\right)(1 - e^{-\varepsilon_6 t}) + T_{33}^0 e^{-\varepsilon_6 t}$ If we take t such that $e^{-\varepsilon_6 t} = \frac{1}{2}$ it results

$T_{33} \geq \left(\frac{(a_{33})^{(6)}(m)^{(6)}}{2}\right), \quad t = log\frac{2}{\varepsilon_6}$ By taking now ε_6 sufficiently small one sees that T_{33} is unbounded. The same property holds for T_{34} if $\lim_{t\to\infty}(b_{34}'')^{(6)}\left((G_{35})(t), t(t), t\right) = (b_{34}')^{(6)}$

We now state a more precise theorem about the behaviors at infinity of the solutions of equations

Analogous inequalities hold also for G_{37} , $G_{38}, T_{36}, T_{37}, T_{38}$ 255

It is now sufficient to take $\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}$, $\frac{(b_i)^{(7)}}{(\hat{M}_{36})^{(7)}} < 1$ and to choose
$(\hat{P}_{36})^{(7)}$ and $(\hat{Q}_{36})^{(7)}$ large to have

$$\frac{(a_i)^{(7)}}{(\hat{M}_{36})^{(7)}}\left[(\hat{P}_{36})^{(7)} + \left((\hat{P}_{36})^{(7)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{36})^{(7)} + G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{36})^{(7)} \quad 256$$

257

$$\frac{(b_i)^{(7)}}{(\widehat{M}_{36})^{(7)}}\left[\left((\hat{Q}_{36})^{(7)}+T_j^0\right)e^{-\left(\frac{(\hat{Q}_{36})^{(7)}+T_j^0}{T_j^0}\right)}+(\hat{Q}_{36})^{(7)}\right]\leq(\hat{Q}_{36})^{(7)}$$

In order that the operator $\mathcal{A}^{(7)}$ transforms the space of sextuples of functions G_i, T_i satisfying Equations into itself

The operator $\mathcal{A}^{(7)}$ is a contraction with respect to the metric 258

$$d\left(((G_{39})^{(1)},(T_{39})^{(1)}),((G_{39})^{(2)},(T_{39})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}|G_i^{(1)}(t)-G_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t},\max_{t\in\mathbb{R}_+}|T_i^{(1)}(t)-T_i^{(2)}(t)|e^{-(\widehat{M}_{36})^{(7)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{39}}),(\widetilde{T_{39}}):\left((\widetilde{G_{39}}),(\widetilde{T_{39}})\right)=\mathcal{A}^{(7)}((G_{39}),(T_{39}))$

It results

$$|\tilde{G}_{36}^{(1)}-\tilde{G}_i^{(2)}|\leq\int_0^t(a_{36})^{(7)}|G_{37}^{(1)}-G_{37}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}ds_{(36)}+$$
$$\int_0^t\{(a'_{36})^{(7)}|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})|G_{36}^{(1)}-G_{36}^{(2)}|e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}+$$
$$G_{36}^{(2)}|(a''_{36})^{(7)}(T_{37}^{(1)},s_{(36)})-(a''_{36})^{(7)}(T_{37}^{(2)},s_{(36)})|\ e^{-(\widehat{M}_{36})^{(7)}s_{(36)}}e^{(\widehat{M}_{36})^{(7)}s_{(36)}}\}ds_{(36)}$$

Where $s_{(36)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$

From the hypotheses on it follows

$$|(G_{39})^{(1)}-(G_{39})^{(2)}|e^{-(\widehat{M}_{36})^{(7)}t}\leq$$ 259
$$\frac{1}{(\widehat{M}_{36})^{(7)}}\left((a_{36})^{(7)}+(a'_{36})^{(7)}+(\widehat{A}_{36})^{(7)}+(\widehat{P}_{36})^{(7)}(\widehat{k}_{36})^{(7)}\right)d\left(((G_{39})^{(1)},(T_{39})^{(1)};\ (G_{39})^{(2)},(T_{39})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 31: The fact that we supposed $(a''_{36})^{(7)}$ and $(b''_{36})^{(7)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}\ and\ (\widehat{Q}_{36})^{(7)}e^{(\widehat{M}_{36})^{(7)}t}$ respectively of $\mathbb{R}_+$. 260

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(7)}$ and $(b_i'')^{(7)}, i=36,37,38$ depend only on T_{37} and respectively on $(G_{39})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 32: There does not exist any t where $G_i\ (t)=0\ and\ T_i\ (t)=0$ 261

it results

$$G_i\ (t)\geq G_i^0e^{\left[-\int_0^t\{(a_i')^{(7)}-(a_i'')^{(7)}(T_{37}(s_{(36)}),s_{(36)})\}ds_{(36)}\right]}\geq 0$$

$T_i\,(t) \geq T_i^0 e^{(-(b_i')^{(7)}t)} > 0$ for t > 0

Definition of $\left((\widehat{M}_{36})^{(7)}\right)_1, \left((\widehat{M}_{36})^{(7)}\right)_2$ *and* $\left((\widehat{M}_{36})^{(7)}\right)_3$: 262

Remark 33: if G_{36} is bounded, the same property have also G_{37} *and* G_{38} . indeed if

$G_{36} < (\widehat{M}_{36})^{(7)}$ it follows $\frac{dG_{37}}{dt} \leq \left((\widehat{M}_{36})^{(7)}\right)_1 - (a_{37}')^{(7)} G_{37}$ and by integrating

$G_{37} \leq \left((\widehat{M}_{36})^{(7)}\right)_2 = G_{37}^0 + 2(a_{37})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_1/(a_{37}')^{(7)}$

In the same way , one can obtain

$G_{38} \leq \left((\widehat{M}_{36})^{(7)}\right)_3 = G_{38}^0 + 2(a_{38})^{(7)}\left((\widehat{M}_{36})^{(7)}\right)_2/(a_{38}')^{(7)}$

If G_{37} *or* G_{38} is bounded, the same property follows for G_{36} , G_{38} and G_{36} , G_{37} respectively.

Remark 34: If G_{36} *is* bounded, from below, the same property holds for G_{37} *and* G_{38} . The proof is analogous with the preceding one. An analogous property is true if G_{37} is bounded from below. 263

Remark 35: If T_{36} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(7)}((G_{39})(t),t)) = (b_{37}')^{(7)}$ then $T_{37} \to \infty$. 264

Definition of $(m)^{(7)}$ and ε_7 :

Indeed let t_7 be so that for $t > t_7$

$(b_{37})^{(7)} - (b_i'')^{(7)}((G_{39})(t),t) < \varepsilon_7, T_{36}\,(t) > (m)^{(7)}$

Then $\frac{dT_{37}}{dt} \geq (a_{37})^{(7)}(m)^{(7)} - \varepsilon_7 T_{37}$ which leads to 265

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{\varepsilon_7}\right)(1 - e^{-\varepsilon_7 t}) + T_{37}^0 e^{-\varepsilon_7 t}$ If we take t such that $e^{-\varepsilon_7 t} = \frac{1}{2}$ it results

$T_{37} \geq \left(\frac{(a_{37})^{(7)}(m)^{(7)}}{2}\right)$, $t = log\,\frac{2}{\varepsilon_7}$ By taking now ε_7 sufficiently small one sees that T_{37} is unbounded. The same property holds for T_{38} if $\lim_{t\to\infty}(b_{38}'')^{(7)}\left((G_{39})(t),t\right) = (b_{38}')^{(7)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations

It is now sufficient to take $\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}$, $\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}} < 1$ and to choose 266
$(\hat{P}_{40})^{(8)}$ and $(\hat{Q}_{40})^{(8)}$ large to have

$$\frac{(a_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[(\hat{P}_{40})^{(8)} + \left((\hat{P}_{40})^{(8)} + G_j^0\right)e^{-\left(\frac{(\hat{P}_{40})^{(8)}+G_j^0}{G_j^0}\right)}\right] \leq (\hat{P}_{40})^{(8)} \quad 267$$

$$\frac{(b_i)^{(8)}}{(\hat{M}_{40})^{(8)}}\left[\left((\hat{Q}_{40})^{(8)} + T_j^0\right)e^{-\left(\frac{(\hat{Q}_{40})^{(8)}+T_j^0}{T_j^0}\right)} + (\hat{Q}_{40})^{(8)}\right] \leq (\hat{Q}_{40})^{(8)} \quad 268$$

In order that the operator $\mathcal{A}^{(8)}$ transforms the space of sextuples of functions G_i , T_i satisfying Equations into itself
The operator $\mathcal{A}^{(8)}$ is a contraction with respect to the metric

$$d\left(((G_{43})^{(1)},(T_{43})^{(1)}),((G_{43})^{(2)},(T_{43})^{(2)})\right)= \quad 269$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{40})^{(8)}t}\}$$

Indeed if we denote 270

Definition of $\widetilde{(G_{43})},\widetilde{(T_{43})}$: $\left(\widetilde{(G_{43})},\widetilde{(T_{43})}\right)=\mathcal{A}^{(8)}((G_{43}),(T_{43}))$

It results 271

$$\left|\tilde{G}_{40}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{40})^{(8)}\left|G_{41}^{(1)}-G_{41}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\,ds_{(40)}+$$
$$\int_0^t\{(a'_{40})^{(8)}\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)\left|G_{40}^{(1)}-G_{40}^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}+$$
$$G_{40}^{(2)}\left|(a''_{40})^{(8)}\left(T_{41}^{(1)},s_{(40)}\right)-(a''_{40})^{(8)}\left(T_{41}^{(2)},s_{(40)}\right)\right|\ e^{-(\widehat{M}_{40})^{(8)}s_{(40)}}e^{(\widehat{M}_{40})^{(8)}s_{(40)}}\}ds_{(40)}$$

Where $s_{(40)}$ represents integrand that is integrated over the interval $[0,\mathrm{t}]$ 272

From the hypotheses it follows

$$\left|(G_{43})^{(1)}-(G_{43})^{(2)}\right|e^{-(\widehat{M}_{40})^{(8)}t}\le \quad 273$$
$$\frac{1}{(\widehat{M}_{40})^{(8)}}\left((a_{40})^{(8)}+(a'_{40})^{(8)}+(\widehat{A}_{40})^{(8)}+(\widehat{P}_{40})^{(8)}(\widehat{k}_{40})^{(8)}\right)d\left(((G_{43})^{(1)},(T_{43})^{(1)};\ (G_{43})^{(2)},(T_{43})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis the result follows

Remark 36: The fact that we supposed $(a''_{40})^{(8)}$ and $(b''_{40})^{(8)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}\ and\ (\widehat{Q}_{40})^{(8)}e^{(\widehat{M}_{40})^{(8)}t}$ respectively of $\mathbb{R}_+$. 274

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(8)}$ and $(b_i'')^{(8)}, i=40{,}41{,}42$ depend only on T_{41} and respectively on $(G_{43})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 37 There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$ 275

it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(8)}-(a_i'')^{(8)}(T_{41}(s_{(40)}),s_{(40)})\}ds_{(40)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(8)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{40})^{(8)}\right)_1,\left((\widehat{M}_{40})^{(8)}\right)_2\ and\ \left((\widehat{M}_{40})^{(8)}\right)_3$: 276

Remark 38: if G_{40} is bounded, the same property have also $G_{41}\ and\ G_{42}$. indeed if

$G_{40} < (\widehat{M}_{40})^{(8)}$ it follows $\frac{dG_{41}}{dt} \leq \left((\widehat{M}_{40})^{(8)}\right)_1 - (a'_{41})^{(8)}G_{41}$ and by integrating

$G_{41} \leq \left((\widehat{M}_{40})^{(8)}\right)_2 = G_{41}^0 + 2(a_{41})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_1/(a'_{41})^{(8)}$

In the same way , one can obtain

$G_{42} \leq \left((\widehat{M}_{40})^{(8)}\right)_3 = G_{42}^0 + 2(a_{42})^{(8)}\left((\widehat{M}_{40})^{(8)}\right)_2/(a'_{42})^{(8)}$

If $G_{41}\ or\ G_{42}$ is bounded, the same property follows for G_{40} , G_{42} and G_{40} , G_{41} respectively.

Remark 39: If $G_{40}\ is$ bounded, from below, the same property holds for $G_{41}\ and\ G_{42}$. The proof is analogous with the preceding one. An analogous property is true if G_{41} is bounded from below. 277

Remark 40: If T_{40} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(8)}((G_{43})(t),t)) = (b'_{41})^{(8)}$ then $T_{41} \to \infty$. 278

Definition of $(m)^{(8)}$ and ε_8 :

Indeed let t_8 be so that for $t > t_8$

$(b_{41})^{(8)} - (b_i'')^{(8)}\left((G_{43})(t),t\right) < \varepsilon_8, T_{40}(t) > (m)^{(8)}$

Then $\frac{dT_{41}}{dt} \geq (a_{41})^{(8)}(m)^{(8)} - \varepsilon_8 T_{41}$ which leads to 279

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{\varepsilon_8}\right)(1 - e^{-\varepsilon_8 t}) + T_{41}^0 e^{-\varepsilon_8 t}$ If we take t such that $e^{-\varepsilon_8 t} = \frac{1}{2}$ it results

$T_{41} \geq \left(\frac{(a_{41})^{(8)}(m)^{(8)}}{2}\right),\ t = log\frac{2}{\varepsilon_8}$ By taking now ε_8 sufficiently small one sees that T_{41} is unbounded. The same property holds for T_{42} if $\lim_{t\to\infty}(b''_{42})^{(8)}\left((G_{43})(t),t(t),t\right) = (b'_{42})^{(8)}$

It is now sufficient to take $\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}, \frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}} < 1$ and to choose $(\widehat{P}_{44})^{(9)}$ and $(\widehat{Q}_{44})^{(9)}$ large to have 279 A

$$\frac{(a_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[(\widehat{P}_{44})^{(9)} + \left((\widehat{P}_{44})^{(9)} + G_j^0\right)e^{-\left(\frac{(\widehat{P}_{44})^{(9)}+G_j^0}{G_j^0}\right)}\right] \leq (\widehat{P}_{44})^{(9)}$$

$$\frac{(b_i)^{(9)}}{(\widehat{M}_{44})^{(9)}}\left[\left((\widehat{Q}_{44})^{(9)} + T_j^0\right)e^{-\left(\frac{(\widehat{Q}_{44})^{(9)}+T_j^0}{T_j^0}\right)} + (\widehat{Q}_{44})^{(9)}\right] \leq (\widehat{Q}_{44})^{(9)}$$

In order that the operator $\mathcal{A}^{(9)}$ transforms the space of sextuples of functions G_i , T_i satisfying 39,35,36 into itself
The operator $\mathcal{A}^{(9)}$ is a contraction with respect to the metric

$$d\left(((G_{47})^{(1)},(T_{47})^{(1)}),((G_{47})^{(2)},(T_{47})^{(2)})\right)=$$
$$\sup_i\{\max_{t\in\mathbb{R}_+}\left|G_i^{(1)}(t)-G_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t},\max_{t\in\mathbb{R}_+}\left|T_i^{(1)}(t)-T_i^{(2)}(t)\right|e^{-(\widehat{M}_{44})^{(9)}t}\}$$

Indeed if we denote

Definition of $(\widetilde{G_{47}}),(\widetilde{T_{47}}):\left((\widetilde{G_{47}}),(\widetilde{T_{47}})\right)=\mathcal{A}^{(9)}\left((G_{47}),(T_{47})\right)$
It results

$$\left|\tilde{G}_{44}^{(1)}-\tilde{G}_i^{(2)}\right|\le\int_0^t(a_{44})^{(9)}\left|G_{45}^{(1)}-G_{45}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\,ds_{(44)}+$$
$$\int_0^t\{(a'_{44})^{(9)}\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)\left|G_{44}^{(1)}-G_{44}^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}+$$
$$G_{44}^{(2)}\left|(a''_{44})^{(9)}\left(T_{45}^{(1)},s_{(44)}\right)-(a''_{44})^{(9)}\left(T_{45}^{(2)},s_{(44)}\right)\right|\ e^{-(\widehat{M}_{44})^{(9)}s_{(44)}}e^{(\widehat{M}_{44})^{(9)}s_{(44)}}\}ds_{(44)}$$

Where $s_{(44)}$ represents integrand that is integrated over the interval $[0,t]$

From the hypotheses on 45,46,47,28 and 29 it follows

$$\left|(G_{47})^{(1)}-G^{(2)}\right|e^{-(\widehat{M}_{44})^{(9)}t}\le$$
$$\frac{1}{(\widehat{M}_{44})^{(9)}}\left((a_{44})^{(9)}+(a'_{44})^{(9)}+(\widehat{A}_{44})^{(9)}+(\widehat{P}_{44})^{(9)}(\widehat{k}_{44})^{(9)}\right)d\left(((G_{47})^{(1)},(T_{47})^{(1)};\ (G_{47})^{(2)},(T_{47})^{(2)})\right)$$

And analogous inequalities for $G_i\ and\ T_i$. Taking into account the hypothesis (39,35,36) the result follows

Remark 41: The fact that we supposed $(a''_{44})^{(9)}$ and $(b''_{44})^{(9)}$ depending also on t can be considered as not conformal with the reality, however we have put this hypothesis ,in order that we can postulate condition necessary to prove the uniqueness of the solution bounded by $(\widehat{P}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}\ and\ (\widehat{Q}_{44})^{(9)}e^{(\widehat{M}_{44})^{(9)}t}$ respectively of $\mathbb{R}_+$.

If instead of proving the existence of the solution on $\mathbb{R}_+$, we have to prove it only on a compact then it suffices to consider that $(a_i'')^{(9)}$ and $(b_i'')^{(9)}, i=44{,}45{,}46$ depend only on T_{45} and respectively on $(G_{47})(and\ not\ on\ \ t)$ and hypothesis can replaced by a usual Lipschitz condition.

Remark 42: There does not exist any t where $G_i\,(t)=0\ and\ T_i\,(t)=0$

From 99 to 44 it results

$$G_i\,(t)\ge G_i^0e^{\left[-\int_0^t\{(a_i')^{(9)}-(a_i'')^{(9)}(T_{45}(s_{(44)}),s_{(44)})\}ds_{(44)}\right]}\ge 0$$

$$T_i\,(t)\ge T_i^0e^{\left(-(b_i')^{(9)}t\right)}>0\quad \text{for t}>0$$

Definition of $\left((\widehat{M}_{44})^{(9)}\right)_1,\left((\widehat{M}_{44})^{(9)}\right)_2\ and\ \left((\widehat{M}_{44})^{(9)}\right)_3$:

Remark 43: if G_{44} is bounded, the same property have also $G_{45}\ and\ G_{46}$. indeed if

$G_{44} < (\widehat{M}_{44})^{(9)}$ it follows $\frac{dG_{45}}{dt} \leq \left((\widehat{M}_{44})^{(9)}\right)_1 - (a'_{45})^{(9)} G_{45}$ and by integrating

$$G_{45} \leq \left((\widehat{M}_{44})^{(9)}\right)_2 = G_{45}^0 + 2(a_{45})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_1/(a'_{45})^{(9)}$$

In the same way , one can obtain

$$G_{46} \leq \left((\widehat{M}_{44})^{(9)}\right)_3 = G_{46}^0 + 2(a_{46})^{(9)}\left((\widehat{M}_{44})^{(9)}\right)_2/(a'_{46})^{(9)}$$

If G_{45} *or* G_{46} is bounded, the same property follows for G_{44} , G_{46} and G_{44} , G_{45} respectively.

Remark 44: If G_{44} *is* bounded, from below, the same property holds for G_{45} *and* G_{46} . The proof is analogous with the preceding one. An analogous property is true if G_{45} is bounded from below.

Remark 45: If T_{44} is bounded from below and $\lim_{t\to\infty}((b_i'')^{(9)}\left((G_{47})(t),t\right) = (b'_{45})^{(9)}$ then $T_{45} \to \infty$.

Definition of $(m)^{(9)}$ and ε_9 :

Indeed let t_9 be so that for $t > t_9$

$$(b_{45})^{(9)} - (b_i'')^{(9)}\left((G_{47})(t),t\right) < \varepsilon_9, T_{44}(t) > (m)^{(9)}$$

Then $\frac{dT_{45}}{dt} \geq (a_{45})^{(9)}(m)^{(9)} - \varepsilon_9 T_{45}$ which leads to

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{\varepsilon_9}\right)(1 - e^{-\varepsilon_9 t}) + T_{45}^0 e^{-\varepsilon_9 t}$ If we take t such that $e^{-\varepsilon_9 t} = \frac{1}{2}$ it results

$T_{45} \geq \left(\frac{(a_{45})^{(9)}(m)^{(9)}}{2}\right)$, $t = log\frac{2}{\varepsilon_9}$ By taking now ε_9 sufficiently small one sees that T_{45} is unbounded. The same property holds for T_{46} if $\lim_{t\to\infty}(b''_{46})^{(9)}\left((G_{47})(t),t\right) = (b'_{46})^{(9)}$
We now state a more precise theorem about the behaviors at infinity of the solutions of equations 37 to 92

Behavior of the solutions of equation 280

Theorem If we denote and define

Definition of $(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$:

$(\sigma_1)^{(1)}, (\sigma_2)^{(1)}, (\tau_1)^{(1)}, (\tau_2)^{(1)}$ four constants satisfying
$-(\sigma_2)^{(1)} \leq -(a'_{13})^{(1)} + (a'_{14})^{(1)} - (a''_{13})^{(1)}(T_{14},t) + (a''_{14})^{(1)}(T_{14},t) \leq -(\sigma_1)^{(1)}$

$-(\tau_2)^{(1)} \leq -(b'_{13})^{(1)} + (b'_{14})^{(1)} - (b''_{13})^{(1)}(G,t) - (b''_{14})^{(1)}(G,t) \leq -(\tau_1)^{(1)}$

Definition of $(v_1)^{(1)}, (v_2)^{(1)}, (u_1)^{(1)}, (u_2)^{(1)}, v^{(1)}, u^{(1)}$: 281

By $(v_1)^{(1)} > 0$, $(v_2)^{(1)} < 0$ and respectively $(u_1)^{(1)} > 0$, $(u_2)^{(1)} < 0$ the roots of the equations $(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_1)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_1)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(\bar{v}_1)^{(1)}, , (\bar{v}_2)^{(1)}, (\bar{u}_1)^{(1)}, (\bar{u}_2)^{(1)}$: 282

By $(\bar{v}_1)^{(1)} > 0$, $(\bar{v}_2)^{(1)} < 0$ and respectively $(\bar{u}_1)^{(1)} > 0$, $(\bar{u}_2)^{(1)} < 0$ the roots of the equations

$(a_{14})^{(1)}\left(v^{(1)}\right)^2 + (\sigma_2)^{(1)}v^{(1)} - (a_{13})^{(1)} = 0$ and $(b_{14})^{(1)}\left(u^{(1)}\right)^2 + (\tau_2)^{(1)}u^{(1)} - (b_{13})^{(1)} = 0$

Definition of $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}, (\nu_0)^{(1)}$:- 283

If we define $(m_1)^{(1)}, (m_2)^{(1)}, (\mu_1)^{(1)}, (\mu_2)^{(1)}$ by

$(m_2)^{(1)} = (\nu_0)^{(1)}, (m_1)^{(1)} = (\nu_1)^{(1)},\ if\ (\nu_0)^{(1)} < (\nu_1)^{(1)}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\bar{\nu}_1)^{(1)}, if\ (\nu_1)^{(1)} < (\nu_0)^{(1)} < (\bar{\nu}_1)^{(1)},$

and $\boxed{(\nu_0)^{(1)} = \frac{G_{13}^0}{G_{14}^0}}$

$(m_2)^{(1)} = (\nu_1)^{(1)}, (m_1)^{(1)} = (\nu_0)^{(1)},\ if\ (\bar{\nu}_1)^{(1)} < (\nu_0)^{(1)}$

and analogously 284

$(\mu_2)^{(1)} = (u_0)^{(1)}, (\mu_1)^{(1)} = (u_1)^{(1)},\ if\ (u_0)^{(1)} < (u_1)^{(1)}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (\bar{u}_1)^{(1)}, if\ (u_1)^{(1)} < (u_0)^{(1)} < (\bar{u}_1)^{(1)},$

and $\boxed{(u_0)^{(1)} = \frac{T_{13}^0}{T_{14}^0}}$

$(\mu_2)^{(1)} = (u_1)^{(1)}, (\mu_1)^{(1)} = (u_0)^{(1)}, if\ (\bar{u}_1)^{(1)} < (u_0)^{(1)}$ where $(u_1)^{(1)}, (\bar{u}_1)^{(1)}$

are defined

Then the solution of global equations satisfies the inequalities 285

$$G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{13}(t) \le G_{13}^0 e^{(S_1)^{(1)}t}$$

where $(p_i)^{(1)}$ is defined by equation

$$\frac{1}{(m_1)^{(1)}} G_{13}^0 e^{((S_1)^{(1)} - (p_{13})^{(1)})t} \le G_{14}(t) \le \frac{1}{(m_2)^{(1)}} G_{13}^0 e^{(S_1)^{(1)}t}$$

$$\left(\frac{(a_{15})^{(1)} G_{13}^0}{(m_1)^{(1)}((S_1)^{(1)} - (p_{13})^{(1)} - (S_2)^{(1)})}\left[e^{((S_1)^{(1)} - (p_{13})^{(1)})t} - e^{-(S_2)^{(1)}t}\right] + G_{15}^0 e^{-(S_2)^{(1)}t} \le G_{15}(t) \le \right. \quad 286$$

$$\left. \frac{(a_{15})^{(1)} G_{13}^0}{(m_2)^{(1)}((S_1)^{(1)} - (a'_{15})^{(1)})}\left[e^{(S_1)^{(1)}t} - e^{-(a'_{15})^{(1)}t}\right] + G_{15}^0 e^{-(a'_{15})^{(1)}t}\right)$$

$$\boxed{T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t}} \quad 287$$

$$\frac{1}{(\mu_1)^{(1)}} T_{13}^0 e^{(R_1)^{(1)}t} \le T_{13}(t) \le \frac{1}{(\mu_2)^{(1)}} T_{13}^0 e^{((R_1)^{(1)} + (r_{13})^{(1)})t} \quad 288$$

$$\frac{(b_{15})^{(1)} T_{13}^0}{(\mu_1)^{(1)}((R_1)^{(1)} - (b'_{15})^{(1)})}\left[e^{(R_1)^{(1)}t} - e^{-(b'_{15})^{(1)}t}\right] + T_{15}^0 e^{-(b'_{15})^{(1)}t} \le T_{15}(t) \le \quad 289$$

$$\frac{(a_{15})^{(1)} T_{13}^0}{(\mu_2)^{(1)}((R_1)^{(1)} + (r_{13})^{(1)} + (R_2)^{(1)})}\left[e^{((R_1)^{(1)} + (r_{13})^{(1)})t} - e^{-(R_2)^{(1)}t}\right] + T_{15}^0 e^{-(R_2)^{(1)}t}$$

Definition of $(S_1)^{(1)}, (S_2)^{(1)}, (R_1)^{(1)}, (R_2)^{(1)}$:- 290

Where $(S_1)^{(1)} = (a_{13})^{(1)}(m_2)^{(1)} - (a'_{13})^{(1)}$

$$(S_2)^{(1)} = (a_{15})^{(1)} - (p_{15})^{(1)}$$

$$(R_1)^{(1)} = (b_{13})^{(1)}(\mu_2)^{(1)} - (b'_{13})^{(1)}$$

$$(R_2)^{(1)} = (b'_{15})^{(1)} - (r_{15})^{(1)}$$

Behavior of the solutions of equation 291

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$: 292

$(\sigma_1)^{(2)}, (\sigma_2)^{(2)}, (\tau_1)^{(2)}, (\tau_2)^{(2)}$ four constants satisfying

$-(\sigma_2)^{(2)} \le -(a'_{16})^{(2)} + (a'_{17})^{(2)} - (a''_{16})^{(2)}(\mathrm{T}_{17}, t) + (a''_{17})^{(2)}(\mathrm{T}_{17}, t) \le -(\sigma_1)^{(2)}$ 293

$-(\tau_2)^{(2)} \le -(b'_{16})^{(2)} + (b'_{17})^{(2)} - (b''_{16})^{(2)}\big((G_{19}), t\big) - (b''_{17})^{(2)}\big((G_{19}), t\big) \le -(\tau_1)^{(2)}$ 294

Definition of $(\nu_1)^{(2)}, (\nu_2)^{(2)}, (u_1)^{(2)}, (u_2)^{(2)}$: 295

By $(\nu_1)^{(2)} > 0$, $(\nu_2)^{(2)} < 0$ and respectively $(u_1)^{(2)} > 0$, $(u_2)^{(2)} < 0$ the roots 296

of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 297

and $(b_{14})^{(2)}\big(u^{(2)}\big)^2 + (\tau_1)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ and 298

Definition of $(\bar{\nu}_1)^{(2)}, , (\bar{\nu}_2)^{(2)}, (\bar{u}_1)^{(2)}, (\bar{u}_2)^{(2)}$: 299

By $(\bar{\nu}_1)^{(2)} > 0$, $(\bar{\nu}_2)^{(2)} < 0$ and respectively $(\bar{u}_1)^{(2)} > 0$, $(\bar{u}_2)^{(2)} < 0$ the 300

roots of the equations $(a_{17})^{(2)}\big(\nu^{(2)}\big)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)} = 0$ 301

and $(b_{17})^{(2)}\big(u^{(2)}\big)^2 + (\tau_2)^{(2)}u^{(2)} - (b_{16})^{(2)} = 0$ 302

Definition of $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$:- 303

If we define $(m_1)^{(2)}, (m_2)^{(2)}, (\mu_1)^{(2)}, (\mu_2)^{(2)}$ by 304

$(m_2)^{(2)} = (\nu_0)^{(2)}, (m_1)^{(2)} = (\nu_1)^{(2)}$, ***if*** $(\nu_0)^{(2)} < (\nu_1)^{(2)}$ 305

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\bar{\nu}_1)^{(2)}$, ***if*** $(\nu_1)^{(2)} < (\nu_0)^{(2)} < (\bar{\nu}_1)^{(2)}$, 306

and $\boxed{(\nu_0)^{(2)} = \frac{\mathrm{G}^0_{16}}{\mathrm{G}^0_{17}}}$

$(m_2)^{(2)} = (\nu_1)^{(2)}, (m_1)^{(2)} = (\nu_0)^{(2)}$, ***if*** $(\bar{\nu}_1)^{(2)} < (\nu_0)^{(2)}$ 307

and analogously 308

$(\mu_2)^{(2)} = (u_0)^{(2)}, (\mu_1)^{(2)} = (u_1)^{(2)}$, ***if*** $(u_0)^{(2)} < (u_1)^{(2)}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (\bar{u}_1)^{(2)}$, ***if*** $(u_1)^{(2)} < (u_0)^{(2)} < (\bar{u}_1)^{(2)}$,

and $\boxed{(u_0)^{(2)} = \frac{\mathrm{T}^0_{16}}{\mathrm{T}^0_{17}}}$

$(\mu_2)^{(2)} = (u_1)^{(2)}, (\mu_1)^{(2)} = (u_0)^{(2)},\ \boldsymbol{if}\ (\bar{u}_1)^{(2)} < (u_0)^{(2)}$ 309

Then the solution of global equations satisfies the inequalities 310

$$G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{16}(t) \le G_{16}^0 e^{(S_1)^{(2)} t}$$

$(p_i)^{(2)}$ is defined by equation

$$\frac{1}{(m_1)^{(2)}} G_{16}^0 e^{((S_1)^{(2)} - (p_{16})^{(2)})t} \le G_{17}(t) \le \frac{1}{(m_2)^{(2)}} G_{16}^0 e^{(S_1)^{(2)} t} \quad 311$$

$$\left(\frac{(a_{18})^{(2)} G_{16}^0}{(m_1)^{(2)}((S_1)^{(2)} - (p_{16})^{(2)} - (S_2)^{(2)})}\left[e^{((S_1)^{(2)} - (p_{16})^{(2)})t} - e^{-(S_2)^{(2)} t}\right] + G_{18}^0 e^{-(S_2)^{(2)} t} \le G_{18}(t) \le \right. \quad 312$$

$$\left. \frac{(a_{18})^{(2)} G_{16}^0}{(m_2)^{(2)}((S_1)^{(2)} - (a'_{18})^{(2)})}\left[e^{(S_1)^{(2)} t} - e^{-(a'_{18})^{(2)} t}\right] + G_{18}^0 e^{-(a'_{18})^{(2)} t}\right)$$

$$\boxed{T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t}} \quad 313$$

$$\frac{1}{(\mu_1)^{(2)}} T_{16}^0 e^{(R_1)^{(2)} t} \le T_{16}(t) \le \frac{1}{(\mu_2)^{(2)}} T_{16}^0 e^{((R_1)^{(2)} + (r_{16})^{(2)})t} \quad 314$$

$$\frac{(b_{18})^{(2)} T_{16}^0}{(\mu_1)^{(2)}((R_1)^{(2)} - (b'_{18})^{(2)})}\left[e^{(R_1)^{(2)} t} - e^{-(b'_{18})^{(2)} t}\right] + T_{18}^0 e^{-(b'_{18})^{(2)} t} \le T_{18}(t) \le \quad 315$$

$$\frac{(a_{18})^{(2)} T_{16}^0}{(\mu_2)^{(2)}((R_1)^{(2)} + (r_{16})^{(2)} + (R_2)^{(2)})}\left[e^{((R_1)^{(2)} + (r_{16})^{(2)})t} - e^{-(R_2)^{(2)} t}\right] + T_{18}^0 e^{-(R_2)^{(2)} t}$$

Definition of $(S_1)^{(2)}, (S_2)^{(2)}, (R_1)^{(2)}, (R_2)^{(2)}$:- 316

Where $(S_1)^{(2)} = (a_{16})^{(2)}(m_2)^{(2)} - (a'_{16})^{(2)}$ 317

$$(S_2)^{(2)} = (a_{18})^{(2)} - (p_{18})^{(2)}$$

$$(R_1)^{(2)} = (b_{16})^{(2)}(\mu_2)^{(1)} - (b'_{16})^{(2)} \quad 318$$

$$(R_2)^{(2)} = (b'_{18})^{(2)} - (r_{18})^{(2)}$$

Behavior of the solutions 319

Theorem 3: If we denote and define

Definition of $(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$:

$(\sigma_1)^{(3)}, (\sigma_2)^{(3)}, (\tau_1)^{(3)}, (\tau_2)^{(3)}$ four constants satisfying

$$-(\sigma_2)^{(3)} \le -(a'_{20})^{(3)} + (a'_{21})^{(3)} - (a''_{20})^{(3)}(T_{21}, t) + (a''_{21})^{(3)}(T_{21}, t) \le -(\sigma_1)^{(3)}$$

$$-(\tau_2)^{(3)} \le -(b'_{20})^{(3)} + (b'_{21})^{(3)} - (b''_{20})^{(3)}(G_{23}, t) - (b''_{21})^{(3)}\big((G_{23}), t\big) \le -(\tau_1)^{(3)}$$

Definition of $(v_1)^{(3)}, (v_2)^{(3)}, (u_1)^{(3)}, (u_2)^{(3)}$: 320

By $(v_1)^{(3)} > 0$, $(v_2)^{(3)} < 0$ and respectively $(u_1)^{(3)} > 0$, $(u_2)^{(3)} < 0$ the roots of the equations
$(a_{21})^{(3)}\big(v^{(3)}\big)^2 + (\sigma_1)^{(3)} v^{(3)} - (a_{20})^{(3)} = 0$
and $(b_{21})^{(3)}\big(u^{(3)}\big)^2 + (\tau_1)^{(3)} u^{(3)} - (b_{20})^{(3)} = 0$ and

By $(\bar{v}_1)^{(3)} > 0$, $(\bar{v}_2)^{(3)} < 0$ and respectively $(\bar{u}_1)^{(3)} > 0$, $(\bar{u}_2)^{(3)} < 0$ the

roots of the equations $(a_{21})^{(3)}\left(v^{(3)}\right)^2 + (\sigma_2)^{(3)}v^{(3)} - (a_{20})^{(3)} = 0$

and $(b_{21})^{(3)}\left(u^{(3)}\right)^2 + (\tau_2)^{(3)}u^{(3)} - (b_{20})^{(3)} = 0$

Definition of $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$:- 321

If we define $(m_1)^{(3)}$, $(m_2)^{(3)}$, $(\mu_1)^{(3)}$, $(\mu_2)^{(3)}$ by

$(m_2)^{(3)} = (v_0)^{(3)}, (m_1)^{(3)} = (v_1)^{(3)}$, $\boldsymbol{if}$ $(v_0)^{(3)} < (v_1)^{(3)}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (\bar{v}_1)^{(3)}$, $\boldsymbol{if}$ $(v_1)^{(3)} < (v_0)^{(3)} < (\bar{v}_1)^{(3)}$,

and $\boxed{(v_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}}$

$(m_2)^{(3)} = (v_1)^{(3)}, (m_1)^{(3)} = (v_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{v}_1)^{(3)} < (v_0)^{(3)}$

and analogously 322

$(\mu_2)^{(3)} = (u_0)^{(3)}, (\mu_1)^{(3)} = (u_1)^{(3)}$, $\boldsymbol{if}$ $(u_0)^{(3)} < (u_1)^{(3)}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (\bar{u}_1)^{(3)}$, $\boldsymbol{if}$ $(u_1)^{(3)} < (u_0)^{(3)} < (\bar{u}_1)^{(3)}$, and $\boxed{(u_0)^{(3)} = \frac{T_{20}^0}{T_{21}^0}}$

$(\mu_2)^{(3)} = (u_1)^{(3)}, (\mu_1)^{(3)} = (u_0)^{(3)}$, $\boldsymbol{if}$ $(\bar{u}_1)^{(3)} < (u_0)^{(3)}$

Then the solution of global equations satisfies the inequalities

$G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{20}(t) \le G_{20}^0 e^{(S_1)^{(3)}t}$

$(p_i)^{(3)}$ is defined by equation

$\frac{1}{(m_1)^{(3)}} G_{20}^0 e^{((S_1)^{(3)} - (p_{20})^{(3)})t} \le G_{21}(t) \le \frac{1}{(m_2)^{(3)}} G_{20}^0 e^{(S_1)^{(3)}t}$ 323

$\left(\frac{(a_{22})^{(3)} G_{20}^0}{(m_1)^{(3)}((S_1)^{(3)} - (p_{20})^{(3)} - (S_2)^{(3)})}\left[e^{((S_1)^{(3)} - (p_{20})^{(3)})t} - e^{-(S_2)^{(3)}t}\right] + G_{22}^0 e^{-(S_2)^{(3)}t} \le G_{22}(t) \le\right.$ 324

$\left.\frac{(a_{22})^{(3)} G_{20}^0}{(m_2)^{(3)}((S_1)^{(3)} - (a'_{22})^{(3)})}\left[e^{(S_1)^{(3)}t} - e^{-(a'_{22})^{(3)}t}\right] + G_{22}^0 e^{-(a'_{22})^{(3)}t}\right)$

$\boxed{T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}}$ 325

$\frac{1}{(\mu_1)^{(3)}} T_{20}^0 e^{(R_1)^{(3)}t} \le T_{20}(t) \le \frac{1}{(\mu_2)^{(3)}} T_{20}^0 e^{((R_1)^{(3)} + (r_{20})^{(3)})t}$ 326

$\frac{(b_{22})^{(3)} T_{20}^0}{(\mu_1)^{(3)}((R_1)^{(3)} - (b'_{22})^{(3)})}\left[e^{(R_1)^{(3)}t} - e^{-(b'_{22})^{(3)}t}\right] + T_{22}^0 e^{-(b'_{22})^{(3)}t} \le T_{22}(t) \le$ 327

$\frac{(a_{22})^{(3)} T_{20}^0}{(\mu_2)^{(3)}((R_1)^{(3)} + (r_{20})^{(3)} + (R_2)^{(3)})}\left[e^{((R_1)^{(3)} + (r_{20})^{(3)})t} - e^{-(R_2)^{(3)}t}\right] + T_{22}^0 e^{-(R_2)^{(3)}t}$

Definition of $(S_1)^{(3)}, (S_2)^{(3)}, (R_1)^{(3)}, (R_2)^{(3)}$:- 328

Where $(S_1)^{(3)} = (a_{20})^{(3)}(m_2)^{(3)} - (a'_{20})^{(3)}$

$$(S_2)^{(3)} = (a_{22})^{(3)} - (p_{22})^{(3)}$$

$$(R_1)^{(3)} = (b_{20})^{(3)}(\mu_2)^{(3)} - (b'_{20})^{(3)}$$

$$(R_2)^{(3)} = (b'_{22})^{(3)} - (r_{22})^{(3)}$$

Behavior of the solutions of equation

Theorem: If we denote and define

Definition of $(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$:

$(\sigma_1)^{(4)}, (\sigma_2)^{(4)}, (\tau_1)^{(4)}, (\tau_2)^{(4)}$ four constants satisfying

$$-(\sigma_2)^{(4)} \le -(a'_{24})^{(4)} + (a'_{25})^{(4)} - (a''_{24})^{(4)}(T_{25}, t) + (a''_{25})^{(4)}(T_{25}, t) \le -(\sigma_1)^{(4)}$$

$$-(\tau_2)^{(4)} \le -(b'_{24})^{(4)} + (b'_{25})^{(4)} - (b''_{24})^{(4)}\big((G_{27}), t\big) - (b''_{25})^{(4)}\big((G_{27}), t\big) \le -(\tau_1)^{(4)}$$

Definition of $(\nu_1)^{(4)}, (\nu_2)^{(4)}, (u_1)^{(4)}, (u_2)^{(4)}, \nu^{(4)}, u^{(4)}$: 329

By $(\nu_1)^{(4)} > 0$, $(\nu_2)^{(4)} < 0$ and respectively $(u_1)^{(4)} > 0$, $(u_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_1)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_1)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$ and

Definition of $(\bar{\nu}_1)^{(4)}, , (\bar{\nu}_2)^{(4)}, (\bar{u}_1)^{(4)}, (\bar{u}_2)^{(4)}$: 330

By $(\bar{\nu}_1)^{(4)} > 0$, $(\bar{\nu}_2)^{(4)} < 0$ and respectively $(\bar{u}_1)^{(4)} > 0$, $(\bar{u}_2)^{(4)} < 0$ the roots of the equations $(a_{25})^{(4)}\big(\nu^{(4)}\big)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)} = 0$
and $(b_{25})^{(4)}\big(u^{(4)}\big)^2 + (\tau_2)^{(4)}u^{(4)} - (b_{24})^{(4)} = 0$

Definition of $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}, (\nu_0)^{(4)}$:-

If we define $(m_1)^{(4)}, (m_2)^{(4)}, (\mu_1)^{(4)}, (\mu_2)^{(4)}$ by

$$(m_2)^{(4)} = (\nu_0)^{(4)}, (m_1)^{(4)} = (\nu_1)^{(4)}, \ \boldsymbol{if}\ (\nu_0)^{(4)} < (\nu_1)^{(4)}$$

$$(m_2)^{(4)} = (\nu_1)^{(4)}, (m_1)^{(4)} = (\bar{\nu}_1)^{(4)}, \boldsymbol{if}\ (\nu_4)^{(4)} < (\nu_0)^{(4)} < (\bar{\nu}_1)^{(4)},$$

and $\boxed{(\nu_0)^{(4)} = \frac{G^0_{24}}{G^0_{25}}}$

$$(m_2)^{(4)} = (\nu_4)^{(4)}, (m_1)^{(4)} = (\nu_0)^{(4)}, \ \boldsymbol{if}\ (\bar{\nu}_4)^{(4)} < (\nu_0)^{(4)}$$

and analogously 331

$$(\mu_2)^{(4)} = (u_0)^{(4)}, (\mu_1)^{(4)} = (u_1)^{(4)}, \ \boldsymbol{if}\ (u_0)^{(4)} < (u_1)^{(4)}$$

$$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (\bar{u}_1)^{(4)}, \boldsymbol{if}\ (u_1)^{(4)} < (u_0)^{(4)} < (\bar{u}_1)^{(4)},$$

and $\boxed{(u_0)^{(4)} = \frac{T^0_{24}}{T^0_{25}}}$

$(\mu_2)^{(4)} = (u_1)^{(4)}, (\mu_1)^{(4)} = (u_0)^{(4)}, \boldsymbol{if}\ (\bar{u}_1)^{(4)} < (u_0)^{(4)}$ where $(u_1)^{(4)}, (\bar{u}_1)^{(4)}$

Then the solution of global equations satisfies the inequalities

332

$$G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{24}(t) \le G_{24}^0 e^{(S_1)^{(4)} t}$$

where $(p_i)^{(4)}$ is defined by equation

$$\frac{1}{(m_1)^{(4)}} G_{24}^0 e^{((S_1)^{(4)} - (p_{24})^{(4)})t} \le G_{25}(t) \le \frac{1}{(m_2)^{(4)}} G_{24}^0 e^{(S_1)^{(4)} t} \quad 333$$

$$\left(\frac{(a_{26})^{(4)} G_{24}^0}{(m_1)^{(4)}((S_1)^{(4)} - (p_{24})^{(4)} - (S_2)^{(4)})} \left[e^{((S_1)^{(4)} - (p_{24})^{(4)})t} - e^{-(S_2)^{(4)} t} \right] + G_{26}^0 e^{-(S_2)^{(4)} t} \le G_{26}(t) \le \right. \quad 334$$

$$\left. \frac{(a_{26})^{(4)} G_{24}^0}{(m_2)^{(4)}((S_1)^{(4)} - (a'_{26})^{(4)})} \left[e^{(S_1)^{(4)} t} - e^{-(a'_{26})^{(4)} t} \right] + G_{26}^0 e^{-(a'_{26})^{(4)} t} \right)$$

$$\boxed{T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t}}$$

$$\frac{1}{(\mu_1)^{(4)}} T_{24}^0 e^{(R_1)^{(4)} t} \le T_{24}(t) \le \frac{1}{(\mu_2)^{(4)}} T_{24}^0 e^{((R_1)^{(4)} + (r_{24})^{(4)})t} \quad 335$$

$$\frac{(b_{26})^{(4)} T_{24}^0}{(\mu_1)^{(4)}((R_1)^{(4)} - (b'_{26})^{(4)})} \left[e^{(R_1)^{(4)} t} - e^{-(b'_{26})^{(4)} t} \right] + T_{26}^0 e^{-(b'_{26})^{(4)} t} \le T_{26}(t) \le \quad 336$$

$$\frac{(a_{26})^{(4)} T_{24}^0}{(\mu_2)^{(4)}((R_1)^{(4)} + (r_{24})^{(4)} + (R_2)^{(4)})} \left[e^{((R_1)^{(4)} + (r_{24})^{(4)})t} - e^{-(R_2)^{(4)} t} \right] + T_{26}^0 e^{-(R_2)^{(4)} t}$$

Definition of $(S_1)^{(4)}, (S_2)^{(4)}, (R_1)^{(4)}, (R_2)^{(4)}$:- 337

Where $(S_1)^{(4)} = (a_{24})^{(4)}(m_2)^{(4)} - (a'_{24})^{(4)}$

$(S_2)^{(4)} = (a_{26})^{(4)} - (p_{26})^{(4)}$

$(R_1)^{(4)} = (b_{24})^{(4)}(\mu_2)^{(4)} - (b'_{24})^{(4)}$

$(R_2)^{(4)} = (b'_{26})^{(4)} - (r_{26})^{(4)}$

Behavior of the solutions of equation 338

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$:

$(\sigma_1)^{(5)}, (\sigma_2)^{(5)}, (\tau_1)^{(5)}, (\tau_2)^{(5)}$ four constants satisfying

$$-(\sigma_2)^{(5)} \le -(a'_{28})^{(5)} + (a'_{29})^{(5)} - (a''_{28})^{(5)}(T_{29}, t) + (a''_{29})^{(5)}(T_{29}, t) \le -(\sigma_1)^{(5)}$$

$$-(\tau_2)^{(5)} \le -(b'_{28})^{(5)} + (b'_{29})^{(5)} - (b''_{28})^{(5)}((G_{31}), t) - (b''_{29})^{(5)}((G_{31}), t) \le -(\tau_1)^{(5)}$$

Definition of $(v_1)^{(5)}, (v_2)^{(5)}, (u_1)^{(5)}, (u_2)^{(5)}, v^{(5)}, u^{(5)}$: 339

By $(v_1)^{(5)} > 0$, $(v_2)^{(5)} < 0$ and respectively $(u_1)^{(5)} > 0$, $(u_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_1)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_1)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$ and

Definition of $(\bar{v}_1)^{(5)},,(\bar{v}_2)^{(5)},(\bar{u}_1)^{(5)},(\bar{u}_2)^{(5)}$: 340

By $(\bar{v}_1)^{(5)} > 0$, $(\bar{v}_2)^{(5)} < 0$ and respectively $(\bar{u}_1)^{(5)} > 0$, $(\bar{u}_2)^{(5)} < 0$ the roots of the equations $(a_{29})^{(5)}\left(v^{(5)}\right)^2 + (\sigma_2)^{(5)}v^{(5)} - (a_{28})^{(5)} = 0$

and $(b_{29})^{(5)}\left(u^{(5)}\right)^2 + (\tau_2)^{(5)}u^{(5)} - (b_{28})^{(5)} = 0$

Definition of $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$, $(v_0)^{(5)}$:-

If we define $(m_1)^{(5)}$, $(m_2)^{(5)}$, $(\mu_1)^{(5)}$, $(\mu_2)^{(5)}$ by

$(m_2)^{(5)} = (v_0)^{(5)}, (m_1)^{(5)} = (v_1)^{(5)}$, $\boldsymbol{if}\ (v_0)^{(5)} < (v_1)^{(5)}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (\bar{v}_1)^{(5)}$, $\boldsymbol{if}\ (v_1)^{(5)} < (v_0)^{(5)} < (\bar{v}_1)^{(5)}$,

and $\boxed{(v_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$

$(m_2)^{(5)} = (v_1)^{(5)}, (m_1)^{(5)} = (v_0)^{(5)}$, $\boldsymbol{if}\ (\bar{v}_1)^{(5)} < (v_0)^{(5)}$

and analogously 341

$(\mu_2)^{(5)} = (u_0)^{(5)}, (\mu_1)^{(5)} = (u_1)^{(5)}$, $\boldsymbol{if}\ (u_0)^{(5)} < (u_1)^{(5)}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (\bar{u}_1)^{(5)}$, $\boldsymbol{if}\ (u_1)^{(5)} < (u_0)^{(5)} < (\bar{u}_1)^{(5)}$,

and $\boxed{(u_0)^{(5)} = \frac{T_{28}^0}{T_{29}^0}}$

$(\mu_2)^{(5)} = (u_1)^{(5)}, (\mu_1)^{(5)} = (u_0)^{(5)}, \boldsymbol{if}\ (\bar{u}_1)^{(5)} < (u_0)^{(5)}$ where $(u_1)^{(5)}, (\bar{u}_1)^{(5)}$

Then the solution of global equations satisfies the inequalities 342

$$G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{28}(t) \le G_{28}^0 e^{(S_1)^{(5)}t}$$

where $(p_i)^{(5)}$ is defined by equation

$$\frac{1}{(m_5)^{(5)}} G_{28}^0 e^{((S_1)^{(5)} - (p_{28})^{(5)})t} \le G_{29}(t) \le \frac{1}{(m_2)^{(5)}} G_{28}^0 e^{(S_1)^{(5)}t} \quad 343$$

$$\left(\frac{(a_{30})^{(5)} G_{28}^0}{(m_1)^{(5)}((S_1)^{(5)} - (p_{28})^{(5)} - (S_2)^{(5)})}\left[e^{((S_1)^{(5)} - (p_{28})^{(5)})t} - e^{-(S_2)^{(5)}t}\right] + G_{30}^0 e^{-(S_2)^{(5)}t} \le G_{30}(t) \le \right. \quad 344$$

$$\left. \frac{(a_{30})^{(5)} G_{28}^0}{(m_2)^{(5)}((S_1)^{(5)} - (a'_{30})^{(5)})}\left[e^{(S_1)^{(5)}t} - e^{-(a'_{30})^{(5)}t}\right] + G_{30}^0 e^{-(a'_{30})^{(5)}t}\right)$$

$$\boxed{T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le T_{28}^0 e^{((R_1)^{(5)} + (r_{28})^{(5)})t}} \quad 345$$

$$\frac{1}{(\mu_1)^{(5)}}T_{28}^0 e^{(R_1)^{(5)}t} \le T_{28}(t) \le \frac{1}{(\mu_2)^{(5)}}T_{28}^0 e^{((R_1)^{(5)}+(r_{28})^{(5)})t} \quad 346$$

$$\frac{(b_{30})^{(5)}T_{28}^0}{(\mu_1)^{(5)}((R_1)^{(5)}-(b'_{30})^{(5)})}\left[e^{(R_1)^{(5)}t} - e^{-(b'_{30})^{(5)}t}\right] + T_{30}^0 e^{-(b'_{30})^{(5)}t} \le T_{30}(t) \le \quad 347$$

$$\frac{(a_{30})^{(5)}T_{28}^0}{(\mu_2)^{(5)}((R_1)^{(5)}+(r_{28})^{(5)}+(R_2)^{(5)})}\left[e^{((R_1)^{(5)}+(r_{28})^{(5)})t} - e^{-(R_2)^{(5)}t}\right] + T_{30}^0 e^{-(R_2)^{(5)}t}$$

Definition of $(S_1)^{(5)}, (S_2)^{(5)}, (R_1)^{(5)}, (R_2)^{(5)}$:- 348

Where $(S_1)^{(5)} = (a_{28})^{(5)}(m_2)^{(5)} - (a'_{28})^{(5)}$

$(S_2)^{(5)} = (a_{30})^{(5)} - (p_{30})^{(5)}$

$(R_1)^{(5)} = (b_{28})^{(5)}(\mu_2)^{(5)} - (b'_{28})^{(5)}$

$(R_2)^{(5)} = (b'_{30})^{(5)} - (r_{30})^{(5)}$

Behavior of the solutions of equation 349

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$:

$(\sigma_1)^{(6)}, (\sigma_2)^{(6)}, (\tau_1)^{(6)}, (\tau_2)^{(6)}$ four constants satisfying

$$-(\sigma_2)^{(6)} \le -(a'_{32})^{(6)} + (a'_{33})^{(6)} - (a''_{32})^{(6)}(T_{33}, t) + (a''_{33})^{(6)}(T_{33}, t) \le -(\sigma_1)^{(6)}$$

$$-(\tau_2)^{(6)} \le -(b'_{32})^{(6)} + (b'_{33})^{(6)} - (b''_{32})^{(6)}\big((G_{35}), t\big) - (b''_{33})^{(6)}\big((G_{35}), t\big) \le -(\tau_1)^{(6)}$$

Definition of $(v_1)^{(6)}, (v_2)^{(6)}, (u_1)^{(6)}, (u_2)^{(6)}, v^{(6)}, u^{(6)}$: 350

By $(v_1)^{(6)} > 0$, $(v_2)^{(6)} < 0$ and respectively $(u_1)^{(6)} > 0$, $(u_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_1)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_1)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$ and

Definition of $(\bar{v}_1)^{(6)},, (\bar{v}_2)^{(6)}, (\bar{u}_1)^{(6)}, (\bar{u}_2)^{(6)}$: 351

By $(\bar{v}_1)^{(6)} > 0$, $(\bar{v}_2)^{(6)} < 0$ and respectively $(\bar{u}_1)^{(6)} > 0$, $(\bar{u}_2)^{(6)} < 0$ the roots of the equations $(a_{33})^{(6)}\big(v^{(6)}\big)^2 + (\sigma_2)^{(6)}v^{(6)} - (a_{32})^{(6)} = 0$
and $(b_{33})^{(6)}\big(u^{(6)}\big)^2 + (\tau_2)^{(6)}u^{(6)} - (b_{32})^{(6)} = 0$

Definition of $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}, (v_0)^{(6)}$:-

If we define $(m_1)^{(6)}, (m_2)^{(6)}, (\mu_1)^{(6)}, (\mu_2)^{(6)}$ by

$(m_2)^{(6)} = (v_0)^{(6)}, (m_1)^{(6)} = (v_1)^{(6)}$, $\boldsymbol{if}$ $(v_0)^{(6)} < (v_1)^{(6)}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (\bar{v}_6)^{(6)}$, $\boldsymbol{if}$ $(v_1)^{(6)} < (v_0)^{(6)} < (\bar{v}_1)^{(6)}$,

and $\boxed{(v_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$

$(m_2)^{(6)} = (v_1)^{(6)}, (m_1)^{(6)} = (v_0)^{(6)},\ \boldsymbol{if}\ (\bar{v}_1)^{(6)} < (v_0)^{(6)}$

and analogously 352

$(\mu_2)^{(6)} = (u_0)^{(6)}, (\mu_1)^{(6)} = (u_1)^{(6)},\ \boldsymbol{if}\ (u_0)^{(6)} < (u_1)^{(6)}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (\bar{u}_1)^{(6)}, \boldsymbol{if}\ (u_1)^{(6)} < (u_0)^{(6)} < (\bar{u}_1)^{(6)},$

and $\boxed{(u_0)^{(6)} = \frac{T_{32}^0}{T_{33}^0}}$

$(\mu_2)^{(6)} = (u_1)^{(6)}, (\mu_1)^{(6)} = (u_0)^{(6)}, \boldsymbol{if}\ (\bar{u}_1)^{(6)} < (u_0)^{(6)}$ where $(u_1)^{(6)}, (\bar{u}_1)^{(6)}$

Then the solution of global equations satisfies the inequalities 353

$$G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{32}(t) \le G_{32}^0 e^{(S_1)^{(6)} t}$$

where $(p_i)^{(6)}$ is defined by equation

$$\frac{1}{(m_1)^{(6)}} G_{32}^0 e^{((S_1)^{(6)} - (p_{32})^{(6)})t} \le G_{33}(t) \le \frac{1}{(m_2)^{(6)}} G_{32}^0 e^{(S_1)^{(6)} t} \quad 354$$

$$\left(\frac{(a_{34})^{(6)} G_{32}^0}{(m_1)^{(6)}((S_1)^{(6)} - (p_{32})^{(6)} - (S_2)^{(6)})}\left[e^{((S_1)^{(6)} - (p_{32})^{(6)})t} - e^{-(S_2)^{(6)} t}\right] + G_{34}^0 e^{-(S_2)^{(6)} t} \le G_{34}(t) \le \right. \quad 355$$

$$\left. \frac{(a_{34})^{(6)} G_{32}^0}{(m_2)^{(6)}((S_1)^{(6)} - (a'_{34})^{(6)})}\left[e^{(S_1)^{(6)} t} - e^{-(a'_{34})^{(6)} t}\right] + G_{34}^0 e^{-(a'_{34})^{(6)} t}\right)$$

$$\boxed{T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t}} \quad 356$$

$$\frac{1}{(\mu_1)^{(6)}} T_{32}^0 e^{(R_1)^{(6)} t} \le T_{32}(t) \le \frac{1}{(\mu_2)^{(6)}} T_{32}^0 e^{((R_1)^{(6)} + (r_{32})^{(6)})t} \quad 357$$

$$\frac{(b_{34})^{(6)} T_{32}^0}{(\mu_1)^{(6)}((R_1)^{(6)} - (b'_{34})^{(6)})}\left[e^{(R_1)^{(6)} t} - e^{-(b'_{34})^{(6)} t}\right] + T_{34}^0 e^{-(b'_{34})^{(6)} t} \le T_{34}(t) \le \quad 358$$

$$\frac{(a_{34})^{(6)} T_{32}^0}{(\mu_2)^{(6)}((R_1)^{(6)} + (r_{32})^{(6)} + (R_2)^{(6)})}\left[e^{((R_1)^{(6)} + (r_{32})^{(6)})t} - e^{-(R_2)^{(6)} t}\right] + T_{34}^0 e^{-(R_2)^{(6)} t}$$

Definition of $(S_1)^{(6)}, (S_2)^{(6)}, (R_1)^{(6)}, (R_2)^{(6)}$:- 359

Where $(S_1)^{(6)} = (a_{32})^{(6)}(m_2)^{(6)} - (a'_{32})^{(6)}$

$(S_2)^{(6)} = (a_{34})^{(6)} - (p_{34})^{(6)}$

$(R_1)^{(6)} = (b_{32})^{(6)}(\mu_2)^{(6)} - (b'_{32})^{(6)}$

$(R_2)^{(6)} = (b'_{34})^{(6)} - (r_{34})^{(6)}$

Behavior of the solutions of equation

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$:

$(\sigma_1)^{(7)}, (\sigma_2)^{(7)}, (\tau_1)^{(7)}, (\tau_2)^{(7)}$ four constants satisfying

$$-(\sigma_2)^{(7)} \le -(a'_{36})^{(7)} + (a'_{37})^{(7)} - (a''_{36})^{(7)}(T_{37}, t) + (a''_{37})^{(7)}(T_{37}, t) \le -(\sigma_1)^{(7)}$$

$$-(\tau_2)^{(7)} \le -(b'_{36})^{(7)} + (b'_{37})^{(7)} - (b''_{36})^{(7)}\big((G_{39}), t\big) - (b''_{37})^{(7)}\big((G_{39}), t\big) \le -(\tau_1)^{(7)}$$

Definition of $(\nu_1)^{(7)}, (\nu_2)^{(7)}, (u_1)^{(7)}, (u_2)^{(7)}, \nu^{(7)}, u^{(7)}$: 361

By $(\nu_1)^{(7)} > 0, (\nu_2)^{(7)} < 0$ and respectively $(u_1)^{(7)} > 0, (u_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_1)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_1)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$ and

Definition of $(\bar{\nu}_1)^{(7)},, (\bar{\nu}_2)^{(7)}, (\bar{u}_1)^{(7)}, (\bar{u}_2)^{(7)}$: 362

By $(\bar{\nu}_1)^{(7)} > 0, (\bar{\nu}_2)^{(7)} < 0$ and respectively $(\bar{u}_1)^{(7)} > 0, (\bar{u}_2)^{(7)} < 0$ the roots of the equations $(a_{37})^{(7)}\big(\nu^{(7)}\big)^2 + (\sigma_2)^{(7)}\nu^{(7)} - (a_{36})^{(7)} = 0$
and $(b_{37})^{(7)}\big(u^{(7)}\big)^2 + (\tau_2)^{(7)}u^{(7)} - (b_{36})^{(7)} = 0$

Definition of $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}, (\nu_0)^{(7)}$:-

If we define $(m_1)^{(7)}, (m_2)^{(7)}, (\mu_1)^{(7)}, (\mu_2)^{(7)}$ by

$$(m_2)^{(7)} = (\nu_0)^{(7)}, (m_1)^{(7)} = (\nu_1)^{(7)}, \ \boldsymbol{if}\ (\nu_0)^{(7)} < (\nu_1)^{(7)}$$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\bar{\nu}_1)^{(7)}, \boldsymbol{if}\ (\nu_1)^{(7)} < (\nu_0)^{(7)} < (\bar{\nu}_1)^{(7)},$$

and $\boxed{(\nu_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$

$$(m_2)^{(7)} = (\nu_1)^{(7)}, (m_1)^{(7)} = (\nu_0)^{(7)}, \ \boldsymbol{if}\ (\bar{\nu}_1)^{(7)} < (\nu_0)^{(7)}$$

and analogously 363

$$(\mu_2)^{(7)} = (u_0)^{(7)}, (\mu_1)^{(7)} = (u_1)^{(7)}, \ \boldsymbol{if}\ (u_0)^{(7)} < (u_1)^{(7)}$$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (\bar{u}_1)^{(7)}, \boldsymbol{if}\ (u_1)^{(7)} < (u_0)^{(7)} < (\bar{u}_1)^{(7)},$$

and $\boxed{(u_0)^{(7)} = \frac{T^0_{36}}{T^0_{37}}}$

$$(\mu_2)^{(7)} = (u_1)^{(7)}, (\mu_1)^{(7)} = (u_0)^{(7)}, \boldsymbol{if}\ (\bar{u}_1)^{(7)} < (u_0)^{(7)} \text{ where } (u_1)^{(7)}, (\bar{u}_1)^{(7)}$$

Then the solution of global equations satisfies the inequalities 364

$$G^0_{36} e^{\left((S_1)^{(7)} - (p_{36})^{(7)}\right)t} \le G_{36}(t) \le G^0_{36} e^{(S_1)^{(7)}t}$$

where $(p_i)^{(7)}$ is defined by equation

$$\frac{1}{(m_7)^{(7)}} G_{36}^0 e^{((S_1)^{(7)}-(p_{36})^{(7)})t} \le G_{37}(t) \le \frac{1}{(m_2)^{(7)}} G_{36}^0 e^{(S_1)^{(7)}t} \quad 365$$

$$\left(\frac{(a_{38})^{(7)} G_{36}^0}{(m_1)^{(7)}((S_1)^{(7)}-(p_{36})^{(7)}-(S_2)^{(7)})}\left[e^{((S_1)^{(7)}-(p_{36})^{(7)})t} - e^{-(S_2)^{(7)}t}\right] + G_{38}^0 e^{-(S_2)^{(7)}t} \le G_{38}(t) \le \right. \quad 366$$

$$\left. \frac{(a_{38})^{(7)} G_{36}^0}{(m_2)^{(7)}((S_1)^{(7)}-(a'_{38})^{(7)})}\left[e^{(S_1)^{(7)}t} - e^{-(a'_{38})^{(7)}t}\right] + G_{38}^0 e^{-(a'_{38})^{(7)}t}\right)$$

$$\boxed{T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t}} \quad 367$$

$$\frac{1}{(\mu_1)^{(7)}} T_{36}^0 e^{(R_1)^{(7)}t} \le T_{36}(t) \le \frac{1}{(\mu_2)^{(7)}} T_{36}^0 e^{((R_1)^{(7)}+(r_{36})^{(7)})t} \quad 368$$

$$\frac{(b_{38})^{(7)} T_{36}^0}{(\mu_1)^{(7)}((R_1)^{(7)}-(b'_{38})^{(7)})}\left[e^{(R_1)^{(7)}t} - e^{-(b'_{38})^{(7)}t}\right] + T_{38}^0 e^{-(b'_{38})^{(7)}t} \le T_{38}(t) \le \quad 369$$

$$\frac{(a_{38})^{(7)} T_{36}^0}{(\mu_2)^{(7)}((R_1)^{(7)}+(r_{36})^{(7)}+(R_2)^{(7)})}\left[e^{((R_1)^{(7)}+(r_{36})^{(7)})t} - e^{-(R_2)^{(7)}t}\right] + T_{38}^0 e^{-(R_2)^{(7)}t}$$

Definition of $(S_1)^{(7)}, (S_2)^{(7)}, (R_1)^{(7)}, (R_2)^{(7)}$:- 370

Where $(S_1)^{(7)} = (a_{36})^{(7)}(m_2)^{(7)} - (a'_{36})^{(7)}$

$(S_2)^{(7)} = (a_{38})^{(7)} - (p_{38})^{(7)}$

$(R_1)^{(7)} = (b_{36})^{(7)}(\mu_2)^{(7)} - (b'_{36})^{(7)}$

$(R_2)^{(7)} = (b'_{38})^{(7)} - (r_{38})^{(7)}$

Behavior of the solutions of equation 371

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$:

$(\sigma_1)^{(8)}, (\sigma_2)^{(8)}, (\tau_1)^{(8)}, (\tau_2)^{(8)}$ four constants satisfying

$$-(\sigma_2)^{(8)} \le -(a'_{40})^{(8)} + (a'_{41})^{(8)} - (a''_{40})^{(8)}(T_{41}, t) + (a''_{41})^{(8)}(T_{41}, t) \le -(\sigma_1)^{(8)}$$

$$-(\tau_2)^{(8)} \le -(b'_{40})^{(8)} + (b'_{41})^{(8)} - (b''_{40})^{(8)}\big((G_{43}), t\big) - (b''_{41})^{(8)}\big((G_{43}), t\big) \le -(\tau_1)^{(8)}$$

Definition of $(\nu_1)^{(8)}, (\nu_2)^{(8)}, (u_1)^{(8)}, (u_2)^{(8)}, \nu^{(8)}, u^{(8)}$: 372

By $(\nu_1)^{(8)} > 0, (\nu_2)^{(8)} < 0$ and respectively $(u_1)^{(8)} > 0, (u_2)^{(8)} < 0$ the roots of the equations $(a_{41})^{(8)}\big(\nu^{(8)}\big)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)} = 0$
and $(b_{41})^{(8)}\big(u^{(8)}\big)^2 + (\tau_1)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$ and

Definition of $(\bar{\nu}_1)^{(8)}, , (\bar{\nu}_2)^{(8)}, (\bar{u}_1)^{(8)}, (\bar{u}_2)^{(8)}$:

By $(\bar{\nu}_1)^{(8)} > 0, (\bar{\nu}_2)^{(8)} < 0$ and respectively $(\bar{u}_1)^{(8)} > 0, (\bar{u}_2)^{(8)} < 0$ the

roots of the equations $(a_{41})^{(8)}\left(v^{(8)}\right)^2 + (\sigma_2)^{(8)}v^{(8)} - (a_{40})^{(8)} = 0$

and $(b_{41})^{(8)}\left(u^{(8)}\right)^2 + (\tau_2)^{(8)}u^{(8)} - (b_{40})^{(8)} = 0$

Definition of $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}, (v_0)^{(8)}$:-

If we define $(m_1)^{(8)}, (m_2)^{(8)}, (\mu_1)^{(8)}, (\mu_2)^{(8)}$ by

$(m_2)^{(8)} = (v_0)^{(8)}, (m_1)^{(8)} = (v_1)^{(8)}$, ***if*** $(v_0)^{(8)} < (v_1)^{(8)}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (\bar{v}_1)^{(8)}$, ***if*** $(v_1)^{(8)} < (v_0)^{(8)} < (\bar{v}_1)^{(8)}$,

and $\boxed{(v_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$

$(m_2)^{(8)} = (v_1)^{(8)}, (m_1)^{(8)} = (v_0)^{(8)}$, ***if*** $(\bar{v}_1)^{(8)} < (v_0)^{(8)}$

and analogously 374

$(\mu_2)^{(8)} = (u_0)^{(8)}, (\mu_1)^{(8)} = (u_1)^{(8)}$, ***if*** $(u_0)^{(8)} < (u_1)^{(8)}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (\bar{u}_1)^{(8)}$, ***if*** $(u_1)^{(8)} < (u_0)^{(8)} < (\bar{u}_1)^{(8)}$,

and $\boxed{(u_0)^{(8)} = \frac{T_{40}^0}{T_{41}^0}}$

$(\mu_2)^{(8)} = (u_1)^{(8)}, (\mu_1)^{(8)} = (u_0)^{(8)}$, ***if*** $(\bar{u}_1)^{(8)} < (u_0)^{(8)}$ where $(u_1)^{(8)}, (\bar{u}_1)^{(8)}$

Then the solution of global equations satisfies the inequalities 375

$$G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{40}(t) \le G_{40}^0 e^{(S_1)^{(8)}t}$$

where $(p_i)^{(8)}$ is defined by equation

$$\frac{1}{(m_1)^{(8)}} G_{40}^0 e^{((S_1)^{(8)} - (p_{40})^{(8)})t} \le G_{41}(t) \le \frac{1}{(m_2)^{(8)}} G_{40}^0 e^{(S_1)^{(8)}t} \quad 376$$

$$\left(\frac{(a_{42})^{(8)} G_{40}^0}{(m_1)^{(8)}((S_1)^{(8)} - (p_{40})^{(8)} - (S_2)^{(8)})}\left[e^{((S_1)^{(8)} - (p_{40})^{(8)})t} - e^{-(S_2)^{(8)}t}\right] + G_{42}^0 e^{-(S_2)^{(8)}t} \le G_{42}(t) \le \right. \quad 377$$

$$\left. \frac{(a_{42})^{(8)} G_{40}^0}{(m_2)^{(8)}((S_1)^{(8)} - (a'_{42})^{(8)})}\left[e^{(S_1)^{(8)}t} - e^{-(a'_{42})^{(8)}t}\right] + G_{42}^0 e^{-(a'_{42})^{(8)}t}\right)$$

$$\boxed{T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t}} \quad 378$$

$$\frac{1}{(\mu_1)^{(8)}} T_{40}^0 e^{(R_1)^{(8)}t} \le T_{40}(t) \le \frac{1}{(\mu_2)^{(8)}} T_{40}^0 e^{((R_1)^{(8)} + (r_{40})^{(8)})t} \quad 379$$

$$\frac{(b_{42})^{(8)} T_{40}^0}{(\mu_1)^{(8)}((R_1)^{(8)} - (b'_{42})^{(8)})}\left[e^{(R_1)^{(8)}t} - e^{-(b'_{42})^{(8)}t}\right] + T_{42}^0 e^{-(b'_{42})^{(8)}t} \le T_{42}(t) \le \quad 380$$

$$\frac{(a_{42})^{(8)} T_{40}^0}{(\mu_2)^{(8)}((R_1)^{(8)} + (r_{40})^{(8)} + (R_2)^{(8)})}\left[e^{((R_1)^{(8)} + (r_{40})^{(8)})t} - e^{-(R_2)^{(8)}t}\right] + T_{42}^0 e^{-(R_2)^{(8)}t}$$

Definition of $(S_1)^{(8)}, (S_2)^{(8)}, (R_1)^{(8)}, (R_2)^{(8)}$:- 381

Where $(S_1)^{(8)} = (a_{40})^{(8)}(m_2)^{(8)} - (a'_{40})^{(8)}$

$$(S_2)^{(8)} = (a_{42})^{(8)} - (p_{42})^{(8)}$$

$$(R_1)^{(8)} = (b_{40})^{(8)}(\mu_2)^{(8)} - (b'_{40})^{(8)}$$

$$(R_2)^{(8)} = (b'_{42})^{(8)} - (r_{42})^{(8)}$$

Behavior of the solutions of equation 37 to 92 382

Theorem 2: If we denote and define

Definition of $(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$:

$(\sigma_1)^{(9)}, (\sigma_2)^{(9)}, (\tau_1)^{(9)}, (\tau_2)^{(9)}$ four constants satisfying

$$-(\sigma_2)^{(9)} \le -(a'_{44})^{(9)} + (a'_{45})^{(9)} - (a''_{44})^{(9)}(T_{45}, t) + (a''_{45})^{(9)}(T_{45}, t) \le -(\sigma_1)^{(9)}$$

$$-(\tau_2)^{(9)} \le -(b'_{44})^{(9)} + (b'_{45})^{(9)} - (b''_{44})^{(9)}\big((G_{47}), t\big) - (b''_{45})^{(9)}\big((G_{47}), t\big) \le -(\tau_1)^{(9)}$$

Definition of $(v_1)^{(9)}, (v_2)^{(9)}, (u_1)^{(9)}, (u_2)^{(9)}, v^{(9)}, u^{(9)}$:

By $(v_1)^{(9)} > 0$, $(v_2)^{(9)} < 0$ and respectively $(u_1)^{(9)} > 0$, $(u_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_1)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_1)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$ and

Definition of $(\bar{v}_1)^{(9)}, , (\bar{v}_2)^{(9)}, (\bar{u}_1)^{(9)}, (\bar{u}_2)^{(9)}$:

By $(\bar{v}_1)^{(9)} > 0$, $(\bar{v}_2)^{(9)} < 0$ and respectively $(\bar{u}_1)^{(9)} > 0$, $(\bar{u}_2)^{(9)} < 0$ the roots of the equations $(a_{45})^{(9)}\big(v^{(9)}\big)^2 + (\sigma_2)^{(9)}v^{(9)} - (a_{44})^{(9)} = 0$
and $(b_{45})^{(9)}\big(u^{(9)}\big)^2 + (\tau_2)^{(9)}u^{(9)} - (b_{44})^{(9)} = 0$

Definition of $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}, (v_0)^{(9)}$:-

If we define $(m_1)^{(9)}, (m_2)^{(9)}, (\mu_1)^{(9)}, (\mu_2)^{(9)}$ by

$$(m_2)^{(9)} = (v_0)^{(9)}, (m_1)^{(9)} = (v_1)^{(9)}, \ \boldsymbol{if}\ (v_0)^{(9)} < (v_1)^{(9)}$$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (\bar{v}_1)^{(9)}, \boldsymbol{if}\ (v_1)^{(9)} < (v_0)^{(9)} < (\bar{v}_1)^{(9)},$$

and $\boxed{(v_0)^{(9)} = \frac{G^0_{44}}{G^0_{45}}}$

$$(m_2)^{(9)} = (v_1)^{(9)}, (m_1)^{(9)} = (v_0)^{(9)}, \ \boldsymbol{if}\ (\bar{v}_1)^{(9)} < (v_0)^{(9)}$$

and analogously

$$(\mu_2)^{(9)} = (u_0)^{(9)}, (\mu_1)^{(9)} = (u_1)^{(9)}, \ \boldsymbol{if}\ (u_0)^{(9)} < (u_1)^{(9)}$$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (\bar{u}_1)^{(9)}$, $\boldsymbol{if}$ $(u_1)^{(9)} < (u_0)^{(9)} < (\bar{u}_1)^{(9)}$,

and $\boxed{(u_0)^{(9)} = \frac{T_{44}^0}{T_{45}^0}}$

$(\mu_2)^{(9)} = (u_1)^{(9)}, (\mu_1)^{(9)} = (u_0)^{(9)}, \boldsymbol{if}$ $(\bar{u}_1)^{(9)} < (u_0)^{(9)}$ where $(u_1)^{(9)}, (\bar{u}_1)^{(9)}$ are defined by 59 and 69 respectively

Then the solution of 19,20,21,22,23 and 24 satisfies the inequalities

$$G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{44}(t) \leq G_{44}^0 e^{(S_1)^{(9)} t}$$

where $(p_i)^{(9)}$ is defined by equation 45

$$\frac{1}{(m_9)^{(9)}} G_{44}^0 e^{((S_1)^{(9)} - (p_{44})^{(9)})t} \leq G_{45}(t) \leq \frac{1}{(m_2)^{(9)}} G_{44}^0 e^{(S_1)^{(9)} t}$$

$$\left(\frac{(a_{46})^{(9)} G_{44}^0}{(m_1)^{(9)}((S_1)^{(9)} - (p_{44})^{(9)} - (S_2)^{(9)})}\left[e^{((S_1)^{(9)} - (p_{44})^{(9)})t} - e^{-(S_2)^{(9)} t}\right] + G_{46}^0 e^{-(S_2)^{(9)} t} \leq G_{46}(t) \leq \right.$$

$$\left.\frac{(a_{46})^{(9)} G_{44}^0}{(m_2)^{(9)}((S_1)^{(9)} - (a'_{46})^{(9)})}\left[e^{(S_1)^{(9)} t} - e^{-(a'_{46})^{(9)} t}\right] + G_{46}^0 e^{-(a'_{46})^{(9)} t}\right)$$

$$\boxed{T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}}$$

$$\frac{1}{(\mu_1)^{(9)}} T_{44}^0 e^{(R_1)^{(9)} t} \leq T_{44}(t) \leq \frac{1}{(\mu_2)^{(9)}} T_{44}^0 e^{((R_1)^{(9)} + (r_{44})^{(9)})t}$$

$$\frac{(b_{46})^{(9)} T_{44}^0}{(\mu_1)^{(9)}((R_1)^{(9)} - (b'_{46})^{(9)})}\left[e^{(R_1)^{(9)} t} - e^{-(b'_{46})^{(9)} t}\right] + T_{46}^0 e^{-(b'_{46})^{(9)} t} \leq T_{46}(t) \leq$$

$$\frac{(a_{46})^{(9)} T_{44}^0}{(\mu_2)^{(9)}((R_1)^{(9)} + (r_{44})^{(9)} + (R_2)^{(9)})}\left[e^{((R_1)^{(9)} + (r_{44})^{(9)})t} - e^{-(R_2)^{(9)} t}\right] + T_{46}^0 e^{-(R_2)^{(9)} t}$$

Definition of $(S_1)^{(9)}, (S_2)^{(9)}, (R_1)^{(9)}, (R_2)^{(9)}$:-

Where $(S_1)^{(9)} = (a_{44})^{(9)}(m_2)^{(9)} - (a'_{44})^{(9)}$

$(S_2)^{(9)} = (a_{46})^{(9)} - (p_{46})^{(9)}$

$(R_1)^{(9)} = (b_{44})^{(9)}(\mu_2)^{(9)} - (b'_{44})^{(9)}$

$(R_2)^{(9)} = (b'_{46})^{(9)} - (r_{46})^{(9)}$

Proof : From global equations we obtain 383

$$\frac{dv^{(1)}}{dt} = (a_{13})^{(1)} - \left((a'_{13})^{(1)} - (a'_{14})^{(1)} + (a''_{13})^{(1)}(T_{14}, t)\right) - (a''_{14})^{(1)}(T_{14}, t)v^{(1)} - (a_{14})^{(1)}v^{(1)}$$

Definition of $v^{(1)}$:- $\boxed{v^{(1)} = \frac{G_{13}}{G_{14}}}$

It follows

$$-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_2)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)\leq\frac{dv^{(1)}}{dt}\leq-\left((a_{14})^{(1)}\left(v^{(1)}\right)^2+(\sigma_1)^{(1)}v^{(1)}-(a_{13})^{(1)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(1)},(\nu_0)^{(1)}$:-

For $0<\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}<(\nu_1)^{(1)}<(\bar{\nu}_1)^{(1)}$

$$v^{(1)}(t)\geq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_0)^{(1)}\right)t\right]}}\quad,\quad\boxed{(C)^{(1)}=\frac{(\nu_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\nu_2)^{(1)}}}$$

$$\text{it follows } (\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\nu_1)^{(1)}$$

In the same manner , we get 384

$$v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(1)}=\frac{(\bar{\nu}_1)^{(1)}-(\nu_0)^{(1)}}{(\nu_0)^{(1)}-(\bar{\nu}_2)^{(1)}}}$$

From which we deduce $(\nu_0)^{(1)}\leq v^{(1)}(t)\leq(\bar{\nu}_1)^{(1)}$

If $0<(\nu_1)^{(1)}<(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}<(\bar{\nu}_1)^{(1)}$ we find like in the previous case, 385

$$(\nu_1)^{(1)}\leq\frac{(\nu_1)^{(1)}+(C)^{(1)}(\nu_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}{1+(C)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\nu_1)^{(1)}-(\nu_2)^{(1)}\right)t\right]}}\leq v^{(1)}(t)\leq$$

$$\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\bar{\nu}_1)^{(1)}$$

If $0<(\nu_1)^{(1)}\leq(\bar{\nu}_1)^{(1)}\leq\boxed{(\nu_0)^{(1)}=\frac{G_{13}^0}{G_{14}^0}}$, we obtain 386

$$(\nu_1)^{(1)}\leq v^{(1)}(t)\leq\frac{(\bar{\nu}_1)^{(1)}+(\bar{C})^{(1)}(\bar{\nu}_2)^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}{1+(\bar{C})^{(1)}e^{\left[-(a_{14})^{(1)}\left((\bar{\nu}_1)^{(1)}-(\bar{\nu}_2)^{(1)}\right)t\right]}}\leq(\nu_0)^{(1)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(1)}(t)$:-

$$(m_2)^{(1)}\leq v^{(1)}(t)\leq(m_1)^{(1)},\quad\boxed{v^{(1)}(t)=\frac{G_{13}(t)}{G_{14}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(1)}(t)$:-

$$(\mu_2)^{(1)}\leq u^{(1)}(t)\leq(\mu_1)^{(1)},\quad\boxed{u^{(1)}(t)=\frac{T_{13}(t)}{T_{14}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the

theorem.

Particular case :

If $(a''_{13})^{(1)} = (a''_{14})^{(1)}, then\ (\sigma_1)^{(1)} = (\sigma_2)^{(1)}$ and in this case $(\nu_1)^{(1)} = (\bar{\nu}_1)^{(1)}$ if in addition $(\nu_0)^{(1)} = (\nu_1)^{(1)}$ then $\nu^{(1)}(t) = (\nu_0)^{(1)}$ and as a consequence $G_{13}(t) = (\nu_0)^{(1)} G_{14}(t)$ this also defines $(\nu_0)^{(1)}$ for the special case

Analogously if $(b''_{13})^{(1)} = (b''_{14})^{(1)}, then\ (\tau_1)^{(1)} = (\tau_2)^{(1)}$ and then

$(u_1)^{(1)} = (\bar{u}_1)^{(1)}$ if in addition $(u_0)^{(1)} = (u_1)^{(1)}$ then $T_{13}(t) = (u_0)^{(1)} T_{14}(t)$ This is an important consequence of the relation between $(\nu_1)^{(1)}$ and $(\bar{\nu}_1)^{(1)}$, and definition of $(u_0)^{(1)}$.

Proof : From global equations we obtain 387

$$\frac{d\nu^{(2)}}{dt} = (a_{16})^{(2)} - \left((a'_{16})^{(2)} - (a'_{17})^{(2)} + (a''_{16})^{(2)}(T_{17}, t)\right) - (a''_{17})^{(2)}(T_{17}, t)\nu^{(2)} - (a_{17})^{(2)}\nu^{(2)}$$

Definition of $\nu^{(2)}$:- $\boxed{\nu^{(2)} = \frac{G_{16}}{G_{17}}}$ 388

It follows 389

$$-\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_2)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right) \le \frac{d\nu^{(2)}}{dt} \le -\left((a_{17})^{(2)}\left(\nu^{(2)}\right)^2 + (\sigma_1)^{(2)}\nu^{(2)} - (a_{16})^{(2)}\right)$$

From which one obtains 390

Definition of $(\bar{\nu}_1)^{(2)}, (\nu_0)^{(2)}$:-

For $0 < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\nu_1)^{(2)} < (\bar{\nu}_1)^{(2)}$

$$\nu^{(2)}(t) \ge \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_0)^{(2)}\right)t\right]}} \quad , \quad \boxed{(C)^{(2)} = \frac{(\nu_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\nu_2)^{(2)}}}$$

it follows $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\nu_1)^{(2)}$

In the same manner , we get 391

$$\nu^{(2)}(t) \le \frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(2)} = \frac{(\bar{\nu}_1)^{(2)} - (\nu_0)^{(2)}}{(\nu_0)^{(2)} - (\bar{\nu}_2)^{(2)}}}$$

From which we deduce $(\nu_0)^{(2)} \le \nu^{(2)}(t) \le (\bar{\nu}_1)^{(2)}$ 392

If $0 < (\nu_1)^{(2)} < (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}} < (\bar{\nu}_1)^{(2)}$ we find like in the previous case, 393

$$(\nu_1)^{(2)} \le \frac{(\nu_1)^{(2)} + (C)^{(2)}(\nu_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}}{1 + (C)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\nu_1)^{(2)} - (\nu_2)^{(2)}\right)t\right]}} \le \nu^{(2)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(2)} + (\bar{C})^{(2)}(\bar{\nu}_2)^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}}{1 + (\bar{C})^{(2)} e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)} - (\bar{\nu}_2)^{(2)}\right)t\right]}} \le (\bar{\nu}_1)^{(2)}$$

If $0 < (\nu_1)^{(2)} \le (\bar{\nu}_1)^{(2)} \le (\nu_0)^{(2)} = \frac{G^0_{16}}{G^0_{17}}$, we obtain 394

$$(\nu_1)^{(2)} \leq \nu^{(2)}(t) \leq \frac{(\bar{\nu}_1)^{(2)}+(\bar{C})^{(2)}(\bar{\nu}_2)^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}}{1+(\bar{C})^{(2)}e^{\left[-(a_{17})^{(2)}\left((\bar{\nu}_1)^{(2)}-(\bar{\nu}_2)^{(2)}\right)t\right]}} \leq (\nu_0)^{(2)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(2)}(t)$:- 395

$$(m_2)^{(2)} \leq \nu^{(2)}(t) \leq (m_1)^{(2)}, \quad \boxed{\nu^{(2)}(t) = \frac{G_{16}(t)}{G_{17}(t)}}$$

In a completely analogous way, we obtain 396

Definition of $u^{(2)}(t)$:-

$$(\mu_2)^{(2)} \leq u^{(2)}(t) \leq (\mu_1)^{(2)}, \quad \boxed{u^{(2)}(t) = \frac{T_{16}(t)}{T_{17}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case : 397

If $(a''_{16})^{(2)} = (a''_{17})^{(2)}, then\ (\sigma_1)^{(2)} = (\sigma_2)^{(2)}$ and in this case $(\nu_1)^{(2)} = (\bar{\nu}_1)^{(2)}$ if in addition $(\nu_0)^{(2)} = (\nu_1)^{(2)}$ then $\nu^{(2)}(t) = (\nu_0)^{(2)}$ and as a consequence $G_{16}(t) = (\nu_0)^{(2)}G_{17}(t)$

Analogously if $(b''_{16})^{(2)} = (b''_{17})^{(2)}, then\ (\tau_1)^{(2)} = (\tau_2)^{(2)}$ and then

$(u_1)^{(2)} = (\bar{u}_1)^{(2)}$if in addition $(u_0)^{(2)} = (u_1)^{(2)}$ then $T_{16}(t) = (u_0)^{(2)}T_{17}(t)$ This is an important consequence of the relation between $(\nu_1)^{(2)}$ and $(\bar{\nu}_1)^{(2)}$

Proof : From global equations we obtain 398

$$\frac{d\nu^{(3)}}{dt} = (a_{20})^{(3)} - \left((a'_{20})^{(3)} - (a'_{21})^{(3)} + (a''_{20})^{(3)}(T_{21}, t)\right) - (a''_{21})^{(3)}(T_{21}, t)\nu^{(3)} - (a_{21})^{(3)}\nu^{(3)}$$

Definition of $\nu^{(3)}$:- $\boxed{\nu^{(3)} = \frac{G_{20}}{G_{21}}}$ 399

It follows

$$-\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_2)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right) \leq \frac{d\nu^{(3)}}{dt} \leq -\left((a_{21})^{(3)}\left(\nu^{(3)}\right)^2 + (\sigma_1)^{(3)}\nu^{(3)} - (a_{20})^{(3)}\right)$$

400

From which one obtains

For $0 < (\nu_0)^{(3)} = \frac{G^0_{20}}{G^0_{21}} < (\nu_1)^{(3)} < (\bar{\nu}_1)^{(3)}$

$$\nu^{(3)}(t) \geq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_0)^{(3)}\right)t\right]}} \quad , \quad \boxed{(C)^{(3)} = \frac{(\nu_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\nu_2)^{(3)}}}$$

it follows $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\nu_1)^{(3)}$

In the same manner , we get 401

$$\nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(3)} = \frac{(\bar{\nu}_1)^{(3)}-(\nu_0)^{(3)}}{(\nu_0)^{(3)}-(\bar{\nu}_2)^{(3)}}}$$

Definition of $(\bar{\nu}_1)^{(3)}$:-

From which we deduce $(\nu_0)^{(3)} \leq \nu^{(3)}(t) \leq (\bar{\nu}_1)^{(3)}$

If $0 < (\nu_1)^{(3)} < (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0} < (\bar{\nu}_1)^{(3)}$ we find like in the previous case, 402

$$(\nu_1)^{(3)} \leq \frac{(\nu_1)^{(3)}+(C)^{(3)}(\nu_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}}{1+(C)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\nu_1)^{(3)}-(\nu_2)^{(3)}\right)t\right]}} \leq \nu^{(3)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\bar{\nu}_1)^{(3)}$$

If $0 < (\nu_1)^{(3)} \leq (\bar{\nu}_1)^{(3)} \leq (\nu_0)^{(3)} = \frac{G_{20}^0}{G_{21}^0}$, we obtain 403

$$(\nu_1)^{(3)} \leq \nu^{(3)}(t) \leq \frac{(\bar{\nu}_1)^{(3)}+(\bar{C})^{(3)}(\bar{\nu}_2)^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}}{1+(\bar{C})^{(3)}e^{\left[-(a_{21})^{(3)}\left((\bar{\nu}_1)^{(3)}-(\bar{\nu}_2)^{(3)}\right)t\right]}} \leq (\nu_0)^{(3)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(3)}(t)$:-

$$(m_2)^{(3)} \leq \nu^{(3)}(t) \leq (m_1)^{(3)}, \quad \boxed{\nu^{(3)}(t) = \frac{G_{20}(t)}{G_{21}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(3)}(t)$:-

$$(\mu_2)^{(3)} \leq u^{(3)}(t) \leq (\mu_1)^{(3)}, \quad \boxed{u^{(3)}(t) = \frac{T_{20}(t)}{T_{21}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{20})^{(3)} = (a''_{21})^{(3)}, then\ (\sigma_1)^{(3)} = (\sigma_2)^{(3)}$ and in this case $(\nu_1)^{(3)} = (\bar{\nu}_1)^{(3)}$ if in addition $(\nu_0)^{(3)} = (\nu_1)^{(3)}$ then $\nu^{(3)}(t) = (\nu_0)^{(3)}$ and as a consequence $G_{20}(t) = (\nu_0)^{(3)}G_{21}(t)$

Analogously if $(b''_{20})^{(3)} = (b''_{21})^{(3)}, then\ (\tau_1)^{(3)} = (\tau_2)^{(3)}$ and then

$(u_1)^{(3)} = (\bar{u}_1)^{(3)}$if in addition $(u_0)^{(3)} = (u_1)^{(3)}$ then $T_{20}(t) = (u_0)^{(3)}T_{21}(t)$ This is an important consequence of the relation between $(\nu_1)^{(3)}$ and $(\bar{\nu}_1)^{(3)}$

Proof : From global equations we obtain 404

$$\frac{d\nu^{(4)}}{dt} = (a_{24})^{(4)} - \left((a'_{24})^{(4)} - (a'_{25})^{(4)} + (a''_{24})^{(4)}(T_{25}, t)\right) - (a''_{25})^{(4)}(T_{25}, t)\nu^{(4)} - (a_{25})^{(4)}\nu^{(4)}$$

Definition of $\nu^{(4)}$:- $\boxed{\nu^{(4)} = \frac{G_{24}}{G_{25}}}$

It follows

$$-\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_2)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right) \leq \frac{d\nu^{(4)}}{dt} \leq -\left((a_{25})^{(4)}\left(\nu^{(4)}\right)^2 + (\sigma_4)^{(4)}\nu^{(4)} - (a_{24})^{(4)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(4)}, (\nu_0)^{(4)}$:-

For $0 < \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}} < (\nu_1)^{(4)} < (\bar{\nu}_1)^{(4)}$

$$\nu^{(4)}(t) \geq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}}{4 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_0)^{(4)}\right)t\right]}} \quad , \quad \boxed{(C)^{(4)} = \frac{(\nu_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\nu_2)^{(4)}}}$$

it follows $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\nu_1)^{(4)}$

In the same manner , we get 405

$$\nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{4 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(4)} = \frac{(\bar{\nu}_1)^{(4)} - (\nu_0)^{(4)}}{(\nu_0)^{(4)} - (\bar{\nu}_2)^{(4)}}}$$

From which we deduce $(\nu_0)^{(4)} \leq \nu^{(4)}(t) \leq (\bar{\nu}_1)^{(4)}$

If $0 < (\nu_1)^{(4)} < (\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0} < (\bar{\nu}_1)^{(4)}$ we find like in the previous case, 406

$$(\nu_1)^{(4)} \leq \frac{(\nu_1)^{(4)} + (C)^{(4)}(\nu_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}}{1 + (C)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\nu_1)^{(4)} - (\nu_2)^{(4)}\right)t\right]}} \leq \nu^{(4)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\bar{\nu}_1)^{(4)}$$

If $0 < (\nu_1)^{(4)} \leq (\bar{\nu}_1)^{(4)} \leq \boxed{(\nu_0)^{(4)} = \frac{G_{24}^0}{G_{25}^0}}$, we obtain 407

$$(\nu_1)^{(4)} \leq \nu^{(4)}(t) \leq \frac{(\bar{\nu}_1)^{(4)} + (\bar{C})^{(4)}(\bar{\nu}_2)^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}}{1 + (\bar{C})^{(4)} e^{\left[-(a_{25})^{(4)}\left((\bar{\nu}_1)^{(4)} - (\bar{\nu}_2)^{(4)}\right)t\right]}} \leq (\nu_0)^{(4)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(4)}(t)$:-

$$(m_2)^{(4)} \leq \nu^{(4)}(t) \leq (m_1)^{(4)}, \quad \boxed{\nu^{(4)}(t) = \frac{G_{24}(t)}{G_{25}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(4)}(t)$:-

$$(\mu_2)^{(4)} \leq u^{(4)}(t) \leq (\mu_1)^{(4)}, \quad \boxed{u^{(4)}(t) = \frac{T_{24}(t)}{T_{25}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{24})^{(4)} = (a''_{25})^{(4)}, then\ (\sigma_1)^{(4)} = (\sigma_2)^{(4)}$ and in this case $(\nu_1)^{(4)} = (\bar{\nu}_1)^{(4)}$ if in addition $(\nu_0)^{(4)} = (\nu_1)^{(4)}$ then $\nu^{(4)}(t) = (\nu_0)^{(4)}$ and as a consequence $G_{24}(t) = (\nu_0)^{(4)} G_{25}(t)$ **this also defines** $(\nu_0)^{(4)}$ **for the special case** .

Analogously if $(b''_{24})^{(4)} = (b''_{25})^{(4)}, then\ (\tau_1)^{(4)} = (\tau_2)^{(4)}$ and then $(u_1)^{(4)} = (\bar{u}_4)^{(4)}$ if in addition $(u_0)^{(4)} = (u_1)^{(4)}$ then $T_{24}(t) = (u_0)^{(4)} T_{25}(t)$ This is an important consequence of the relation between $(\nu_1)^{(4)}$ and $(\bar{\nu}_1)^{(4)}$, **and definition of** $(u_0)^{(4)}$.

408

Proof : From global equations we obtain

$$\frac{d\nu^{(5)}}{dt} = (a_{28})^{(5)} - \left((a'_{28})^{(5)} - (a'_{29})^{(5)} + (a''_{28})^{(5)}(T_{29}, t)\right) - (a''_{29})^{(5)}(T_{29}, t)\nu^{(5)} - (a_{29})^{(5)}\nu^{(5)}$$

Definition of $\nu^{(5)}$:- $\boxed{\nu^{(5)} = \frac{G_{28}}{G_{29}}}$

It follows

$$-\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_2)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right) \leq \frac{d\nu^{(5)}}{dt} \leq -\left((a_{29})^{(5)}\left(\nu^{(5)}\right)^2 + (\sigma_1)^{(5)}\nu^{(5)} - (a_{28})^{(5)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(5)}, (\nu_0)^{(5)}$:-

For $0 < \boxed{(\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}}} < (\nu_1)^{(5)} < (\bar{\nu}_1)^{(5)}$

$$\nu^{(5)}(t) \geq \frac{(\nu_1)^{(5)} + (C)^{(5)}(\nu_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}}{5 + (C)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)} - (\nu_0)^{(5)}\right)t\right]}} \quad , \quad \boxed{(C)^{(5)} = \frac{(\nu_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\nu_2)^{(5)}}}$$

it follows $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\nu_1)^{(5)}$

In the same manner , we get 409

$$\nu^{(5)}(t) \leq \frac{(\bar{\nu}_1)^{(5)} + (\bar{C})^{(5)}(\bar{\nu}_2)^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}}{5 + (\bar{C})^{(5)} e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)} - (\bar{\nu}_2)^{(5)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(5)} = \frac{(\bar{\nu}_1)^{(5)} - (\nu_0)^{(5)}}{(\nu_0)^{(5)} - (\bar{\nu}_2)^{(5)}}}$$

From which we deduce $(\nu_0)^{(5)} \leq \nu^{(5)}(t) \leq (\bar{\nu}_5)^{(5)}$

If $0 < (\nu_1)^{(5)} < (\nu_0)^{(5)} = \frac{G^0_{28}}{G^0_{29}} < (\bar{\nu}_1)^{(5)}$ we find like in the previous case, 410

$$(\nu_1)^{(5)} \le \frac{(\nu_1)^{(5)}+(C)^{(5)}(\nu_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}}{1+(C)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\nu_1)^{(5)}-(\nu_2)^{(5)}\right)t\right]}} \le \nu^{(5)}(t) \le$$

$$\frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\bar{\nu}_1)^{(5)}$$

If $0 < (\nu_1)^{(5)} \le (\bar{\nu}_1)^{(5)} \le \boxed{(\nu_0)^{(5)} = \frac{G_{28}^0}{G_{29}^0}}$, we obtain 411

$$(\nu_1)^{(5)} \le \nu^{(5)}(t) \le \frac{(\bar{\nu}_1)^{(5)}+(\bar{C})^{(5)}(\bar{\nu}_2)^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}}{1+(\bar{C})^{(5)}e^{\left[-(a_{29})^{(5)}\left((\bar{\nu}_1)^{(5)}-(\bar{\nu}_2)^{(5)}\right)t\right]}} \le (\nu_0)^{(5)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(5)}(t)$:-

$(m_2)^{(5)} \le \nu^{(5)}(t) \le (m_1)^{(5)}$, $\boxed{\nu^{(5)}(t) = \frac{G_{28}(t)}{G_{29}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(5)}(t)$:-

$(\mu_2)^{(5)} \le u^{(5)}(t) \le (\mu_1)^{(5)}$, $\boxed{u^{(5)}(t) = \frac{T_{28}(t)}{T_{29}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a_{28}'')^{(5)} = (a_{29}'')^{(5)}, then\ (\sigma_1)^{(5)} = (\sigma_2)^{(5)}$ and in this case $(\nu_1)^{(5)} = (\bar{\nu}_1)^{(5)}$ if in addition $(\nu_0)^{(5)} = (\nu_5)^{(5)}$ then $\nu^{(5)}(t) = (\nu_0)^{(5)}$ and as a consequence $G_{28}(t) = (\nu_0)^{(5)}G_{29}(t)$ **this also defines** $(\nu_0)^{(5)}$ **for the special case** .

Analogously if $(b_{28}'')^{(5)} = (b_{29}'')^{(5)}, then\ (\tau_1)^{(5)} = (\tau_2)^{(5)}$ and then $(u_1)^{(5)} = (\bar{u}_1)^{(5)}$if in addition $(u_0)^{(5)} = (u_1)^{(5)}$ then $T_{28}(t) = (u_0)^{(5)}T_{29}(t)$ This is an important consequence of the relation between $(\nu_1)^{(5)}$ and $(\bar{\nu}_1)^{(5)}$, **and definition of** $(u_0)^{(5)}$.

Proof : From global equations we obtain 412

$$\frac{d\nu^{(6)}}{dt} = (a_{32})^{(6)} - \left((a_{32}')^{(6)} - (a_{33}')^{(6)} + (a_{32}'')^{(6)}(T_{33}, t)\right) - (a_{33}'')^{(6)}(T_{33}, t)\nu^{(6)} - (a_{33})^{(6)}\nu^{(6)}$$

Definition of $\nu^{(6)}$:- $\boxed{\nu^{(6)} = \frac{G_{32}}{G_{33}}}$

It follows

$$-\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_2)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right) \le \frac{d\nu^{(6)}}{dt} \le -\left((a_{33})^{(6)}\left(\nu^{(6)}\right)^2 + (\sigma_1)^{(6)}\nu^{(6)} - (a_{32})^{(6)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(6)}, (\nu_0)^{(6)}$:-

For $0 < \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}} < (\nu_1)^{(6)} < (\bar{\nu}_1)^{(6)}$

$$\nu^{(6)}(t) \geq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_0)^{(6)}\right)t\right]}} \quad , \quad \boxed{(C)^{(6)} = \frac{(\nu_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\nu_2)^{(6)}}}$$

it follows $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\nu_1)^{(6)}$

In the same manner , we get 413

$$\nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(6)} = \frac{(\bar{\nu}_1)^{(6)}-(\nu_0)^{(6)}}{(\nu_0)^{(6)}-(\bar{\nu}_2)^{(6)}}}$$

From which we deduce $(\nu_0)^{(6)} \leq \nu^{(6)}(t) \leq (\bar{\nu}_1)^{(6)}$

If $0 < (\nu_1)^{(6)} < (\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0} < (\bar{\nu}_1)^{(6)}$ we find like in the previous case, 414

$$(\nu_1)^{(6)} \leq \frac{(\nu_1)^{(6)}+(C)^{(6)}(\nu_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}}{1+(C)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\nu_1)^{(6)}-(\nu_2)^{(6)}\right)t\right]}} \leq \nu^{(6)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\bar{\nu}_1)^{(6)}$$

If $0 < (\nu_1)^{(6)} \leq (\bar{\nu}_1)^{(6)} \leq \boxed{(\nu_0)^{(6)} = \frac{G_{32}^0}{G_{33}^0}}$, we obtain 415

$$(\nu_1)^{(6)} \leq \nu^{(6)}(t) \leq \frac{(\bar{\nu}_1)^{(6)}+(\bar{C})^{(6)}(\bar{\nu}_2)^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}}{1+(\bar{C})^{(6)}e^{\left[-(a_{33})^{(6)}\left((\bar{\nu}_1)^{(6)}-(\bar{\nu}_2)^{(6)}\right)t\right]}} \leq (\nu_0)^{(6)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(6)}(t)$:-

$$(m_2)^{(6)} \leq \nu^{(6)}(t) \leq (m_1)^{(6)}, \quad \boxed{\nu^{(6)}(t) = \frac{G_{32}(t)}{G_{33}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(6)}(t)$:-

$$(\mu_2)^{(6)} \leq u^{(6)}(t) \leq (\mu_1)^{(6)}, \quad \boxed{u^{(6)}(t) = \frac{T_{32}(t)}{T_{33}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{32})^{(6)} = (a''_{33})^{(6)}, then\ (\sigma_1)^{(6)} = (\sigma_2)^{(6)}$ and in this case $(\nu_1)^{(6)} = (\bar{\nu}_1)^{(6)}$ if in addition $(\nu_0)^{(6)} = (\nu_1)^{(6)}$ then $\nu^{(6)}(t) = (\nu_0)^{(6)}$ and as a consequence $G_{32}(t) = (\nu_0)^{(6)}G_{33}(t)$ **this also defines** $(\nu_0)^{(6)}$ **for the special case** .

Analogously if $(b''_{32})^{(6)} = (b''_{33})^{(6)}, then\ (\tau_1)^{(6)} = (\tau_2)^{(6)}$ and then

$(u_1)^{(6)} = (\bar{u}_1)^{(6)}$if in addition $(u_0)^{(6)} = (u_1)^{(6)}$ then $T_{32}(t) = (u_0)^{(6)}T_{33}(t)$ This is an important consequence of the relation between $(v_1)^{(6)}$ and $(\bar{v}_1)^{(6)}$, **and definition of** $(u_0)^{(6)}$.

416

Proof : From global equations we obtain

$$\frac{dv^{(7)}}{dt} = (a_{36})^{(7)} - \left((a'_{36})^{(7)} - (a'_{37})^{(7)} + (a''_{36})^{(7)}(T_{37}, t)\right) - (a''_{37})^{(7)}(T_{37}, t)v^{(7)} - (a_{37})^{(7)}v^{(7)}$$

Definition of $v^{(7)}$:- $\boxed{v^{(7)} = \frac{G_{36}}{G_{37}}}$

It follows

$$-\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_2)^{(7)}v^{(7)} - (a_{36})^{(7)}\right) \leq \frac{dv^{(7)}}{dt} \leq -\left((a_{37})^{(7)}\left(v^{(7)}\right)^2 + (\sigma_1)^{(7)}v^{(7)} - (a_{36})^{(7)}\right)$$

From which one obtains

Definition of $(\bar{v}_1)^{(7)}, (v_0)^{(7)}$:-

For $0 < \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}} < (v_1)^{(7)} < (\bar{v}_1)^{(7)}$

$$v^{(7)}(t) \geq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_0)^{(7)}\right)t\right]}} \quad , \quad \boxed{(C)^{(7)} = \frac{(v_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (v_2)^{(7)}}}$$

it follows $(v_0)^{(7)} \leq v^{(7)}(t) \leq (v_1)^{(7)}$

In the same manner , we get 417

$$v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(7)} = \frac{(\bar{v}_1)^{(7)} - (v_0)^{(7)}}{(v_0)^{(7)} - (\bar{v}_2)^{(7)}}}$$

From which we deduce $(v_0)^{(7)} \leq v^{(7)}(t) \leq (\bar{v}_1)^{(7)}$

If $0 < (v_1)^{(7)} < (v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}} < (\bar{v}_1)^{(7)}$ we find like in the previous case, 418

$$(v_1)^{(7)} \leq \frac{(v_1)^{(7)} + (C)^{(7)}(v_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}}{1 + (C)^{(7)}e^{\left[-(a_{37})^{(7)}\left((v_1)^{(7)} - (v_2)^{(7)}\right)t\right]}} \leq v^{(7)}(t) \leq$$

$$\frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (\bar{v}_1)^{(7)}$$

If $0 < (v_1)^{(7)} \leq (\bar{v}_1)^{(7)} \leq \boxed{(v_0)^{(7)} = \frac{G^0_{36}}{G^0_{37}}}$, we obtain 419

$$(v_1)^{(7)} \leq v^{(7)}(t) \leq \frac{(\bar{v}_1)^{(7)} + (\bar{C})^{(7)}(\bar{v}_2)^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}}{1 + (\bar{C})^{(7)}e^{\left[-(a_{37})^{(7)}\left((\bar{v}_1)^{(7)} - (\bar{v}_2)^{(7)}\right)t\right]}} \leq (v_0)^{(7)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(7)}(t)$:-

$(m_2)^{(7)} \leq \nu^{(7)}(t) \leq (m_1)^{(7)}$, $\boxed{\nu^{(7)}(t) = \frac{G_{36}(t)}{G_{37}(t)}}$

In a completely analogous way, we obtain

Definition of $u^{(7)}(t)$:- 420

$(\mu_2)^{(7)} \leq u^{(7)}(t) \leq (\mu_1)^{(7)}$, $\boxed{u^{(7)}(t) = \frac{T_{36}(t)}{T_{37}(t)}}$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{36})^{(7)} = (a''_{37})^{(7)}, then\ (\sigma_1)^{(7)} = (\sigma_2)^{(7)}$ and in this case $(\nu_1)^{(7)} = (\bar{\nu}_1)^{(7)}$ if in addition $(\nu_0)^{(7)} = (\nu_1)^{(7)}$ then $\nu^{(7)}(t) = (\nu_0)^{(7)}$ and as a consequence $G_{36}(t) = (\nu_0)^{(7)} G_{37}(t)$ **this also defines** $(\nu_0)^{(7)}$ **for the special case** .

Analogously if $(b''_{36})^{(7)} = (b''_{37})^{(7)}, then\ (\tau_1)^{(7)} = (\tau_2)^{(7)}$ and then $(u_1)^{(7)} = (\bar{u}_1)^{(7)}$if in addition $(u_0)^{(7)} = (u_1)^{(7)}$ then $T_{36}(t) = (u_0)^{(7)} T_{37}(t)$ This is an important consequence of the relation between $(\nu_1)^{(7)}$ and $(\bar{\nu}_1)^{(7)}$, **and definition of** $(u_0)^{(7)}$.

Proof : From global equations we obtain 421

$$\frac{d\nu^{(8)}}{dt} = (a_{40})^{(8)} - \left((a'_{40})^{(8)} - (a'_{41})^{(8)} + (a''_{40})^{(8)}(T_{41}, t)\right) - (a''_{41})^{(8)}(T_{41}, t)\nu^{(8)} - (a_{41})^{(8)}\nu^{(8)}$$

Definition of $\nu^{(8)}$:- $\boxed{\nu^{(8)} = \frac{G_{40}}{G_{41}}}$

It follows

$$-\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_2)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right) \leq \frac{d\nu^{(8)}}{dt} \leq -\left((a_{41})^{(8)}\left(\nu^{(8)}\right)^2 + (\sigma_1)^{(8)}\nu^{(8)} - (a_{40})^{(8)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(8)}, (\nu_0)^{(8)}$:-

For $0 < \boxed{(\nu_0)^{(8)} = \frac{G^0_{40}}{G^0_{41}}} < (\nu_1)^{(8)} < (\bar{\nu}_1)^{(8)}$

$$\nu^{(8)}(t) \geq \frac{(\nu_1)^{(8)} + (C)^{(8)}(\nu_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}}{1 + (C)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)} - (\nu_0)^{(8)}\right)t\right]}} \quad , \quad \boxed{(C)^{(8)} = \frac{(\nu_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\nu_2)^{(8)}}}$$

it follows $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\nu_1)^{(8)}$

In the same manner , we get 422

$$\nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)} + (\bar{C})^{(8)}(\bar{\nu}_2)^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}}{1 + (\bar{C})^{(8)} e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)} - (\bar{\nu}_2)^{(8)}\right)t\right]}} \quad , \quad \boxed{(\bar{C})^{(8)} = \frac{(\bar{\nu}_1)^{(8)} - (\nu_0)^{(8)}}{(\nu_0)^{(8)} - (\bar{\nu}_2)^{(8)}}}$$

From which we deduce $(\nu_0)^{(8)} \leq \nu^{(8)}(t) \leq (\bar{\nu}_8)^{(8)}$

If $0 < (\nu_1)^{(8)} < (\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0} < (\bar{\nu}_1)^{(8)}$ we find like in the previous case, 423

$$(\nu_1)^{(8)} \leq \frac{(\nu_1)^{(8)}+(C)^{(8)}(\nu_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}}{1+(C)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\nu_1)^{(8)}-(\nu_2)^{(8)}\right)t\right]}} \leq \nu^{(8)}(t) \leq$$

$$\frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\bar{\nu}_1)^{(8)}$$

If $0 < (\nu_1)^{(8)} \leq (\bar{\nu}_1)^{(8)} \leq \boxed{(\nu_0)^{(8)} = \frac{G_{40}^0}{G_{41}^0}}$, we obtain 424

$$(\nu_1)^{(8)} \leq \nu^{(8)}(t) \leq \frac{(\bar{\nu}_1)^{(8)}+(\bar{C})^{(8)}(\bar{\nu}_2)^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}}{1+(\bar{C})^{(8)}e^{\left[-(a_{41})^{(8)}\left((\bar{\nu}_1)^{(8)}-(\bar{\nu}_2)^{(8)}\right)t\right]}} \leq (\nu_0)^{(8)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $\nu^{(8)}(t)$:-

$$(m_2)^{(8)} \leq \nu^{(8)}(t) \leq (m_1)^{(8)}, \quad \boxed{\nu^{(8)}(t) = \frac{G_{40}(t)}{G_{41}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(8)}(t)$:-

$$(\mu_2)^{(8)} \leq u^{(8)}(t) \leq (\mu_1)^{(8)}, \quad \boxed{u^{(8)}(t) = \frac{T_{40}(t)}{T_{41}(t)}}$$

Now, using this result and replacing it in global equations we get easily the result stated in the theorem.

Particular case :

If $(a''_{40})^{(8)} = (a''_{41})^{(8)}, then\ (\sigma_1)^{(8)} = (\sigma_2)^{(8)}$ and in this case $(\nu_1)^{(8)} = (\bar{\nu}_1)^{(8)}$ if in addition $(\nu_0)^{(8)} = (\nu_1)^{(8)}$ then $\nu^{(8)}(t) = (\nu_0)^{(8)}$ and as a consequence $G_{40}(t) = (\nu_0)^{(8)}G_{41}(t)$ **this also defines** $(\nu_0)^{(8)}$ **for the special case** .

Analogously if $(b''_{40})^{(8)} = (b''_{41})^{(8)}, then\ (\tau_1)^{(8)} = (\tau_2)^{(8)}$ and then
$(u_1)^{(8)} = (\bar{u}_1)^{(8)}$if in addition $(u_0)^{(8)} = (u_1)^{(8)}$ then $T_{40}(t) = (u_0)^{(8)}T_{41}(t)$ This is an important consequence of the relation between $(\nu_1)^{(8)}$ and $(\bar{\nu}_1)^{(8)}$, **and definition of** $(u_0)^{(8)}$.

424 A

Proof : From 99,20,44,22,23,44 we obtain

$$\frac{d\nu^{(9)}}{dt} = (a_{44})^{(9)} - \left((a'_{44})^{(9)} - (a'_{45})^{(9)} + (a''_{44})^{(9)}(T_{45}, t)\right) - (a''_{45})^{(9)}(T_{45}, t)\nu^{(9)} - (a_{45})^{(9)}\nu^{(9)}$$

Definition of $\nu^{(9)}$:- $\quad \boxed{\nu^{(9)} = \frac{G_{44}}{G_{45}}}$

It follows

$$-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_2)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)\le\frac{dv^{(9)}}{dt}\le-\left((a_{45})^{(9)}\left(v^{(9)}\right)^2+(\sigma_1)^{(9)}v^{(9)}-(a_{44})^{(9)}\right)$$

From which one obtains

Definition of $(\bar{\nu}_1)^{(9)},(\nu_0)^{(9)}$:-

For $0<\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}<(\nu_1)^{(9)}<(\bar{\nu}_1)^{(9)}$

$$v^{(9)}(t)\ge\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_0)^{(9)}\right)t\right]}}\quad,\quad\boxed{(C)^{(9)}=\frac{(\nu_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\nu_2)^{(9)}}}$$

it follows $(\nu_0)^{(9)}\le v^{(9)}(t)\le(\nu_9)^{(9)}$

In the same manner , we get

$$v^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\quad,\quad\boxed{(\bar{C})^{(9)}=\frac{(\bar{\nu}_1)^{(9)}-(\nu_0)^{(9)}}{(\nu_0)^{(9)}-(\bar{\nu}_2)^{(9)}}}$$

From which we deduce $(\nu_0)^{(9)}\le v^{(9)}(t)\le(\bar{\nu}_1)^{(9)}$

If $0<(\nu_1)^{(9)}<(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}<(\bar{\nu}_1)^{(9)}$ we find like in the previous case,

$$(\nu_1)^{(9)}\le\frac{(\nu_1)^{(9)}+(C)^{(9)}(\nu_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}{1+(C)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\nu_1)^{(9)}-(\nu_2)^{(9)}\right)t\right]}}\le v^{(9)}(t)\le$$

$$\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\bar{\nu}_1)^{(9)}$$

If $0<(\nu_1)^{(9)}\le(\bar{\nu}_1)^{(9)}\le\boxed{(\nu_0)^{(9)}=\frac{G_{44}^0}{G_{45}^0}}$, we obtain

$$(\nu_1)^{(9)}\le v^{(9)}(t)\le\frac{(\bar{\nu}_1)^{(9)}+(\bar{C})^{(9)}(\bar{\nu}_2)^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}{1+(\bar{C})^{(9)}e^{\left[-(a_{45})^{(9)}\left((\bar{\nu}_1)^{(9)}-(\bar{\nu}_2)^{(9)}\right)t\right]}}\le(\nu_0)^{(9)}$$

And so with the notation of the first part of condition (c) , we have

Definition of $v^{(9)}(t)$:-

$$(m_2)^{(9)}\le v^{(9)}(t)\le(m_1)^{(9)},\quad\boxed{v^{(9)}(t)=\frac{G_{44}(t)}{G_{45}(t)}}$$

In a completely analogous way, we obtain

Definition of $u^{(9)}(t)$:-

$$(\mu_2)^{(9)}\le u^{(9)}(t)\le(\mu_1)^{(9)},\quad\boxed{u^{(9)}(t)=\frac{T_{44}(t)}{T_{45}(t)}}$$

Now, using this result and replacing it in 99, 20,44,22,23, and 44 we get easily the result stated in the theorem.

Particular case :

If $(a''_{44})^{(9)} = (a''_{45})^{(9)}, then\ (\sigma_1)^{(9)} = (\sigma_2)^{(9)}$ and in this case $(\nu_1)^{(9)} = (\bar{\nu}_1)^{(9)}$ if in addition $(\nu_0)^{(9)} = (\nu_1)^{(9)}$ then $\nu^{(9)}(t) = (\nu_0)^{(9)}$ and as a consequence $G_{44}(t) = (\nu_0)^{(9)} G_{45}(t)$ **this also defines** $(\nu_0)^{(9)}$ **for the special case** .

Analogously if $(b''_{44})^{(9)} = (b''_{45})^{(9)}, then\ (\tau_1)^{(9)} = (\tau_2)^{(9)}$ and then $(u_1)^{(9)} = (\bar{u}_1)^{(9)}$if in addition $(u_0)^{(9)} = (u_1)^{(9)}$ then $T_{44}(t) = (u_0)^{(9)} T_{45}(t)$ This is an important consequence of the relation between $(\nu_1)^{(9)}$ and $(\bar{\nu}_1)^{(9)}$, **and definition of** $(u_0)^{(9)}$.
We can prove the following 425

Theorem : If $(a''_i)^{(1)} and\ (b''_i)^{(1)}$ are independent on t , and the conditions with the notations

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} < 0$

$(a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a_{13})^{(1)}(p_{13})^{(1)} + (a'_{14})^{(1)}(p_{14})^{(1)} + (p_{13})^{(1)}(p_{14})^{(1)} > 0$

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} > 0$,

$(b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} - (b'_{13})^{(1)}(r_{14})^{(1)} - (b'_{14})^{(1)}(r_{14})^{(1)} + (r_{13})^{(1)}(r_{14})^{(1)} < 0$

$with\ (p_{13})^{(1)}, (r_{14})^{(1)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(2)} and\ (b''_i)^{(2)}$ are independent on t , and the conditions with the notations 426

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} < 0$ 427

$(a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a_{16})^{(2)}(p_{16})^{(2)} + (a'_{17})^{(2)}(p_{17})^{(2)} + (p_{16})^{(2)}(p_{17})^{(2)} > 0$ 428

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} > 0$, 429

$(b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} - (b'_{16})^{(2)}(r_{17})^{(2)} - (b'_{17})^{(2)}(r_{17})^{(2)} + (r_{16})^{(2)}(r_{17})^{(2)} < 0$ 430

$with\ (p_{16})^{(2)}, (r_{17})^{(2)}$ as defined by equation are satisfied , then the system

Theorem : If $(a''_i)^{(3)} and\ (b''_i)^{(3)}$ are independent on t , and the conditions with the notations 431

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} < 0$

$(a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a_{20})^{(3)}(p_{20})^{(3)} + (a'_{21})^{(3)}(p_{21})^{(3)} + (p_{20})^{(3)}(p_{21})^{(3)} > 0$

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} > 0$,

$(b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} - (b'_{20})^{(3)}(r_{21})^{(3)} - (b'_{21})^{(3)}(r_{21})^{(3)} + (r_{20})^{(3)}(r_{21})^{(3)} < 0$

$with\ (p_{20})^{(3)}, (r_{21})^{(3)}$ as defined by equation are satisfied , then the system

We can prove the following 432

Theorem : If $(a''_i)^{(4)} and\ (b''_i)^{(4)}$ are independent on t , and the conditions with the notations

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} < 0$

$(a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a_{24})^{(4)}(p_{24})^{(4)} + (a'_{25})^{(4)}(p_{25})^{(4)} + (p_{24})^{(4)}(p_{25})^{(4)} > 0$

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} > 0$,

$(b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} - (b'_{24})^{(4)}(r_{25})^{(4)} - (b'_{25})^{(4)}(r_{25})^{(4)} + (r_{24})^{(4)}(r_{25})^{(4)} < 0$

$with\ \ (p_{24})^{(4)}, (r_{25})^{(4)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(5)} and\ (b_i'')^{(5)}$ are independent on t , and the conditions with the notations 433

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} < 0$

$(a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a_{28})^{(5)}(p_{28})^{(5)} + (a'_{29})^{(5)}(p_{29})^{(5)} + (p_{28})^{(5)}(p_{29})^{(5)} > 0$

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} > 0$,

$(b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} - (b'_{28})^{(5)}(r_{29})^{(5)} - (b'_{29})^{(5)}(r_{29})^{(5)} + (r_{28})^{(5)}(r_{29})^{(5)} < 0$

$with\ \ (p_{28})^{(5)}, (r_{29})^{(5)}$ as defined by equation are satisfied , then the system

Theorem If $(a_i'')^{(6)} and\ (b_i'')^{(6)}$ are independent on t , and the conditions with the notations 434

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} < 0$

$(a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a_{32})^{(6)}(p_{32})^{(6)} + (a'_{33})^{(6)}(p_{33})^{(6)} + (p_{32})^{(6)}(p_{33})^{(6)} > 0$

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} > 0$,

$(b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} - (b'_{32})^{(6)}(r_{33})^{(6)} - (b'_{33})^{(6)}(r_{33})^{(6)} + (r_{32})^{(6)}(r_{33})^{(6)} < 0$

$with\ \ (p_{32})^{(6)}, (r_{33})^{(6)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(7)} and\ (b_i'')^{(7)}$ are independent on t , and the conditions with the notations 435

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} < 0$

$(a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a_{36})^{(7)}(p_{36})^{(7)} + (a'_{37})^{(7)}(p_{37})^{(7)} + (p_{36})^{(7)}(p_{37})^{(7)} > 0$

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} > 0$,

$(b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} - (b'_{36})^{(7)}(r_{37})^{(7)} - (b'_{37})^{(7)}(r_{37})^{(7)} + (r_{36})^{(7)}(r_{37})^{(7)} < 0$

$with\ \ (p_{36})^{(7)}, (r_{37})^{(7)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(8)} and\ (b_i'')^{(8)}$ are independent on t , and the conditions with the notations 436

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} < 0$

$(a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a_{40})^{(8)}(p_{40})^{(8)} + (a'_{41})^{(8)}(p_{41})^{(8)} + (p_{40})^{(8)}(p_{41})^{(8)} > 0$

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} > 0$,

$(b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} - (b'_{40})^{(8)}(r_{41})^{(8)} - (b'_{41})^{(8)}(r_{41})^{(8)} + (r_{40})^{(8)}(r_{41})^{(8)} < 0$

$with\ \ (p_{40})^{(8)}, (r_{41})^{(8)}$ as defined by equation are satisfied , then the system

Theorem : If $(a_i'')^{(9)} and\ (b_i'')^{(9)}$ are independent on t , and the conditions (with the notations 45,46,27,28) 436 A

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} < 0$

$(a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a_{44})^{(9)}(p_{44})^{(9)} + (a'_{45})^{(9)}(p_{45})^{(9)} + (p_{44})^{(9)}(p_{45})^{(9)} > 0$

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} > 0$,

$(b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} - (b'_{44})^{(9)}(r_{45})^{(9)} - (b'_{45})^{(9)}(r_{45})^{(9)} + (r_{44})^{(9)}(r_{45})^{(9)} < 0$

$with\ \ (p_{44})^{(9)}, (r_{45})^{(9)}$ as defined by equation 45 are satisfied , then the system

$$(a_{13})^{(1)}G_{14} - \left[(a'_{13})^{(1)} + (a''_{13})^{(1)}(T_{14})\right]G_{13} = 0 \quad 437$$

$$(a_{14})^{(1)}G_{13} - \left[(a'_{14})^{(1)} + (a''_{14})^{(1)}(T_{14})\right]G_{14} = 0 \quad 438$$

$$(a_{15})^{(1)}G_{14} - \left[(a'_{15})^{(1)} + (a''_{15})^{(1)}(T_{14})\right]G_{15} = 0 \quad 439$$

$$(b_{13})^{(1)}T_{14} - [(b'_{13})^{(1)} - (b''_{13})^{(1)}(G)\ ]T_{13} = 0 \quad 440$$

$$(b_{14})^{(1)}T_{13} - [(b'_{14})^{(1)} - (b''_{14})^{(1)}(G)\ ]T_{14} = 0 \quad 441$$

$$(b_{15})^{(1)}T_{14} - [(b'_{15})^{(1)} - (b''_{15})^{(1)}(G)\ ]T_{15} = 0 \quad 442$$

has a unique positive solution , which is an equilibrium solution for the system

$$(a_{16})^{(2)}G_{17} - \left[(a'_{16})^{(2)} + (a''_{16})^{(2)}(T_{17})\right]G_{16} = 0 \quad 443$$

$$(a_{17})^{(2)}G_{16} - \left[(a'_{17})^{(2)} + (a''_{17})^{(2)}(T_{17})\right]G_{17} = 0 \quad 444$$

$$(a_{18})^{(2)}G_{17} - \left[(a'_{18})^{(2)} + (a''_{18})^{(2)}(T_{17})\right]G_{18} = 0 \quad 445$$

$$(b_{16})^{(2)}T_{17} - [(b'_{16})^{(2)} - (b''_{16})^{(2)}(G_{19})\ ]T_{16} = 0 \quad 446$$

$$(b_{17})^{(2)}T_{16} - [(b'_{17})^{(2)} - (b''_{17})^{(2)}(G_{19})\ ]T_{17} = 0 \quad 447$$

$$(b_{18})^{(2)}T_{17} - [(b'_{18})^{(2)} - (b''_{18})^{(2)}(G_{19})\ ]T_{18} = 0 \quad 448$$

has a unique positive solution , which is an equilibrium solution

$$(a_{20})^{(3)}G_{21} - \left[(a'_{20})^{(3)} + (a''_{20})^{(3)}(T_{21})\right]G_{20} = 0 \quad 449$$

$$(a_{21})^{(3)}G_{20} - \left[(a'_{21})^{(3)} + (a''_{21})^{(3)}(T_{21})\right]G_{21} = 0 \quad 450$$

$$(a_{22})^{(3)}G_{21} - \left[(a'_{22})^{(3)} + (a''_{22})^{(3)}(T_{21})\right]G_{22} = 0 \quad 451$$

$$(b_{20})^{(3)}T_{21} - [(b'_{20})^{(3)} - (b''_{20})^{(3)}(G_{23})\ ]T_{20} = 0 \quad 452$$

$$(b_{21})^{(3)}T_{20} - [(b'_{21})^{(3)} - (b''_{21})^{(3)}(G_{23})\ ]T_{21} = 0 \quad 453$$

$$(b_{22})^{(3)}T_{21} - [(b'_{22})^{(3)} - (b''_{22})^{(3)}(G_{23})\ ]T_{22} = 0 \quad 454$$

has a unique positive solution , which is an equilibrium solution

$$(a_{24})^{(4)}G_{25} - \left[(a'_{24})^{(4)} + (a''_{24})^{(4)}(T_{25})\right]G_{24} = 0 \quad 455$$

$$(a_{25})^{(4)}G_{24} - \left[(a'_{25})^{(4)} + (a''_{25})^{(4)}(T_{25})\right]G_{25} = 0 \quad 456$$

$$(a_{26})^{(4)}G_{25} - \left[(a'_{26})^{(4)} + (a''_{26})^{(4)}(T_{25})\right]G_{26} = 0 \quad 457$$

$$(b_{24})^{(4)}T_{25} - [(b'_{24})^{(4)} - (b''_{24})^{(4)}\left((G_{27})\right)]T_{24} = 0 \quad 458$$

$$(b_{25})^{(4)}T_{24} - [(b'_{25})^{(4)} - (b''_{25})^{(4)}\left((G_{27})\right)]T_{25} = 0 \quad 459$$

$$(b_{26})^{(4)}T_{25} - [(b'_{26})^{(4)} - (b''_{26})^{(4)}\left((G_{27})\right)]T_{26} = 0 \quad 460$$

has a unique positive solution , which is an equilibrium solution

$$(a_{28})^{(5)}G_{29} - \left[(a'_{28})^{(5)} + (a''_{28})^{(5)}(T_{29})\right]G_{28} = 0 \quad 461$$

$$(a_{29})^{(5)}G_{28} - \left[(a'_{29})^{(5)} + (a''_{29})^{(5)}(T_{29})\right]G_{29} = 0 \quad 462$$

$$(a_{30})^{(5)}G_{29} - \left[(a'_{30})^{(5)} + (a''_{30})^{(5)}(T_{29})\right]G_{30} = 0 \quad 463$$

$$(b_{28})^{(5)}T_{29} - [(b'_{28})^{(5)} - (b''_{28})^{(5)}(G_{31})]T_{28} = 0 \quad 464$$

$$(b_{29})^{(5)}T_{28} - [(b'_{29})^{(5)} - (b''_{29})^{(5)}(G_{31})]T_{29} = 0 \quad 465$$

$$(b_{30})^{(5)}T_{29} - [(b'_{30})^{(5)} - (b''_{30})^{(5)}(G_{31})]T_{30} = 0 \quad 466$$

has a unique positive solution , which is an equilibrium solution

$$(a_{32})^{(6)}G_{33} - \left[(a'_{32})^{(6)} + (a''_{32})^{(6)}(T_{33})\right]G_{32} = 0 \quad 467$$

$$(a_{33})^{(6)}G_{32} - \left[(a'_{33})^{(6)} + (a''_{33})^{(6)}(T_{33})\right]G_{33} = 0 \quad 468$$

$$(a_{34})^{(6)}G_{33} - \left[(a'_{34})^{(6)} + (a''_{34})^{(6)}(T_{33})\right]G_{34} = 0 \quad 469$$

$$(b_{32})^{(6)}T_{33} - [(b'_{32})^{(6)} - (b''_{32})^{(6)}(G_{35})]T_{32} = 0 \quad 470$$

$$(b_{33})^{(6)}T_{32} - [(b'_{33})^{(6)} - (b''_{33})^{(6)}(G_{35})]T_{33} = 0 \quad 471$$

$$(b_{34})^{(6)}T_{33} - [(b'_{34})^{(6)} - (b''_{34})^{(6)}(G_{35})]T_{34} = 0 \quad 472$$

has a unique positive solution , which is an equilibrium solution

$$(a_{36})^{(7)}G_{37} - \left[(a'_{36})^{(7)} + (a''_{36})^{(7)}(T_{37})\right]G_{36} = 0 \quad 473$$

$$(a_{37})^{(7)}G_{36} - \left[(a'_{37})^{(7)} + (a''_{37})^{(7)}(T_{37})\right]G_{37} = 0 \quad 474$$

$$(a_{38})^{(7)}G_{37} - \left[(a'_{38})^{(7)} + (a''_{38})^{(7)}(T_{37})\right]G_{38} = 0 \quad 475$$

$$(b_{36})^{(7)}T_{37} - [(b'_{36})^{(7)} - (b''_{36})^{(7)}(G_{39})]T_{36} = 0 \quad 476$$

$$(b_{37})^{(7)}T_{36} - [(b'_{37})^{(7)} - (b''_{37})^{(7)}(G_{39})\,]T_{37} = 0 \quad 477$$

$$(b_{38})^{(7)}T_{37} - [(b'_{38})^{(7)} - (b''_{38})^{(7)}(G_{39})\,]T_{38} = 0 \quad 478$$

$$(a_{40})^{(8)}G_{41} - \left[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41})\right]G_{40} = 0 \quad 479$$

$$(a_{41})^{(8)}G_{40} - \left[(a'_{41})^{(8)} + (a''_{41})^{(8)}(T_{41})\right]G_{41} = 0 \quad 480$$

$$(a_{42})^{(8)}G_{41} - \left[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41})\right]G_{42} = 0 \quad 481$$

$$(b_{40})^{(8)}T_{41} - [(b'_{40})^{(8)} - (b''_{40})^{(8)}(G_{43})\,]T_{40} = 0 \quad 482$$

$$(b_{41})^{(8)}T_{40} - [(b'_{41})^{(8)} - (b''_{41})^{(8)}(G_{43})\,]T_{41} = 0 \quad 483$$

$$(b_{42})^{(8)}T_{41} - [(b'_{42})^{(8)} - (b''_{42})^{(8)}(G_{43})\,]T_{42} = 0 \quad 484$$

$$(a_{44})^{(9)}G_{45} - \left[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45})\right]G_{44} = 0 \quad 484\text{A}$$

$$(a_{45})^{(9)}G_{44} - \left[(a'_{45})^{(9)} + (a''_{45})^{(9)}(T_{45})\right]G_{45} = 0$$

$$(a_{46})^{(9)}G_{45} - \left[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45})\right]G_{46} = 0$$

$$(b_{44})^{(9)}T_{45} - [(b'_{44})^{(9)} - (b''_{44})^{(9)}(G_{47})\,]T_{44} = 0$$

$$(b_{45})^{(9)}T_{44} - [(b'_{45})^{(9)} - (b''_{45})^{(9)}(G_{47})\,]T_{45} = 0$$

$$(b_{46})^{(9)}T_{45} - [(b'_{46})^{(9)} - (b''_{46})^{(9)}(G_{47})\,]T_{46} = 0$$

Proof: 485

(a) Indeed the first two equations have a nontrivial solution G_{13}, G_{14} if

$$F(T) = (a'_{13})^{(1)}(a'_{14})^{(1)} - (a_{13})^{(1)}(a_{14})^{(1)} + (a'_{13})^{(1)}(a''_{14})^{(1)}(T_{14}) + (a'_{14})^{(1)}(a''_{13})^{(1)}(T_{14}) + (a''_{13})^{(1)}(T_{14})(a''_{14})^{(1)}(T_{14}) = 0$$

Proof: 486

(p) Indeed the first two equations have a nontrivial solution G_{16}, G_{17} if

$$\mathrm{F}(T_{19}) = (a'_{16})^{(2)}(a'_{17})^{(2)} - (a_{16})^{(2)}(a_{17})^{(2)} + (a'_{16})^{(2)}(a''_{17})^{(2)}(T_{17}) + (a'_{17})^{(2)}(a''_{16})^{(2)}(T_{17}) + (a''_{16})^{(2)}(T_{17})(a''_{17})^{(2)}(T_{17}) = 0$$

Proof: 487

(a) Indeed the first two equations have a nontrivial solution G_{20}, G_{21} if

$F(T_{23}) = (a'_{20})^{(3)}(a'_{21})^{(3)} - (a_{20})^{(3)}(a_{21})^{(3)} + (a'_{20})^{(3)}(a''_{21})^{(3)}(T_{21}) + (a'_{21})^{(3)}(a''_{20})^{(3)}(T_{21}) + (a''_{20})^{(3)}(T_{21})(a''_{21})^{(3)}(T_{21}) = 0$

Proof: 488

(a) Indeed the first two equations have a nontrivial solution G_{24}, G_{25} if

$F(T_{27}) = (a'_{24})^{(4)}(a'_{25})^{(4)} - (a_{24})^{(4)}(a_{25})^{(4)} + (a'_{24})^{(4)}(a''_{25})^{(4)}(T_{25}) + (a'_{25})^{(4)}(a''_{24})^{(4)}(T_{25}) + (a''_{24})^{(4)}(T_{25})(a''_{25})^{(4)}(T_{25}) = 0$

Proof: 489

(a) Indeed the first two equations have a nontrivial solution G_{28}, G_{29} if

$F(T_{31}) = (a'_{28})^{(5)}(a'_{29})^{(5)} - (a_{28})^{(5)}(a_{29})^{(5)} + (a'_{28})^{(5)}(a''_{29})^{(5)}(T_{29}) + (a'_{29})^{(5)}(a''_{28})^{(5)}(T_{29}) + (a''_{28})^{(5)}(T_{29})(a''_{29})^{(5)}(T_{29}) = 0$

Proof: 490

(a) Indeed the first two equations have a nontrivial solution G_{32}, G_{33} if

$F(T_{35}) = (a'_{32})^{(6)}(a'_{33})^{(6)} - (a_{32})^{(6)}(a_{33})^{(6)} + (a'_{32})^{(6)}(a''_{33})^{(6)}(T_{33}) + (a'_{33})^{(6)}(a''_{32})^{(6)}(T_{33}) + (a''_{32})^{(6)}(T_{33})(a''_{33})^{(6)}(T_{33}) = 0$

Proof: 491

(a) Indeed the first two equations have a nontrivial solution G_{36}, G_{37} if

$F(T_{39}) = (a'_{36})^{(7)}(a'_{37})^{(7)} - (a_{36})^{(7)}(a_{37})^{(7)} + (a'_{36})^{(7)}(a''_{37})^{(7)}(T_{37}) + (a'_{37})^{(7)}(a''_{36})^{(7)}(T_{37}) + (a''_{36})^{(7)}(T_{37})(a''_{37})^{(7)}(T_{37}) = 0$

Proof: 492

(a) Indeed the first two equations have a nontrivial solution G_{40}, G_{41} if

$F(T_{43}) = (a'_{40})^{(8)}(a'_{41})^{(8)} - (a_{40})^{(8)}(a_{41})^{(8)} + (a'_{40})^{(8)}(a''_{41})^{(8)}(T_{41}) + (a'_{41})^{(8)}(a''_{40})^{(8)}(T_{41}) + (a''_{40})^{(8)}(T_{41})(a''_{41})^{(8)}(T_{41}) = 0$

Proof: 492 A

(a) Indeed the first two equations have a nontrivial solution G_{44}, G_{45} if

$F(T_{47}) = (a'_{44})^{(9)}(a'_{45})^{(9)} - (a_{44})^{(9)}(a_{45})^{(9)} + (a'_{44})^{(9)}(a''_{45})^{(9)}(T_{45}) + (a'_{45})^{(9)}(a''_{44})^{(9)}(T_{45}) + (a''_{44})^{(9)}(T_{45})(a''_{45})^{(9)}(T_{45}) = 0$

Definition and uniqueness of T_{14}^* :- 493

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(1)}(T_{14})$ being increasing, it follows that there exists a unique T_{14}^* for which $f(T_{14}^*) = 0$. With this value , we obtain from the three first equations

$$G_{13} = \frac{(a_{13})^{(1)}G_{14}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T^*_{14})]} \quad , \quad G_{15} = \frac{(a_{15})^{(1)}G_{14}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T^*_{14})]}$$

Definition and uniqueness of T^*_{17} :- 494

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(2)}(T_{17})$ being increasing, it follows that there exists a unique T^*_{17} for which $f(T^*_{17}) = 0$. With this value , we obtain from the three first equations

$$G_{16} = \frac{(a_{16})^{(2)}G_{17}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T^*_{17})]} \quad , \quad G_{18} = \frac{(a_{18})^{(2)}G_{17}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T^*_{17})]} \qquad 495$$

Definition and uniqueness of T^*_{21} :- 496

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(1)}(T_{21})$ being increasing, it follows that there exists a unique T^*_{21} for which $f(T^*_{21}) = 0$. With this value , we obtain from the three first equations

$$G_{20} = \frac{(a_{20})^{(3)}G_{21}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T^*_{21})]} \quad , \quad G_{22} = \frac{(a_{22})^{(3)}G_{21}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T^*_{21})]}$$

Definition and uniqueness of T^*_{25} :- 497

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(4)}(T_{25})$ being increasing, it follows that there exists a unique T^*_{25} for which $f(T^*_{25}) = 0$. With this value , we obtain from the three first equations

$$G_{24} = \frac{(a_{24})^{(4)}G_{25}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T^*_{25})]} \quad , \quad G_{26} = \frac{(a_{26})^{(4)}G_{25}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T^*_{25})]}$$

Definition and uniqueness of T^*_{29} :- 498

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(5)}(T_{29})$ being increasing, it follows that there exists a unique T^*_{29} for which $f(T^*_{29}) = 0$. With this value , we obtain from the three first equations

$$G_{28} = \frac{(a_{28})^{(5)}G_{29}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T^*_{29})]} \quad , \quad G_{30} = \frac{(a_{30})^{(5)}G_{29}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T^*_{29})]}$$

Definition and uniqueness of T^*_{33} :- 499

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(6)}(T_{33})$ being increasing, it follows that there exists a unique T^*_{33} for which $f(T^*_{33}) = 0$. With this value , we obtain from the three first equations

$$G_{32} = \frac{(a_{32})^{(6)}G_{33}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T^*_{33})]} \quad , \quad G_{34} = \frac{(a_{34})^{(6)}G_{33}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T^*_{33})]}$$

Definition and uniqueness of T^*_{37} :- 500

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a''_i)^{(7)}(T_{37})$ being increasing, it follows that there exists a unique T^*_{37} for which $f(T^*_{37}) = 0$. With this value , we obtain from the three first equations

$$G_{36} = \frac{(a_{36})^{(7)}G_{37}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T^*_{37})]} \quad , \quad G_{38} = \frac{(a_{38})^{(7)}G_{37}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T^*_{37})]}$$

Definition and uniqueness of T_{41}^* :- 501

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(8)}(T_{41})$ being increasing, it follows that there exists a unique T_{41}^* for which $f(T_{41}^*) = 0$. With this value , we obtain from the three first equations

$$G_{40} = \frac{(a_{40})^{(8)} G_{41}}{[(a'_{40})^{(8)} + (a''_{40})^{(8)}(T_{41}^*)]} \quad , \quad G_{42} = \frac{(a_{42})^{(8)} G_{41}}{[(a'_{42})^{(8)} + (a''_{42})^{(8)}(T_{41}^*)]}$$

Definition and uniqueness of T_{45}^* :- 501 A

After hypothesis $f(0) < 0, f(\infty) > 0$ and the functions $(a_i'')^{(9)}(T_{45})$ being increasing, it follows that there exists a unique T_{45}^* for which $f(T_{45}^*) = 0$. With this value , we obtain from the three first equations

$$G_{44} = \frac{(a_{44})^{(9)} G_{45}}{[(a'_{44})^{(9)} + (a''_{44})^{(9)}(T_{45}^*)]} \quad , \quad G_{46} = \frac{(a_{46})^{(9)} G_{45}}{[(a'_{46})^{(9)} + (a''_{46})^{(9)}(T_{45}^*)]}$$

By the same argument, the equations admit solutions G_{13}, G_{14} if 502

$$\varphi(G) = (b'_{13})^{(1)}(b'_{14})^{(1)} - (b_{13})^{(1)}(b_{14})^{(1)} -$$

$$\left[(b'_{13})^{(1)}(b''_{14})^{(1)}(G) + (b'_{14})^{(1)}(b''_{13})^{(1)}(G)\right] + (b''_{13})^{(1)}(G)(b''_{14})^{(1)}(G) = 0$$

Where in $G(G_{13}, G_{14}, G_{15}), G_{13}, G_{15}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{14} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi(G^*) = 0$

By the same argument, the equations admit solutions G_{16}, G_{17} if 503

$$\varphi(G_{19}) = (b'_{16})^{(2)}(b'_{17})^{(2)} - (b_{16})^{(2)}(b_{17})^{(2)} -$$

$$\left[(b'_{16})^{(2)}(b''_{17})^{(2)}(G_{19}) + (b'_{17})^{(2)}(b''_{16})^{(2)}(G_{19})\right] + (b''_{16})^{(2)}(G_{19})(b''_{17})^{(2)}(G_{19}) = 0$$

Where in $(G_{19})(G_{16}, G_{17}, G_{18}), G_{16}, G_{18}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{17} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{14}^* such that $\varphi((G_{19})^*) = 0$ 504

By the same argument, the equations admit solutions G_{20}, G_{21} if 505

$$\varphi(G_{23}) = (b'_{20})^{(3)}(b'_{21})^{(3)} - (b_{20})^{(3)}(b_{21})^{(3)} -$$

$$\left[(b'_{20})^{(3)}(b''_{21})^{(3)}(G_{23}) + (b'_{21})^{(3)}(b''_{20})^{(3)}(G_{23})\right] + (b''_{20})^{(3)}(G_{23})(b''_{21})^{(3)}(G_{23}) = 0$$

Where in $G_{23}(G_{20}, G_{21}, G_{22}), G_{20}, G_{22}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{21} taking into account the hypothesis $\varphi(0) > 0$, $\varphi(\infty) < 0$ it follows that there exists a unique G_{21}^* such that $\varphi((G_{23})^*) = 0$

By the same argument, the equations admit solutions G_{24}, G_{25} if 506

$$\varphi(G_{27}) = (b'_{24})^{(4)}(b'_{25})^{(4)} - (b_{24})^{(4)}(b_{25})^{(4)} -$$

$$\left[(b'_{24})^{(4)}(b''_{25})^{(4)}(G_{27}) + (b'_{25})^{(4)}(b''_{24})^{(4)}(G_{27})\right] + (b''_{24})^{(4)}(G_{27})(b''_{25})^{(4)}(G_{27}) = 0$$

Where in $(G_{27})(G_{24}, G_{25}, G_{26}), G_{24}, G_{26}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{25} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{25} such that $\varphi((G_{27})^*) = 0$

By the same argument, the equations admit solutions G_{28}, G_{29} if 507
$\varphi(G_{31}) = (b'_{28})^{(5)}(b'_{29})^{(5)} - (b_{28})^{(5)}(b_{29})^{(5)} -$

$$\left[(b'_{28})^{(5)}(b''_{29})^{(5)}(G_{31}) + (b'_{29})^{(5)}(b''_{28})^{(5)}(G_{31})\right] + (b''_{28})^{(5)}(G_{31})(b''_{29})^{(5)}(G_{31}) = 0$$

Where in $(G_{31})(G_{28}, G_{29}, G_{30}), G_{28}, G_{30}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{29} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{29} such that $\varphi((G_{31})^*) = 0$

By the same argument, the equations admit solutions G_{32}, G_{33} if 508
$\varphi(G_{35}) = (b'_{32})^{(6)}(b'_{33})^{(6)} - (b_{32})^{(6)}(b_{33})^{(6)} -$

$$\left[(b'_{32})^{(6)}(b''_{33})^{(6)}(G_{35}) + (b'_{33})^{(6)}(b''_{32})^{(6)}(G_{35})\right] + (b''_{32})^{(6)}(G_{35})(b''_{33})^{(6)}(G_{35}) = 0$$

Where in $(G_{35})(G_{32}, G_{33}, G_{34}), G_{32}, G_{34}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{33} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{33} such that $\varphi({G_{35}}^*) = 0$

By the same argument, the equations admit solutions G_{36}, G_{37} if 509
$\varphi(G_{39}) = (b'_{36})^{(7)}(b'_{37})^{(7)} - (b_{36})^{(7)}(b_{37})^{(7)} -$
$\left[(b'_{36})^{(7)}(b''_{37})^{(7)}(G_{39}) + (b'_{37})^{(7)}(b''_{36})^{(7)}(G_{39})\right] + (b''_{36})^{(7)}(G_{39})(b''_{37})^{(7)}(G_{39}) = 0$

Where in $(G_{39})(G_{36}, G_{37}, G_{38}), G_{36}, G_{38}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{37} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{37} such that $\varphi({G_{39}}^*) = 0$

By the same argument, the equations admit solutions G_{40}, G_{41} if 510
$\varphi(G_{43}) = (b'_{40})^{(8)}(b'_{41})^{(8)} - (b_{40})^{(8)}(b_{41})^{(8)} -$
$\left[(b'_{40})^{(8)}(b''_{41})^{(8)}(G_{43}) + (b'_{41})^{(8)}(b''_{40})^{(8)}(G_{43})\right] + (b''_{40})^{(8)}(G_{43})(b''_{41})^{(8)}(G_{43}) = 0$

Where in $(G_{43})(G_{40}, G_{41}, G_{42}), G_{40}, G_{42}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{41} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{41} such that $\varphi({G_{43}}^*) = 0$

By the same argument, the equations 92,93 admit solutions G_{44}, G_{45} if
$\varphi(G_{47}) = (b'_{44})^{(9)}(b'_{45})^{(9)} - (b_{44})^{(9)}(b_{45})^{(9)} -$

$$\left[(b'_{44})^{(9)}(b''_{45})^{(9)}(G_{47}) + (b'_{45})^{(9)}(b''_{44})^{(9)}(G_{47})\right] + (b''_{44})^{(9)}(G_{47})(b''_{45})^{(9)}(G_{47}) = 0$$

Where in $(G_{47})(G_{44}, G_{45}, G_{46}), G_{44}, G_{46}$ must be replaced by their values from 96. It is easy to see that φ is a decreasing function in G_{45} taking into account the hypothesis $\varphi(0) > 0\,, \varphi(\infty) < 0$ it follows that there exists a unique G^*_{45} such that $\varphi((G_{47})^*) = 0$

Finally we obtain the unique solution 511

G^*_{14} given by $\varphi(G^*) = 0$, T^*_{14} given by $f(T^*_{14}) = 0$ and

$G_{13}^{*} = \frac{(a_{13})^{(1)}G_{14}^{*}}{[(a'_{13})^{(1)}+(a''_{13})^{(1)}(T_{14}^{*})]}$, $G_{15}^{*} = \frac{(a_{15})^{(1)}G_{14}^{*}}{[(a'_{15})^{(1)}+(a''_{15})^{(1)}(T_{14}^{*})]}$

$T_{13}^{*} = \frac{(b_{13})^{(1)}T_{14}^{*}}{[(b'_{13})^{(1)}-(b''_{13})^{(1)}(G^{*})]}$, $T_{15}^{*} = \frac{(b_{15})^{(1)}T_{14}^{*}}{[(b'_{15})^{(1)}-(b''_{15})^{(1)}(G^{*})]}$

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution

G_{17}^{*} given by $\varphi((G_{19})^{*}) = 0$, T_{17}^{*} given by $f(T_{17}^{*}) = 0$ and 512

$G_{16}^{*} = \frac{(a_{16})^{(2)}G_{17}^{*}}{[(a'_{16})^{(2)}+(a''_{16})^{(2)}(T_{17}^{*})]}$, $G_{18}^{*} = \frac{(a_{18})^{(2)}G_{17}^{*}}{[(a'_{18})^{(2)}+(a''_{18})^{(2)}(T_{17}^{*})]}$ 513

$T_{16}^{*} = \frac{(b_{16})^{(2)}T_{17}^{*}}{[(b'_{16})^{(2)}-(b''_{16})^{(2)}((G_{19})^{*})]}$, $T_{18}^{*} = \frac{(b_{18})^{(2)}T_{17}^{*}}{[(b'_{18})^{(2)}-(b''_{18})^{(2)}((G_{19})^{*})]}$ 514

Obviously, these values represent an equilibrium solution

Finally we obtain the unique solution 515

G_{21}^{*} given by $\varphi((G_{23})^{*}) = 0$, T_{21}^{*} given by $f(T_{21}^{*}) = 0$ and

$G_{20}^{*} = \frac{(a_{20})^{(3)}G_{21}^{*}}{[(a'_{20})^{(3)}+(a''_{20})^{(3)}(T_{21}^{*})]}$, $G_{22}^{*} = \frac{(a_{22})^{(3)}G_{21}^{*}}{[(a'_{22})^{(3)}+(a''_{22})^{(3)}(T_{21}^{*})]}$

$T_{20}^{*} = \frac{(b_{20})^{(3)}T_{21}^{*}}{[(b'_{20})^{(3)}-(b''_{20})^{(3)}(G_{23}{}^{*})]}$, $T_{22}^{*} = \frac{(b_{22})^{(3)}T_{21}^{*}}{[(b'_{22})^{(3)}-(b''_{22})^{(3)}(G_{23}{}^{*})]}$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 516

G_{25}^{*} given by $\varphi(G_{27}) = 0$, T_{25}^{*} given by $f(T_{25}^{*}) = 0$ and

$G_{24}^{*} = \frac{(a_{24})^{(4)}G_{25}^{*}}{[(a'_{24})^{(4)}+(a''_{24})^{(4)}(T_{25}^{*})]}$, $G_{26}^{*} = \frac{(a_{26})^{(4)}G_{25}^{*}}{[(a'_{26})^{(4)}+(a''_{26})^{(4)}(T_{25}^{*})]}$

$T_{24}^{*} = \frac{(b_{24})^{(4)}T_{25}^{*}}{[(b'_{24})^{(4)}-(b''_{24})^{(4)}((G_{27})^{*})]}$, $T_{26}^{*} = \frac{(b_{26})^{(4)}T_{25}^{*}}{[(b'_{26})^{(4)}-(b''_{26})^{(4)}((G_{27})^{*})]}$ 517

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 518

G_{29}^{*} given by $\varphi((G_{31})^{*}) = 0$, T_{29}^{*} given by $f(T_{29}^{*}) = 0$ and

$G_{28}^{*} = \frac{(a_{28})^{(5)}G_{29}^{*}}{[(a'_{28})^{(5)}+(a''_{28})^{(5)}(T_{29}^{*})]}$, $G_{30}^{*} = \frac{(a_{30})^{(5)}G_{29}^{*}}{[(a'_{30})^{(5)}+(a''_{30})^{(5)}(T_{29}^{*})]}$

$T_{28}^{*} = \frac{(b_{28})^{(5)}T_{29}^{*}}{[(b'_{28})^{(5)}-(b''_{28})^{(5)}((G_{31})^{*})]}$, $T_{30}^{*} = \frac{(b_{30})^{(5)}T_{29}^{*}}{[(b'_{30})^{(5)}-(b''_{30})^{(5)}((G_{31})^{*})]}$ 519

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 520

G_{33}^{*} given by $\varphi((G_{35})^{*}) = 0$, T_{33}^{*} given by $f(T_{33}^{*}) = 0$ and

$$G_{32}^{*}=\frac{(a_{32})^{(6)}G_{33}^{*}}{[(a'_{32})^{(6)}+(a''_{32})^{(6)}(T_{33}^{*})]}\quad,\quad G_{34}^{*}=\frac{(a_{34})^{(6)}G_{33}^{*}}{[(a'_{34})^{(6)}+(a''_{34})^{(6)}(T_{33}^{*})]}$$

$$T_{32}^{*}=\frac{(b_{32})^{(6)}T_{33}^{*}}{[(b'_{32})^{(6)}-(b''_{32})^{(6)}((G_{35})^{*})]}\quad,\quad T_{34}^{*}=\frac{(b_{34})^{(6)}T_{33}^{*}}{[(b'_{34})^{(6)}-(b''_{34})^{(6)}((G_{35})^{*})]} \qquad 521$$

Obviously, these values represent an equilibrium solution of global equations

Finally we obtain the unique solution 522

G_{37}^{*} given by $\varphi((G_{39})^{*})=0$, T_{37}^{*} given by $f(T_{37}^{*})=0$ and

$$G_{36}^{*}=\frac{(a_{36})^{(7)}G_{37}^{*}}{[(a'_{36})^{(7)}+(a''_{36})^{(7)}(T_{37}^{*})]}\quad,\quad G_{38}^{*}=\frac{(a_{38})^{(7)}G_{37}^{*}}{[(a'_{38})^{(7)}+(a''_{38})^{(7)}(T_{37}^{*})]}$$

$$T_{36}^{*}=\frac{(b_{36})^{(7)}T_{37}^{*}}{[(b'_{36})^{(7)}-(b''_{36})^{(7)}((G_{39})^{*})]}\quad,\quad T_{38}^{*}=\frac{(b_{38})^{(7)}T_{37}^{*}}{[(b'_{38})^{(7)}-(b''_{38})^{(7)}((G_{39})^{*})]}$$

Finally we obtain the unique solution 523

G_{41}^{*} given by $\varphi((G_{43})^{*})=0$, T_{41}^{*} given by $f(T_{41}^{*})=0$ and

$$G_{40}^{*}=\frac{(a_{40})^{(8)}G_{41}^{*}}{[(a'_{40})^{(8)}+(a''_{40})^{(8)}(T_{41}^{*})]}\quad,\quad G_{42}^{*}=\frac{(a_{42})^{(8)}G_{41}^{*}}{[(a'_{42})^{(8)}+(a''_{42})^{(8)}(T_{41}^{*})]}$$

$$T_{40}^{*}=\frac{(b_{40})^{(8)}T_{41}^{*}}{[(b'_{40})^{(8)}-(b''_{40})^{(8)}((G_{43})^{*})]}\quad,\quad T_{42}^{*}=\frac{(b_{42})^{(8)}T_{41}^{*}}{[(b'_{42})^{(8)}-(b''_{42})^{(8)}((G_{43})^{*})]}$$

Finally we obtain the unique solution of 89 to 99 523 A

G_{45}^{*} given by $\varphi((G_{47})^{*})=0$, T_{45}^{*} given by $f(T_{45}^{*})=0$ and

$$G_{44}^{*}=\frac{(a_{44})^{(9)}G_{45}^{*}}{[(a'_{44})^{(9)}+(a''_{44})^{(9)}(T_{45}^{*})]}\quad,\quad G_{46}^{*}=\frac{(a_{46})^{(9)}G_{45}^{*}}{[(a'_{46})^{(9)}+(a''_{46})^{(9)}(T_{45}^{*})]}$$

$$T_{44}^{*}=\frac{(b_{44})^{(9)}T_{45}^{*}}{[(b'_{44})^{(9)}-(b''_{44})^{(9)}((G_{47})^{*})]}\quad,\quad T_{46}^{*}=\frac{(b_{46})^{(9)}T_{45}^{*}}{[(b'_{46})^{(9)}-(b''_{46})^{(9)}((G_{47})^{*})]}$$

ASYMPTOTIC STABILITY ANALYSIS 524

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(1)}$ *and* $(b_i'')^{(1)}$ Belong to $C^{(1)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof:_Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i=G_i^{*}+\mathbb{G}_i \qquad , T_i=T_i^{*}+\mathbb{T}_i$$

$$\frac{\partial(a''_{14})^{(1)}}{\partial T_{14}}(T_{14}^{*})=(q_{14})^{(1)}\quad,\quad \frac{\partial(b_i'')^{(1)}}{\partial G_j}(G^{*})=s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{13}}{dt} = -\left((a'_{13})^{(1)} + (p_{13})^{(1)}\right)\mathbb{G}_{13} + (a_{13})^{(1)}\mathbb{G}_{14} - (q_{13})^{(1)}G^*_{13}\mathbb{T}_{14} \quad 525$$

$$\frac{d\mathbb{G}_{14}}{dt} = -\left((a'_{14})^{(1)} + (p_{14})^{(1)}\right)\mathbb{G}_{14} + (a_{14})^{(1)}\mathbb{G}_{13} - (q_{14})^{(1)}G^*_{14}\mathbb{T}_{14} \quad 526$$

$$\frac{d\mathbb{G}_{15}}{dt} = -\left((a'_{15})^{(1)} + (p_{15})^{(1)}\right)\mathbb{G}_{15} + (a_{15})^{(1)}\mathbb{G}_{14} - (q_{15})^{(1)}G^*_{15}\mathbb{T}_{14} \quad 527$$

$$\frac{d\mathbb{T}_{13}}{dt} = -\left((b'_{13})^{(1)} - (r_{13})^{(1)}\right)\mathbb{T}_{13} + (b_{13})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(13)(j)}T^*_{13}\mathbb{G}_j\right) \quad 528$$

$$\frac{d\mathbb{T}_{14}}{dt} = -\left((b'_{14})^{(1)} - (r_{14})^{(1)}\right)\mathbb{T}_{14} + (b_{14})^{(1)}\mathbb{T}_{13} + \sum_{j=13}^{15}\left(s_{(14)(j)}T^*_{14}\mathbb{G}_j\right) \quad 529$$

$$\frac{d\mathbb{T}_{15}}{dt} = -\left((b'_{15})^{(1)} - (r_{15})^{(1)}\right)\mathbb{T}_{15} + (b_{15})^{(1)}\mathbb{T}_{14} + \sum_{j=13}^{15}\left(s_{(15)(j)}T^*_{15}\mathbb{G}_j\right) \quad 530$$

ASYMPTOTIC STABILITY ANALYSIS 531

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(2)}$ and $(b''_i)^{(2)}$ Belong to $C^{(2)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i \quad 532$$

$$\frac{\partial(a''_{17})^{(2)}}{\partial T_{17}}(T^*_{17}) = (q_{17})^{(2)} \quad , \frac{\partial(b''_i)^{(2)}}{\partial G_j}((G_{19})^*) = s_{ij} \quad 533$$

taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{16}}{dt} = -\left((a'_{16})^{(2)} + (p_{16})^{(2)}\right)\mathbb{G}_{16} + (a_{16})^{(2)}\mathbb{G}_{17} - (q_{16})^{(2)}G^*_{16}\mathbb{T}_{17} \quad 534$$

$$\frac{d\mathbb{G}_{17}}{dt} = -\left((a'_{17})^{(2)} + (p_{17})^{(2)}\right)\mathbb{G}_{17} + (a_{17})^{(2)}\mathbb{G}_{16} - (q_{17})^{(2)}G^*_{17}\mathbb{T}_{17} \quad 535$$

$$\frac{d\mathbb{G}_{18}}{dt} = -\left((a'_{18})^{(2)} + (p_{18})^{(2)}\right)\mathbb{G}_{18} + (a_{18})^{(2)}\mathbb{G}_{17} - (q_{18})^{(2)}G^*_{18}\mathbb{T}_{17} \quad 536$$

$$\frac{d\mathbb{T}_{16}}{dt} = -\left((b'_{16})^{(2)} - (r_{16})^{(2)}\right)\mathbb{T}_{16} + (b_{16})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(16)(j)}T^*_{16}\mathbb{G}_j\right) \quad 537$$

$$\frac{d\mathbb{T}_{17}}{dt} = -\left((b'_{17})^{(2)} - (r_{17})^{(2)}\right)\mathbb{T}_{17} + (b_{17})^{(2)}\mathbb{T}_{16} + \sum_{j=16}^{18}\left(s_{(17)(j)}T^*_{17}\mathbb{G}_j\right) \quad 538$$

$$\frac{d\mathbb{T}_{18}}{dt} = -\left((b'_{18})^{(2)} - (r_{18})^{(2)}\right)\mathbb{T}_{18} + (b_{18})^{(2)}\mathbb{T}_{17} + \sum_{j=16}^{18}\left(s_{(18)(j)}T^*_{18}\mathbb{G}_j\right) \quad 539$$

ASYMPTOTIC STABILITY ANALYSIS 540

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a''_i)^{(3)}$ *and* $(b''_i)^{(3)}$ Belong to $C^{(3)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G^*_i + \mathbb{G}_i \quad , T_i = T^*_i + \mathbb{T}_i$$

$$\frac{\partial(a_{21}'')^{(3)}}{\partial T_{21}}(T_{21}^*) = (q_{21})^{(3)} \quad , \frac{\partial(b_i'')^{(3)}}{\partial G_j}((G_{23})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{20}}{dt} = -\left((a_{20}')^{(3)} + (p_{20})^{(3)}\right)\mathbb{G}_{20} + (a_{20})^{(3)}\mathbb{G}_{21} - (q_{20})^{(3)}G_{20}^*\mathbb{T}_{21} \quad 541$$

$$\frac{d\mathbb{G}_{21}}{dt} = -\left((a_{21}')^{(3)} + (p_{21})^{(3)}\right)\mathbb{G}_{21} + (a_{21})^{(3)}\mathbb{G}_{20} - (q_{21})^{(3)}G_{21}^*\mathbb{T}_{21} \quad 542$$

$$\frac{d\mathbb{G}_{22}}{dt} = -\left((a_{22}')^{(3)} + (p_{22})^{(3)}\right)\mathbb{G}_{22} + (a_{22})^{(3)}\mathbb{G}_{21} - (q_{22})^{(3)}G_{22}^*\mathbb{T}_{21} \quad 543$$

$$\frac{d\mathbb{T}_{20}}{dt} = -\left((b_{20}')^{(3)} - (r_{20})^{(3)}\right)\mathbb{T}_{20} + (b_{20})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(20)(j)}T_{20}^*\mathbb{G}_j\right) \quad 544$$

$$\frac{d\mathbb{T}_{21}}{dt} = -\left((b_{21}')^{(3)} - (r_{21})^{(3)}\right)\mathbb{T}_{21} + (b_{21})^{(3)}\mathbb{T}_{20} + \sum_{j=20}^{22}\left(s_{(21)(j)}T_{21}^*\mathbb{G}_j\right) \quad 545$$

$$\frac{d\mathbb{T}_{22}}{dt} = -\left((b_{22}')^{(3)} - (r_{22})^{(3)}\right)\mathbb{T}_{22} + (b_{22})^{(3)}\mathbb{T}_{21} + \sum_{j=20}^{22}\left(s_{(22)(j)}T_{22}^*\mathbb{G}_j\right) \quad 546$$

ASYMPTOTIC STABILITY ANALYSIS 547

Theorem 4: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(4)}$ *and* $(b_i'')^{(4)}$ Belong to $C^{(4)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 548

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a_{25}'')^{(4)}}{\partial T_{25}}(T_{25}^*) = (q_{25})^{(4)} \quad , \frac{\partial(b_i'')^{(4)}}{\partial G_j}((G_{27})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{24}}{dt} = -\left((a_{24}')^{(4)} + (p_{24})^{(4)}\right)\mathbb{G}_{24} + (a_{24})^{(4)}\mathbb{G}_{25} - (q_{24})^{(4)}G_{24}^*\mathbb{T}_{25} \quad 549$$

$$\frac{d\mathbb{G}_{25}}{dt} = -\left((a_{25}')^{(4)} + (p_{25})^{(4)}\right)\mathbb{G}_{25} + (a_{25})^{(4)}\mathbb{G}_{24} - (q_{25})^{(4)}G_{25}^*\mathbb{T}_{25} \quad 550$$

$$\frac{d\mathbb{G}_{26}}{dt} = -\left((a_{26}')^{(4)} + (p_{26})^{(4)}\right)\mathbb{G}_{26} + (a_{26})^{(4)}\mathbb{G}_{25} - (q_{26})^{(4)}G_{26}^*\mathbb{T}_{25} \quad 551$$

$$\frac{d\mathbb{T}_{24}}{dt} = -\left((b_{24}')^{(4)} - (r_{24})^{(4)}\right)\mathbb{T}_{24} + (b_{24})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(24)(j)}T_{24}^*\mathbb{G}_j\right) \quad 552$$

$$\frac{d\mathbb{T}_{25}}{dt} = -\left((b_{25}')^{(4)} - (r_{25})^{(4)}\right)\mathbb{T}_{25} + (b_{25})^{(4)}\mathbb{T}_{24} + \sum_{j=24}^{26}\left(s_{(25)(j)}T_{25}^*\mathbb{G}_j\right) \quad 553$$

$$\frac{d\mathbb{T}_{26}}{dt} = -\left((b_{26}')^{(4)} - (r_{26})^{(4)}\right)\mathbb{T}_{26} + (b_{26})^{(4)}\mathbb{T}_{25} + \sum_{j=24}^{26}\left(s_{(26)(j)}T_{26}^*\mathbb{G}_j\right) \quad 554$$

ASYMPTOTIC STABILITY ANALYSIS 555

Theorem 5: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(5)}$ *and* $(b_i'')^{(5)}$ Belong to $C^{(5)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 556

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{29}'')^{(5)}}{\partial T_{29}}(T_{29}^*) = (q_{29})^{(5)} \ , \ \frac{\partial (b_i'')^{(5)}}{\partial G_j}((G_{31})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{28}}{dt} = -\left((a_{28}')^{(5)} + (p_{28})^{(5)}\right)\mathbb{G}_{28} + (a_{28})^{(5)}\mathbb{G}_{29} - (q_{28})^{(5)}G_{28}^*\mathbb{T}_{29} \quad 557$$

$$\frac{d\mathbb{G}_{29}}{dt} = -\left((a_{29}')^{(5)} + (p_{29})^{(5)}\right)\mathbb{G}_{29} + (a_{29})^{(5)}\mathbb{G}_{28} - (q_{29})^{(5)}G_{29}^*\mathbb{T}_{29} \quad 558$$

$$\frac{d\mathbb{G}_{30}}{dt} = -\left((a_{30}')^{(5)} + (p_{30})^{(5)}\right)\mathbb{G}_{30} + (a_{30})^{(5)}\mathbb{G}_{29} - (q_{30})^{(5)}G_{30}^*\mathbb{T}_{29} \quad 559$$

$$\frac{d\mathbb{T}_{28}}{dt} = -\left((b_{28}')^{(5)} - (r_{28})^{(5)}\right)\mathbb{T}_{28} + (b_{28})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(28)(j)}T_{28}^*\mathbb{G}_j\right) \quad 560$$

$$\frac{d\mathbb{T}_{29}}{dt} = -\left((b_{29}')^{(5)} - (r_{29})^{(5)}\right)\mathbb{T}_{29} + (b_{29})^{(5)}\mathbb{T}_{28} + \sum_{j=28}^{30}\left(s_{(29)(j)}T_{29}^*\mathbb{G}_j\right) \quad 561$$

$$\frac{d\mathbb{T}_{30}}{dt} = -\left((b_{30}')^{(5)} - (r_{30})^{(5)}\right)\mathbb{T}_{30} + (b_{30})^{(5)}\mathbb{T}_{29} + \sum_{j=28}^{30}\left(s_{(30)(j)}T_{30}^*\mathbb{G}_j\right) \quad 562$$

ASYMPTOTIC STABILITY ANALYSIS 563

Theorem 6: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(6)}$ *and* $(b_i'')^{(6)}$ Belong to $C^{(6)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 564

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{33}'')^{(6)}}{\partial T_{33}}(T_{33}^*) = (q_{33})^{(6)} \ , \ \frac{\partial (b_i'')^{(6)}}{\partial G_j}((G_{35})^*) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{32}}{dt} = -\left((a_{32}')^{(6)} + (p_{32})^{(6)}\right)\mathbb{G}_{32} + (a_{32})^{(6)}\mathbb{G}_{33} - (q_{32})^{(6)}G_{32}^*\mathbb{T}_{33} \quad 565$$

$$\frac{d\mathbb{G}_{33}}{dt} = -\left((a_{33}')^{(6)} + (p_{33})^{(6)}\right)\mathbb{G}_{33} + (a_{33})^{(6)}\mathbb{G}_{32} - (q_{33})^{(6)}G_{33}^*\mathbb{T}_{33} \quad 566$$

$$\frac{d\mathbb{G}_{34}}{dt} = -\left((a_{34}')^{(6)} + (p_{34})^{(6)}\right)\mathbb{G}_{34} + (a_{34})^{(6)}\mathbb{G}_{33} - (q_{34})^{(6)}G_{34}^*\mathbb{T}_{33} \quad 567$$

$$\frac{d\mathbb{T}_{32}}{dt} = -\left((b_{32}')^{(6)} - (r_{32})^{(6)}\right)\mathbb{T}_{32} + (b_{32})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(32)(j)}T_{32}^*\mathbb{G}_j\right) \quad 568$$

$$\frac{d\mathbb{T}_{33}}{dt} = -\left((b_{33}')^{(6)} - (r_{33})^{(6)}\right)\mathbb{T}_{33} + (b_{33})^{(6)}\mathbb{T}_{32} + \sum_{j=32}^{34}\left(s_{(33)(j)}T_{33}^*\mathbb{G}_j\right) \quad 569$$

$$\frac{d\mathbb{T}_{34}}{dt} = -\left((b_{34}')^{(6)} - (r_{34})^{(6)}\right)\mathbb{T}_{34} + (b_{34})^{(6)}\mathbb{T}_{33} + \sum_{j=32}^{34}\left(s_{(34)(j)}T_{34}^*\mathbb{G}_j\right) \quad 570$$

ASYMPTOTIC STABILITY ANALYSIS 571

Theorem 7: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(7)}$ *and* $(b_i'')^{(7)}$ Belong to $C^{(7)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 572

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{37}'')^{(7)}}{\partial T_{37}}(T_{37}^*) = (q_{37})^{(7)} \ , \ \frac{\partial (b_i'')^{(7)}}{\partial G_j}\left((G_{39})^{**} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain from

$$\frac{d\mathbb{G}_{36}}{dt} = -\left((a_{36}')^{(7)} + (p_{36})^{(7)}\right)\mathbb{G}_{36} + (a_{36})^{(7)}\mathbb{G}_{37} - (q_{36})^{(7)}G_{36}^*\mathbb{T}_{37} \quad 573$$

$$\frac{d\mathbb{G}_{37}}{dt} = -\left((a_{37}')^{(7)} + (p_{37})^{(7)}\right)\mathbb{G}_{37} + (a_{37})^{(7)}\mathbb{G}_{36} - (q_{37})^{(7)}G_{37}^*\mathbb{T}_{37} \quad 574$$

$$\frac{d\mathbb{G}_{38}}{dt} = -\left((a_{38}')^{(7)} + (p_{38})^{(7)}\right)\mathbb{G}_{38} + (a_{38})^{(7)}\mathbb{G}_{37} - (q_{38})^{(7)}G_{38}^*\mathbb{T}_{37} \quad 575$$

$$\frac{d\mathbb{T}_{36}}{dt} = -\left((b_{36}')^{(7)} - (r_{36})^{(7)}\right)\mathbb{T}_{36} + (b_{36})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(36)(j)}T_{36}^*\mathbb{G}_j\right) \quad 576$$

$$\frac{d\mathbb{T}_{37}}{dt} = -\left((b_{37}')^{(7)} - (r_{37})^{(7)}\right)\mathbb{T}_{37} + (b_{37})^{(7)}\mathbb{T}_{36} + \sum_{j=36}^{38}\left(s_{(37)(j)}T_{37}^*\mathbb{G}_j\right) \quad 578$$

$$\frac{d\mathbb{T}_{38}}{dt} = -\left((b_{38}')^{(7)} - (r_{38})^{(7)}\right)\mathbb{T}_{38} + (b_{38})^{(7)}\mathbb{T}_{37} + \sum_{j=36}^{38}\left(s_{(38)(j)}T_{38}^*\mathbb{G}_j\right) \quad 579$$

Obviously, these values represent an equilibrium solution

ASYMPTOTIC STABILITY ANALYSIS

Theorem 8: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(8)}$ *and* $(b_i'')^{(8)}$ Belong to $C^{(8)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:- 580

$$G_i = G_i^* + \mathbb{G}_i \quad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial (a_{41}'')^{(8)}}{\partial T_{41}}(T_{41}^*) = (q_{41})^{(8)} \ , \ \frac{\partial (b_i'')^{(8)}}{\partial G_j}\left((G_{43})^{*} \right) = s_{ij}$$

Then taking into account equations and neglecting the terms of power 2, we obtain

$$\frac{d\mathbb{G}_{40}}{dt} = -\left((a_{40}')^{(8)} + (p_{40})^{(8)}\right)\mathbb{G}_{40} + (a_{40})^{(8)}\mathbb{G}_{41} - (q_{40})^{(8)}G_{40}^*\mathbb{T}_{41} \quad 581$$

$$\frac{d\mathbb{G}_{41}}{dt} = -\left((a_{41}')^{(8)} + (p_{41})^{(8)}\right)\mathbb{G}_{41} + (a_{41})^{(8)}\mathbb{G}_{40} - (q_{41})^{(8)}G_{41}^*\mathbb{T}_{41} \quad 582$$

$$\frac{d\mathbb{G}_{42}}{dt} = -\left((a_{42}')^{(8)} + (p_{42})^{(8)}\right)\mathbb{G}_{42} + (a_{42})^{(8)}\mathbb{G}_{41} - (q_{42})^{(8)}G_{42}^*\mathbb{T}_{41} \quad 583$$

$$\frac{d\mathbb{T}_{40}}{dt} = -\left((b'_{40})^{(8)} - (r_{40})^{(8)}\right)\mathbb{T}_{40} + (b_{40})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(40)(j)}T^*_{40}\mathbb{G}_j\right) \quad 584$$

$$\frac{d\mathbb{T}_{41}}{dt} = -\left((b'_{41})^{(8)} - (r_{41})^{(8)}\right)\mathbb{T}_{41} + (b_{41})^{(8)}\mathbb{T}_{40} + \sum_{j=40}^{42}\left(s_{(41)(j)}T^*_{41}\mathbb{G}_j\right) \quad 585$$

$$\frac{d\mathbb{T}_{42}}{dt} = -\left((b'_{42})^{(8)} - (r_{42})^{(8)}\right)\mathbb{T}_{42} + (b_{42})^{(8)}\mathbb{T}_{41} + \sum_{j=40}^{42}\left(s_{(42)(j)}T^*_{42}\mathbb{G}_j\right) \quad 586$$

586

ASYMPTOTIC STABILITY ANALYSIS A

Theorem 9: If the conditions of the previous theorem are satisfied and if the functions $(a_i'')^{(9)}$ *and* $(b_i'')^{(9)}$ Belong to $C^{(9)}(\mathbb{R}_+)$ then the above equilibrium point is asymptotically stable.

Proof: Denote

Definition of $\mathbb{G}_i, \mathbb{T}_i$:-

$$G_i = G_i^* + \mathbb{G}_i \qquad , T_i = T_i^* + \mathbb{T}_i$$

$$\frac{\partial(a''_{45})^{(9)}}{\partial T_{45}}(T^*_{45}) = (q_{45})^{(9)} \quad , \frac{\partial(b''_i)^{(9)}}{\partial G_j}((G_{47})^*) = s_{ij}$$

Then taking into account equations 89 to 99 and neglecting the terms of power 2, we obtain from 99 to 44

$$\frac{d\mathbb{G}_{44}}{dt} = -\left((a'_{44})^{(9)} + (p_{44})^{(9)}\right)\mathbb{G}_{44} + (a_{44})^{(9)}\mathbb{G}_{45} - (q_{44})^{(9)}G^*_{44}\mathbb{T}_{45} \quad 586\,B$$

$$\frac{d\mathbb{G}_{45}}{dt} = -\left((a'_{45})^{(9)} + (p_{45})^{(9)}\right)\mathbb{G}_{45} + (a_{45})^{(9)}\mathbb{G}_{44} - (q_{45})^{(9)}G^*_{45}\mathbb{T}_{45} \quad 586\,C$$

$$\frac{d\mathbb{G}_{46}}{dt} = -\left((a'_{46})^{(9)} + (p_{46})^{(9)}\right)\mathbb{G}_{46} + (a_{46})^{(9)}\mathbb{G}_{45} - (q_{46})^{(9)}G^*_{46}\mathbb{T}_{45} \quad 586\,D$$

$$\frac{d\mathbb{T}_{44}}{dt} = -\left((b'_{44})^{(9)} - (r_{44})^{(9)}\right)\mathbb{T}_{44} + (b_{44})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(44)(j)}T^*_{44}\mathbb{G}_j\right) \quad 586\,E$$

$$\frac{d\mathbb{T}_{45}}{dt} = -\left((b'_{45})^{(9)} - (r_{45})^{(9)}\right)\mathbb{T}_{45} + (b_{45})^{(9)}\mathbb{T}_{44} + \sum_{j=44}^{46}\left(s_{(45)(j)}T^*_{45}\mathbb{G}_j\right) \quad 586\,F$$

$$\frac{d\mathbb{T}_{46}}{dt} = -\left((b'_{46})^{(9)} - (r_{46})^{(9)}\right)\mathbb{T}_{46} + (b_{46})^{(9)}\mathbb{T}_{45} + \sum_{j=44}^{46}\left(s_{(46)(j)}T^*_{46}\mathbb{G}_j\right) \quad 586\,G$$

The characteristic equation of this system is 587

$$\left((\lambda)^{(1)} + (b'_{15})^{(1)} - (r_{15})^{(1)}\right)\{\left((\lambda)^{(1)} + (a'_{15})^{(1)} + (p_{15})^{(1)}\right)$$

$$\left[\left(\left((\lambda)^{(1)} + (a'_{13})^{(1)} + (p_{13})^{(1)}\right)(q_{14})^{(1)}G^*_{14} + (a_{14})^{(1)}(q_{13})^{(1)}G^*_{13}\right)\right]$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(14)}T^*_{14} + (b_{14})^{(1)}s_{(13),(14)}T^*_{14}\right)$$

$$+\left(\left((\lambda)^{(1)} + (a'_{14})^{(1)} + (p_{14})^{(1)}\right)(q_{13})^{(1)}G^*_{13} + (a_{13})^{(1)}(q_{14})^{(1)}G^*_{14}\right)$$

$$\left(\left((\lambda)^{(1)} + (b'_{13})^{(1)} - (r_{13})^{(1)}\right)s_{(14),(13)}T^*_{14} + (b_{14})^{(1)}s_{(13),(13)}T^*_{13}\right)$$

$$\left(\left((\lambda)^{(1)}\right)^2 + \left((a'_{13})^{(1)} + (a'_{14})^{(1)} + (p_{13})^{(1)} + (p_{14})^{(1)}\right)(\lambda)^{(1)}\right)$$

$$\left((\lambda)^{(1)}\right)^2+\left((b'_{13})^{(1)}+(b'_{14})^{(1)}-(r_{13})^{(1)}+(r_{14})^{(1)}\right)(\lambda)^{(1)}\Big)$$

$$+\left(\left((\lambda)^{(1)}\right)^2+\left((a'_{13})^{(1)}+(a'_{14})^{(1)}+(p_{13})^{(1)}+(p_{14})^{(1)}\right)(\lambda)^{(1)}\right)(q_{15})^{(1)}G_{15}$$

$$+\left((\lambda)^{(1)}+(a'_{13})^{(1)}+(p_{13})^{(1)}\right)\left((a_{15})^{(1)}(q_{14})^{(1)}G^*_{14}+(a_{14})^{(1)}(a_{15})^{(1)}(q_{13})^{(1)}G^*_{13}\right)$$

$$\left((\lambda)^{(1)}+(b'_{13})^{(1)}-(r_{13})^{(1)}\right)s_{(14),(15)}T^*_{14}+(b_{14})^{(1)}s_{(13),(15)}T^*_{13}\Big)\}=0$$

$$+$$

$$(\lambda)^{(2)}+(b'_{18})^{(2)}-(r_{18})^{(2)}\}\{\left((\lambda)^{(2)}+(a'_{18})^{(2)}+(p_{18})^{(2)}\right)$$

$$\left[\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)\Big]$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(17)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(17)}\mathrm{T}^*_{17}\Big)$$

$$+\left(\left((\lambda)^{(2)}+(a'_{17})^{(2)}+(p_{17})^{(2)}\right)(q_{16})^{(2)}\mathrm{G}^*_{16}+(a_{16})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(16)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(16)}\mathrm{T}^*_{16}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$\left((\lambda)^{(2)}\right)^2+\left((b'_{16})^{(2)}+(b'_{17})^{(2)}-(r_{16})^{(2)}+(r_{17})^{(2)}\right)(\lambda)^{(2)}\Big)$$

$$+\left(\left((\lambda)^{(2)}\right)^2+\left((a'_{16})^{(2)}+(a'_{17})^{(2)}+(p_{16})^{(2)}+(p_{17})^{(2)}\right)(\lambda)^{(2)}\right)(q_{18})^{(2)}\mathrm{G}_{18}$$

$$+\left((\lambda)^{(2)}+(a'_{16})^{(2)}+(p_{16})^{(2)}\right)\left((a_{18})^{(2)}(q_{17})^{(2)}\mathrm{G}^*_{17}+(a_{17})^{(2)}(a_{18})^{(2)}(q_{16})^{(2)}\mathrm{G}^*_{16}\right)$$

$$\left((\lambda)^{(2)}+(b'_{16})^{(2)}-(r_{16})^{(2)}\right)s_{(17),(18)}\mathrm{T}^*_{17}+(b_{17})^{(2)}s_{(16),(18)}\mathrm{T}^*_{16}\Big)\}=0$$

$$+$$

$$(\lambda)^{(3)}+(b'_{22})^{(3)}-(r_{22})^{(3)}\}\{\left((\lambda)^{(3)}+(a'_{22})^{(3)}+(p_{22})^{(3)}\right)$$

$$\left[\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)(q_{21})^{(3)}G^*_{21}+(a_{21})^{(3)}(q_{20})^{(3)}G^*_{20}\right)\Big]$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(21)}T^*_{21}+(b_{21})^{(3)}s_{(20),(21)}T^*_{21}\Big)$$

$$+\left(\left((\lambda)^{(3)}+(a'_{21})^{(3)}+(p_{21})^{(3)}\right)(q_{20})^{(3)}G^*_{20}+(a_{20})^{(3)}(q_{21})^{(1)}G^*_{21}\right)$$

$$\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(20)}T^*_{21}+(b_{21})^{(3)}s_{(20),(20)}T^*_{20}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$\left((\lambda)^{(3)}\right)^2+\left((b'_{20})^{(3)}+(b'_{21})^{(3)}-(r_{20})^{(3)}+(r_{21})^{(3)}\right)(\lambda)^{(3)}\Big)$$

$$+\left(\left((\lambda)^{(3)}\right)^2+\left((a'_{20})^{(3)}+(a'_{21})^{(3)}+(p_{20})^{(3)}+(p_{21})^{(3)}\right)(\lambda)^{(3)}\right)(q_{22})^{(3)}G_{22}$$

$$+\left((\lambda)^{(3)}+(a'_{20})^{(3)}+(p_{20})^{(3)}\right)\left((a_{22})^{(3)}(q_{21})^{(3)}G^{*}_{21}+(a_{21})^{(3)}(a_{22})^{(3)}(q_{20})^{(3)}G^{*}_{20}\right)$$

$$\left(\left((\lambda)^{(3)}+(b'_{20})^{(3)}-(r_{20})^{(3)}\right)s_{(21),(22)}T^{*}_{21}+(b_{21})^{(3)}s_{(20),(22)}T^{*}_{20}\right)\}=0$$

$$+$$

$$(\lambda)^{(4)}+(b'_{26})^{(4)}-(r_{26})^{(4)}\{\left((\lambda)^{(4)}+(a'_{26})^{(4)}+(p_{26})^{(4)}\right)$$

$$\left[\left(\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)\right]$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(25)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(25)}T^{*}_{25}\right)$$

$$+\left(\left((\lambda)^{(4)}+(a'_{25})^{(4)}+(p_{25})^{(4)}\right)(q_{24})^{(4)}G^{*}_{24}+(a_{24})^{(4)}(q_{25})^{(4)}G^{*}_{25}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(24)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(24)}T^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$\left(\left((\lambda)^{(4)}\right)^{2}+\left((b'_{24})^{(4)}+(b'_{25})^{(4)}-(r_{24})^{(4)}+(r_{25})^{(4)}\right)(\lambda)^{(4)}\right)$$

$$+\left(\left((\lambda)^{(4)}\right)^{2}+\left((a'_{24})^{(4)}+(a'_{25})^{(4)}+(p_{24})^{(4)}+(p_{25})^{(4)}\right)(\lambda)^{(4)}\right)(q_{26})^{(4)}G_{26}$$

$$+\left((\lambda)^{(4)}+(a'_{24})^{(4)}+(p_{24})^{(4)}\right)\left((a_{26})^{(4)}(q_{25})^{(4)}G^{*}_{25}+(a_{25})^{(4)}(a_{26})^{(4)}(q_{24})^{(4)}G^{*}_{24}\right)$$

$$\left(\left((\lambda)^{(4)}+(b'_{24})^{(4)}-(r_{24})^{(4)}\right)s_{(25),(26)}T^{*}_{25}+(b_{25})^{(4)}s_{(24),(26)}T^{*}_{24}\right)\}=0$$

$$+$$

$$(\lambda)^{(5)}+(b'_{30})^{(5)}-(r_{30})^{(5)}\{\left((\lambda)^{(5)}+(a'_{30})^{(5)}+(p_{30})^{(5)}\right)$$

$$\left[\left(\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)\right]$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(29)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(29)}T^{*}_{29}\right)$$

$$+\left(\left((\lambda)^{(5)}+(a'_{29})^{(5)}+(p_{29})^{(5)}\right)(q_{28})^{(5)}G^{*}_{28}+(a_{28})^{(5)}(q_{29})^{(5)}G^{*}_{29}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(28)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(28)}T^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$\left(\left((\lambda)^{(5)}\right)^{2}+\left((b'_{28})^{(5)}+(b'_{29})^{(5)}-(r_{28})^{(5)}+(r_{29})^{(5)}\right)(\lambda)^{(5)}\right)$$

$$+\left(\left((\lambda)^{(5)}\right)^{2}+\left((a'_{28})^{(5)}+(a'_{29})^{(5)}+(p_{28})^{(5)}+(p_{29})^{(5)}\right)(\lambda)^{(5)}\right)(q_{30})^{(5)}G_{30}$$

$$+\left((\lambda)^{(5)}+(a'_{28})^{(5)}+(p_{28})^{(5)}\right)\left((a_{30})^{(5)}(q_{29})^{(5)}G^{*}_{29}+(a_{29})^{(5)}(a_{30})^{(5)}(q_{28})^{(5)}G^{*}_{28}\right)$$

$$\left(\left((\lambda)^{(5)}+(b'_{28})^{(5)}-(r_{28})^{(5)}\right)s_{(29),(30)}T^{*}_{29}+(b_{29})^{(5)}s_{(28),(30)}T^{*}_{28}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(6)}+(b'_{34})^{(6)}-(r_{34})^{(6)}\right)\{\left((\lambda)^{(6)}+(a'_{34})^{(6)}+(p_{34})^{(6)}\right)$$

$$\left[\left(\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)\right]$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(33)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(33)}T^{*}_{33}\right)$$

$$+\left(\left((\lambda)^{(6)}+(a'_{33})^{(6)}+(p_{33})^{(6)}\right)(q_{32})^{(6)}G^{*}_{32}+(a_{32})^{(6)}(q_{33})^{(6)}G^{*}_{33}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(32)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(32)}T^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$\left(\left((\lambda)^{(6)}\right)^{2}+\left((b'_{32})^{(6)}+(b'_{33})^{(6)}-(r_{32})^{(6)}+(r_{33})^{(6)}\right)(\lambda)^{(6)}\right)$$

$$+\left(\left((\lambda)^{(6)}\right)^{2}+\left((a'_{32})^{(6)}+(a'_{33})^{(6)}+(p_{32})^{(6)}+(p_{33})^{(6)}\right)(\lambda)^{(6)}\right)(q_{34})^{(6)}G_{34}$$

$$+\left((\lambda)^{(6)}+(a'_{32})^{(6)}+(p_{32})^{(6)}\right)\left((a_{34})^{(6)}(q_{33})^{(6)}G^{*}_{33}+(a_{33})^{(6)}(a_{34})^{(6)}(q_{32})^{(6)}G^{*}_{32}\right)$$

$$\left(\left((\lambda)^{(6)}+(b'_{32})^{(6)}-(r_{32})^{(6)}\right)s_{(33),(34)}T^{*}_{33}+(b_{33})^{(6)}s_{(32),(34)}T^{*}_{32}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(7)}+(b'_{38})^{(7)}-(r_{38})^{(7)}\right)\{\left((\lambda)^{(7)}+(a'_{38})^{(7)}+(p_{38})^{(7)}\right)$$

$$\left[\left(\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)\right]$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(37)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(37)}T^{*}_{37}\right)$$

$$+\left(\left((\lambda)^{(7)}+(a'_{37})^{(7)}+(p_{37})^{(7)}\right)(q_{36})^{(7)}G^{*}_{36}+(a_{36})^{(7)}(q_{37})^{(7)}G^{*}_{37}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(36)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(36)}T^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$\left(\left((\lambda)^{(7)}\right)^{2}+\left((b'_{36})^{(7)}+(b'_{37})^{(7)}-(r_{36})^{(7)}+(r_{37})^{(7)}\right)(\lambda)^{(7)}\right)$$

$$+\left(\left((\lambda)^{(7)}\right)^{2}+\left((a'_{36})^{(7)}+(a'_{37})^{(7)}+(p_{36})^{(7)}+(p_{37})^{(7)}\right)(\lambda)^{(7)}\right)(q_{38})^{(7)}G_{38}$$

$$+\left((\lambda)^{(7)}+(a'_{36})^{(7)}+(p_{36})^{(7)}\right)\left((a_{38})^{(7)}(q_{37})^{(7)}G^{*}_{37}+(a_{37})^{(7)}(a_{38})^{(7)}(q_{36})^{(7)}G^{*}_{36}\right)$$

$$\left(\left((\lambda)^{(7)}+(b'_{36})^{(7)}-(r_{36})^{(7)}\right)s_{(37),(38)}T^{*}_{37}+(b_{37})^{(7)}s_{(36),(38)}T^{*}_{36}\right)\}=0$$

$$+$$

$$\left((\lambda)^{(8)}+(b'_{42})^{(8)}-(r_{42})^{(8)}\right)\{\left((\lambda)^{(8)}+(a'_{42})^{(8)}+(p_{42})^{(8)}\right)$$

$$\left[\left(\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)\right]$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(41)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(41)}T^{*}_{41}\right)$$

$$+\left(\left((\lambda)^{(8)}+(a'_{41})^{(8)}+(p_{41})^{(8)}\right)(q_{40})^{(8)}G^{*}_{40}+(a_{40})^{(8)}(q_{41})^{(8)}G^{*}_{41}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(40)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(40)}T^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$\left(\left((\lambda)^{(8)}\right)^{2}+\left((b'_{40})^{(8)}+(b'_{41})^{(8)}-(r_{40})^{(8)}+(r_{41})^{(8)}\right)(\lambda)^{(8)}\right)$$

$$+\left(\left((\lambda)^{(8)}\right)^{2}+\left((a'_{40})^{(8)}+(a'_{41})^{(8)}+(p_{40})^{(8)}+(p_{41})^{(8)}\right)(\lambda)^{(8)}\right)(q_{42})^{(8)}G_{42}$$

$$+\left((\lambda)^{(8)}+(a'_{40})^{(8)}+(p_{40})^{(8)}\right)\left((a_{42})^{(8)}(q_{41})^{(8)}G^{*}_{41}+(a_{41})^{(8)}(a_{42})^{(8)}(q_{40})^{(8)}G^{*}_{40}\right)$$

$$\left(\left((\lambda)^{(8)}+(b'_{40})^{(8)}-(r_{40})^{(8)}\right)s_{(41),(42)}T^{*}_{41}+(b_{41})^{(8)}s_{(40),(42)}T^{*}_{40}\right)\}=0$$

$$+$$

$$(\lambda)^{(9)}+(b'_{46})^{(9)}-(r_{46})^{(9)}\{\left((\lambda)^{(9)}+(a'_{46})^{(9)}+(p_{46})^{(9)}\right)$$

$$\left[\left(\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)\right]$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(45)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(45)}T^{*}_{45}\right)$$

$$+\left(\left((\lambda)^{(9)}+(a'_{45})^{(9)}+(p_{45})^{(9)}\right)(q_{44})^{(9)}G^{*}_{44}+(a_{44})^{(9)}(q_{45})^{(9)}G^{*}_{45}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(44)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(44)}T^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$\left(\left((\lambda)^{(9)}\right)^{2}+\left((b'_{44})^{(9)}+(b'_{45})^{(9)}-(r_{44})^{(9)}+(r_{45})^{(9)}\right)(\lambda)^{(9)}\right)$$

$$+\left(\left((\lambda)^{(9)}\right)^{2}+\left((a'_{44})^{(9)}+(a'_{45})^{(9)}+(p_{44})^{(9)}+(p_{45})^{(9)}\right)(\lambda)^{(9)}\right)(q_{46})^{(9)}G_{46}$$

$$+\left((\lambda)^{(9)}+(a'_{44})^{(9)}+(p_{44})^{(9)}\right)\left((a_{46})^{(9)}(q_{45})^{(9)}G^{*}_{45}+(a_{45})^{(9)}(a_{46})^{(9)}(q_{44})^{(9)}G^{*}_{44}\right)$$

$$\left(\left((\lambda)^{(9)}+(b'_{44})^{(9)}-(r_{44})^{(9)}\right)s_{(45),(46)}T^{*}_{45}+(b_{45})^{(9)}s_{(44),(46)}T^{*}_{44}\right)\}=0$$

And as one sees, all the coefficients are positive. It follows that all the roots have negative real part, and this proves the theorem.

References:

(1) ^ a b C.B. Parker (1994). McGraw Hill Encyclopaedia of Physics (2nd ed.). McGraw Hill. p. 542. ISBN 0-07-051400-3.
(2) ^ C.B. Parker (1994). McGraw Hill Encyclopaedia of Physics (2nd ed.). McGraw Hill. p. 541. ISBN 0-07-051400-3.
(3) ^ a b c d e f g Bell, John (1964). "On the Einstein Podolsky Rosen Paradox". Physics 1 (3): 195–200.
(4) ^ Bohm, David Quantum Theory. Prentice–Hall, 1951.
(5) ^ a b c d Article on Bell's Theorem by Abner Shimony in the Stanford Encyclopedia of Philosophy, (2004).
(6) ^ Leggett, Anthony (2003). "Nonlocal Hidden-Variable Theories and Quantum Mechanics: An Incompatibility Theorem". Foundations of Physics 33 (10): 1469–1493.
(7) ^ Griffiths, David J. (1998). Introduction to Quantum Mechanics (2nd ed.). Pearson/Prentice Hall. pp. 423.
(8) ^ Merzbacher, Eugene (2005). Quantum Mechanics (3rd ed.). John Wiley & Sons. pp. 18, 362.
(9) ^ Stapp, 1975
(10) ^ Bell, JS, "On the impossible pilot wave." Foundations of Physics (1982) 12:989–99. Reprinted in Speakable and unspeakable in quantum mechanics: collected papers on quantum philosophy. CUP, 2004, p. 160.
(11) ^ Bell, JS, "On the impossible pilot wave." Foundations of Physics (1982) 12:989–99. Reprinted in Speakable and unspeakable in quantum mechanics: collected papers on quantum philosophy. CUP, 2004, p. 161.
(12) ^ Einstein, A.; Podolsky, B.; Rosen, N. (1935). "Can Quantum-Mechanical Description of Physical Reality Be Considered Complete?". Physical Review 47 (10): 777. Bibcode 1935PhRv...47..777E. doi:10.1103/PhysRev.47.777.
(13) ^ a b c d Clauser, John F. (1974). "Experimental consequences of objective local theories". Physical Review D 10 (2): 526. Bibcode 1974PhRvD..10..526C. doi:10.1103/PhysRevD.10.526.
(14) ^ Eberhard, P. H. (1977). "Bell's theorem without hidden variables". Nuovo Cimento B 38: 75–80. Bibcode 1977NCimB..38...75E. doi:10.1007/BF02726212.
(15) ^ Bell, JS, Speakable and unspeakable in quantum mechanics: Introduction remarks at Naples–Amalfi meeting., 1984. Reprinted in Speakable and unspeakable in quantum mechanics: collected papers on quantum philosophy. CUP, 2004, p. 29.
(16) ^ a b c Clauser, John; Horne, Michael; Shimony, Abner; Holt, Richard (1969). "Proposed Experiment to Test Local Hidden-Variable Theories". Physical Review Letters 23 (15): 880. Bibcode 1969PhRvL..23..880C. doi:10.1103/PhysRevLett.23.880.
(17) ^ M. Redhead, Incompleteness, Nonlocality and Realism, Clarendon Press (1987)
(18) ^ A. Einstein in Correspondance Einstein–Besso, p.265 (Herman, Paris, 1979)
(19) ^ Marshall and Santos, Semiclassical optics as an alternative to nonlocality Recent Research Developments in Optics 2:683–717 (2002) ISBN 81-7736-140-6
(20) ^ a b Freedman, Stuart J.; Clauser, John F. (1972). "Experimental Test of Local Hidden-Variable Theories". Physical Review Letters 28 (14): 938. Bibcode 1972PhRvL..28..938F. doi:10.1103/PhysRevLett.28.938.
(21) ^ Giorgio Brida; Marco Genovese; Marco Gramegna; Fabrizio Piacentini; Enrico Predazzi; Ivano Ruo-Berchera (2007). "Experimental tests of hidden variable theories from dBB to Stochastic Electrodynamics". Journal of Physics: Conference Series 67 (12047): 012047. arXiv:quant-ph/0612075. Bibcode 2007JPhCS..67a2047G. doi:10.1088/1742-6596/67/1/012047.
(22) ^ Gammaitoni, Luca; Hänggi, Peter; Jung, Peter; Marchesoni, Fabio (1998). "Stochastic resonance". Reviews of Modern Physics 70: 223. Bibcode 1998RvMP...70..223G. doi:10.1103/RevModPhys.70.223.
(23) ^ Gröblacher, Simon; Paterek, Tomasz; Kaltenbaek, Rainer; Brukner, Časlav; Żukowski, Marek; Aspelmeyer, Markus; Zeilinger, Anton (2007). "An experimental test of non-local realism". Nature 446 (7138): 871–5. arXiv:0704.2529. Bibcode 2007Natur.446..871G. doi:10.1038/nature05677. PMID 17443179.

(24)^ Cramer, John (1986). "The transactional interpretation of quantum mechanics". Reviews of Modern Physics 58 (3): 647. Bibcode 1986RvMP...58..647C. doi:10.1103/RevModPhys.58.647.
(25)^ Gerard 't Hooft (2009). "Entangled quantum states in a local deterministic theory". arXiv:0908.3408 [quant-ph].
(26)^ Gerard 't Hooft (2007). "The Free-Will Postulate in Quantum Mechanics". arXiv:quant-ph/0701097 [quant-ph].
(27)^ Jaynes, E. T. (1989). "Clearing up Mysteries—The Original Goal". Maximum Entropy and Bayesian Methods: 12.
(28)^ Roger Penrose (2007). The Road to Reality. Vintage books. p. 583. ISBN 0-679-77631-1.
(29)^ E. Abers (2004). Quantum Mechanics. Addison Wesley. pp. 193–195. ISBN 9780131461000.
(30)^ R.G. Lerner, G.L. Trigg (1991). Encyclopaedia of Physics (2nd ed.). VHC publishers. p. 495. ISBN 3-527-26954-1 (Verlagsgesellschaft) 0-89573-752-3 (VHC Inc.).
(31) A. Aspect et al., Experimental Tests of Realistic Local Theories via Bell's Theorem, Phys. Rev. Lett. 47, 460 (1981)
(32) A. Aspect et al., Experimental Realization of Einstein–Podolsky–Rosen–Bohm Gedankenexperiment: A New Violation of Bell's Inequalities, Phys. Rev. Lett. 49, 91 (1982).
(33) A. Aspect et al., Experimental Test of Bell's Inequalities Using Time-Varying Analyzers, Phys. Rev. Lett. 49, 1804 (1982).
(34) A. Aspect and P. Grangier, About resonant scattering and other hypothetical effects in the Orsay atomic-cascade experiment tests of Bell inequalities: a discussion and some new experimental data, Lettere al Nuovo Cimento 43, 345 (1985)
(35) B. D'Espagnat, The Quantum Theory and Reality, Scientific American, 241, 158 (1979)
(36) J. S. Bell, On the problem of hidden variables in quantum mechanics, Rev. Mod. Phys. 38, 447 (1966)
(37) J. S. Bell, On the Einstein Podolsky Rosen Paradox, Physics 1, 3, 195–200 (1964)
(38) J. S. Bell, Introduction to the hidden variable question, Proceedings of the International School of Physics 'Enrico Fermi', Course IL, Foundations of Quantum Mechanics (1971) 171–81
(39) J. S. Bell, Bertlmann's socks and the nature of reality, Journal de Physique, Colloque C2, suppl. au numero 3, Tome 42 (1981) pp C2 41–61
(40) J. S. Bell, Speakable and Unspeakable in Quantum Mechanics (Cambridge University Press 1987) [A collection of Bell's papers, including all of the above.]
(41) J. F. Clauser and A. Shimony, Bell's theorem: experimental tests and implications, Reports on Progress in Physics 41, 1881 (1978)
(42) J. F. Clauser and M. A. Horne, Phys. Rev D 10, 526–535 (1974)
(43) E. S. Fry, T. Walther and S. Li, Proposal for a loophole-free test of the Bell inequalities, Phys. Rev. A 52, 4381 (1995)
(44) E. S. Fry, and T. Walther, Atom based tests of the Bell Inequalities — the legacy of John Bell continues, pp 103–117 of Quantum [Un]speakables, R.A. Bertlmann and A. Zeilinger (eds.) (Springer, Berlin-Heidelberg-New York, 2002)
(45) R. B. Griffiths, Consistent Quantum Theory', Cambridge University Press (2002).
(46) L. Hardy, Nonlocality for 2 particles without inequalities for almost all entangled states. Physical Review Letters 71 (11) 1665–1668 (1993)
(47) M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information, Cambridge University Press (2000)
(48) P. Pearle, Hidden-Variable Example Based upon Data Rejection, Physical Review D 2, 1418–25 (1970)
(49) A. Peres, Quantum Theory: Concepts and Methods, Kluwer, Dordrecht, 1993.
(50) P. Pluch, Theory of Quantum Probability, PhD Thesis, University of Klagenfurt, 2006.
(51) B. C. van Frassen, Quantum Mechanics, Clarendon Press, 1991.
(52) M.A. Rowe, D. Kielpinski, V. Meyer, C.A. Sackett, W.M. Itano, C. Monroe, and D.J. Wineland, Experimental violation of Bell's inequalities with efficient detection,(Nature, 409, 791–794, 2001).
(53) S. Sulcs, The Nature of Light and Twentieth Century Experimental Physics, Foundations of Science 8, 365–391 (2003)

(54) S. Gröblacher et al., An experimental test of non-local realism,(Nature, 446, 871–875, 2007).
(55) D. N. Matsukevich, P. Maunz, D. L. Moehring, S. Olmschenk, and C. Monroe, Bell Inequality Violation with Two Remote Atomic Qubits, Phys. Rev. Lett. 100, 150404 (2008).
(56) The comic Dilbert, by Scott Adams, refers to Bell's Theorem in the 1992-09-21 and 1992-09-22 strips.
(57) ^ Oerter, Robert (2006-09-26). The Theory of Almost Everything: The Standard Model, the Unsung Triumph of Modern Physics (p. 2). Penguin Group. Kindle Edition.
(58) ^ S.L. Glashow (1961). "Partial-symmetries of weak interactions". Nuclear Physics 22 (4): 579–588. Bibcode 1961NucPh..22..579G. doi:10.1016/0029-5582(61)90469-2.
(59) ^ S. Weinberg (1967). "A Model of Leptons". Physical Review Letters 19 (21): 1264–1266. Bibcode 1967PhRvL..19.1264W. doi:10.1103/PhysRevLett.19.1264.
(60) ^ A. Salam (1968). N. Svartholm. ed. Elementary Particle Physics: Relativistic Groups and Analyticity. Eighth Nobel Symposium. Stockholm: Almquvist and Wiksell. pp. 367.
(61) ^ F. Englert, R. Brout (1964). "Broken Symmetry and the Mass of Gauge Vector Mesons". Physical Review Letters 13 (9): 321–323. Bibcode 1964PhRvL..13..321E. doi:10.1103/PhysRevLett.13.321.
(62) ^ P.W. Higgs (1964). "Broken Symmetries and the Masses of Gauge Bosons". Physical Review Letters 13 (16): 508–509. Bibcode 1964PhRvL..13..508H. doi:10.1103/PhysRevLett.13.508.
(63) ^ G.S. Guralnik, C.R. Hagen, T.W.B. Kibble (1964). "Global Conservation Laws and Massless Particles". Physical Review Letters 13 (20): 585–587. Bibcode 1964PhRvL..13..585G. doi:10.1103/PhysRevLett.13.585.
(64) ^ F.J. Hasert et al.; Faissner, H.; Krenz, W.; Von Krogh, J.; Lanske, D.; Morfin, J.; Schultze, K.; Weerts, H. et al. (1973). "Search for elastic muon-neutrino electron scattering". Physics Letters B 46: 121. Bibcode 1973PhLB...46..121H. doi:10.1016/0370-2693(73)90494-2.
(65) ^ F.J. Hasert et al.; Kabe, S.; Krenz, W.; Von Krogh, J.; Lanske, D.; Morfin, J.; Schultze, K.; Weerts, H. et al. (1973). "Observation of neutrino-like interactions without muon or electron in the gargamelle neutrino experiment". Physics Letters B 46: 138. Bibcode 1973PhLB...46..138H. doi:10.1016/0370-2693(73)90499-1.
(66) ^ F.J. Hasert et al.; Kabe, S.; Krenz, W.; Von Krogh, J.; Lanske, D.; Morfin, J.; Schultze, K.; Weerts, H. et al. (1974). "Observation of neutrino-like interactions without muon or electron in the Gargamelle neutrino experiment". Nuclear Physics B 73: 1. Bibcode 1974NuPhB..73....1H. doi:10.1016/0550-3213(74)90038-8.
(67) ^ D. Haidt (4 October 2004). "The discovery of the weak neutral currents". CERN Courier. Retrieved 8 May 2008.
(68) ^ "Details can be worked out if the situation is simple enough for us to make an approximation, which is almost never, but often we can understand more or less what is happening." from The Feynman Lectures on Physics, Vol 1. pp. 2–7
(69) ^ S. Braibant, G. Giacomelli, M. Spurio (2009). Particles and Fundamental Interactions: An Introduction to Particle Physics. Springer. pp. 313–314. ISBN 978-94-007-2463-1.
(70) ^ F. Englert, R. Brout (1964). "Broken Symmetry and the Mass of Gauge Vector Mesons". Physical Review Letters 13 (9): 321–323. Bibcode 1964PhRvL..13..321E. doi:10.1103/PhysRevLett.13.321.
(71) ^ P.W. Higgs (1964). "Broken Symmetries and the Masses of Gauge Bosons". Physical Review Letters 13 (16): 508–509. Bibcode 1964PhRvL..13..508H. doi:10.1103/PhysRevLett.13.508.
(72) ^ G.S. Guralnik, C.R. Hagen, T.W.B. Kibble (1964). "Global Conservation Laws and Massless Particles". Physical Review Letters 13 (20): 585–587. Bibcode 1964PhRvL..13..585G. doi:10.1103/PhysRevLett.13.585.
(73) ^ G.S. Guralnik (2009). "The History of the Guralnik, Hagen and Kibble development of the Theory of Spontaneous Symmetry Breaking and Gauge Particles". International Journal of Modern Physics A 24 (14): 2601–2627. arXiv:0907.3466. Bibcode 2009IJMPA..24.2601G. doi:10.1142/S0217751X09045431.
(74) ^ Lee, Benjamin W.; Quigg, C.; Thacker, H. B. (1977). "Weak interactions at very high energies: The role of the Higgs-boson mass". Physical Review D 16 (5): 1519–1531. Bibcode 1977PhRvD..16.1519L. doi:10.1103/PhysRevD.16.1519.

(75)^ "Huge $10 billion collider resumes hunt for 'God particle' - CNN.com". CNN. 11 November 2009. Retrieved 4 May 2010.
(76)^ http://profmattstrassler.com/articles-and-posts/the-higgs-particle/the-discovery-of-the-higgs/higgs-discovery-is-it-a-higgs
(77)^ "CERN experiments observe particle consistent with long-sought Higgs boson". CERN. 4 July 2012. Retrieved 4 July 2012.
(78)^ "Observation of a New Particle with a Mass of 125 GeV". Cms.web.cern.ch. Retrieved 2012-07-05.
(79)^ "ATLAS Experiment". Atlas.ch. 2006-01-01. Retrieved 2012-07-05.
(80)^ Video (04:38) - CERN Announcement (4 July 2012) Of Higgs Boson Discovery.
(81)^ Overbye, Dennis (July 4, 2012). "A New Particle Could Be Physics' Holy Grail". New York Times. Retrieved July 4, 2012.
(82)^ BABAR Data in Tension with the Standard Model (SLAC press-release).
(83)^ BaBar Collaboration, Evidence for an excess of B -> D(*) Tau Nu decays, arXiv:1205.5442.
(84)^ BaBar data hint at cracks in the Standard Model (EScienceNews.com).
(85)^ Bagdonaitel, Julija; Jensen, Paul; Henkel, Christian; Bethlem, Hendrick L.; Menten, Karl M.; Ubachs, Wim (December 13, 2012). "A Stringent Limit on a Drifting Proton-to-Electron Mass Ratio from Alcohol in the Early Universe". Science (journal) 339 (6115): 46. doi:10.1126/science.1224898. Retrieved December 14, 2012.
(86)^ Moskowitz, Clara (December 13, 2012). "Phew! Universe's Constant Has Stayed Constant". Space.com. Retrieved December 14, 2012.
(87)^ "CERN Press Release". Press.web.cern.ch. 2010-05-31. Retrieved 2012-07-05.
(88)^ S. Weinberg (1979). "Baryon and Lepton Nonconserving Processes". Physical Review Letters 43 (21): 1566. doi:10.1103/PhysRevLett.43.1566.
(89)^ P. Minkowski (1977). "mu --> e gamma at a Rate of One Out of 1-Billion Muon Decays?". Physics Letters B 67 (4): 421. Bibcode 1977PhLB...67..421M. doi:10.1016/0370-2693(77)90435-X.
(90)^ R. N. Mohapatra, G. Senjanovic (1980). "Neutrino Mass and Spontaneous Parity Nonconservation". Phys. Rev. Lett. 44 (14): 912–915. Bibcode 1980PhRvL..44..912M. doi:10.1103/PhysRevLett.44.912.
(91)^ M. Gell-Mann, P. Ramond and R. Slansky, in Supergravity, ed. by D. Freedman et al., North Holland (1979).
(92) News bulletins from CERM that came to me