

Antimicrobial Activity of *Catharanthus Roseus* .

Sheeraz ahmad wagay¹, Sudhansud Dhar Dwivedi², Manik Sharma³ Jagrati tripathi⁴, Mushtaq ahmad⁵

1. Govt. Benezee College Bhopal,
2. Bhoj Mahavadehaliya Bhopal.
3. Unique college Bhopal

Abstract

The aim of the present study is to investigate the antimicrobial activity and phytochemical analysis of Acetone extract of *Catharanthus roseus* whole plant against the wound isolates. Two different solvents such as ethanol and methanol were used to extract the bioactive compounds from the whole plant of *Catharanthus roseus* and screened for their antimicrobial activity against the isolated wound pathogens under well diffusion method. The maximum antibacterial activity was observed in crude Ethanolic extract of *Catharanthus roseus* against *Pseudomonas aeruginosa*. Qualitative analysis of phytochemical screening reveals the presence of Flavonoids, Tannin, Alkaloids and Terpenoids.

Keywords: - *Catharanthus roseus*, Antimicrobial activity and Phytochemical analysis

INTRODUCTION

A wound is a disruption in the normal anatomical structure and function of living tissue that can be caused by physical, chemical, Microbiological or immunological injury. Wounds represent a significant burden on the patients and health care professionals worldwide. Current estimates indicate that worldwide nearly 6 million people suffer from chronic wounds (Kumar et al., 2007). Chronic wounds may even lead to multiple organ failure or death of the patient. Wound infections often contain multiple organisms, including both aerobic and anaerobic gram-positive cocci and gram- negative bacilli and yeast. Organisms like Streptococcus sp, Staphylococcus aureus, Escherichia coli, Pseudomonas aeruginosa, Proteus sp, Klebsiella, Enterobacter, Clostridium, Peptostreptococcus, Fusobacterium and Aeromonas are highly predominant. (Henry, John, 2001). Many antibiotics used to treat wound pathogens but they also cause undesirable side effects. Bacterial resistance to antibiotic is a major therapeutic problem (Russell, 2002). Search towards safety medicines is must, scientists turn towards the herbal medicines to overcome side effects.

According to World Health Organization medicinal plants with various life sustaining constituents would be the best source to obtain a variety of potential, safe and novel drugs. (Natarajan et al., 2003). Bioactive compounds are exploited on a large scale because of their more systemic and no toxic effects. *Catharanthus roseus* L (apocyanaceae) also known as Vinca Rosea, is native to the Caribbean Basin and has historically been used to treat a wide assortment of diseases. European herbalists used the plant for conditions as varied as headache to a folk remedy for diabetes. It has more than 400 known alkaloids, some of which are approved as antineoplastic agents to treat leukemia, Hodgkin's disease, malignant lymphomas, neuroblastoma, rhabdomyosarcoma, Wilms' tumor, and other cancers.

Its vasodilating and memory-enhancing properties have been shown to alleviate vascular dementia and Alzheimer's disease .it also act as a wound healer (Fischhof et al., 1996 and Hindmarch et al., 1991). In our investigation an attempt was made to analyze the antibacterial activity of *Catharanthus roseus* against the clinical wound isolates.

MATERIAL AND METHODS

Plant Materials

Matured leaves of *Catharanthus roseus* were collected from Outside of Bhopal and identified by experts of Institute. Leaves were surface sterilized with 70% ethyl alcohol followed by 0.1% mercuric chloride. *Catharanthus roseus* leaves were shade dried and powdered with mechanical blender. The air dried plants powders (100 g) were successively extracted with water and alcohol. The extracts were dried in vacuum desiccator and were stored in a sterile container for further use (Kelmanson et al, 2000).

Determination antibacterial activity

Collection of Samples

Fifty Pus samples were collected from the Wound of patients admitted in Bhopal Hospital. Pus samples were collected from wall of an abscess with the help of sterile swab. Samples were transported to the laboratory for microbiological analysis.

Isolation and identification of wound samples

For the isolation of causative agents, the wound samples were inoculated in Blood agar and MacConkey agar. Plates were incubated at 37°C for 24-48 hrs. Colonies were analyzed by physiological and biochemical test for conformation (Konemanetal., 1998).

Agar Well Diffusion Assay

Agar well diffusion method was followed by using Muller-Hinton Agar (MHA). The plates were seeded with 18 hours old culture of the isolates. The organic fractions were dissolved in Dimethyl sulfoxide (DMSO) and sterilized by using sortorius syringe filter of pore size 0.22µm. Various concentrations of the extracts (250µl, 500µl, 750µl and 1000µl) were added into the sterile 8mm diameter well. Incubation was made at 37°C for 24hrs. Antibacterial activity was assayed by measuring the diameter of the inhibition zone formed around the well using standard (Hi-Media) scale. The experiment was repeated thrice and the average values were calculated for antibacterial activity (Perezetal., 1990)

Phytochemical screening

Ethanollic extracts were analyzed for the presence of alkaloids, saponins, triterpenes, and/or steroids, flavonoids, and tannins according to standard methods (Harborne, 1973)

Thin layer chromatography Preparation of TLC Plates

25x10 cm glass plates were washed with distilled water followed by smearing with acetone. After drying the plates were placed on the template in row. The slurry of silica gel G prepared with glass distilled water in the ratio 1:2 (w/v) was poured in the applicator. The glass plates were immediately coated with a layer of silica gel in 500µm thickness. The coated plates were activated at 80°C for 3 hours. Then the plates were stored in a plate chamber for further study. In that study chloroform and methanol (solvent) was used in 96:4 ratio.

Loading of substances:

The concentrated plant extract of 2.5 mg was loaded on the TLC plates just above 2 cm from the bottom using a capillary tube. The plates were reserved in a developing jar containing the solvent mixture. After, the solvent front reached approximately 18cm height. The plates were removed and allowed at room temperature for 30 min. Then the plates were also observed under UV light (240 and 300 nm) and recorded the R_f value of fluorescence substances (Anushia et al., 2009).

RESULTS & DISCUSSION

Antibacterial activity:

Different concentrations (250µg to 1000µg) of all the four extracts were recorded. Crude ethanolic fractions of *Catharanthus roseus* were tested against all the isolates. *Pseudomonas aeruginosa* (29mm), were highly sensitive to the ethanol fraction followed by *Staphylococcus aureus* (25mm), *Escherichia coli* (24mm), *Klebsiella pneumoniae* (18mm) and *Streptococcus pyogenes* (15mm). In case of Crude methanolic extract *Staphylococcus aureus* (25mm), *Klebsiella pneumoniae* (24mm), *Escherichia coli* (21mm), *Pseudomonas aeruginosa* (20mm), *Streptococcus pyogenes* (16mm) also shows their sensitivity. Soxhlet methanolic extract shows highest sensitivity against *Staphylococcus aureus* (16mm) followed by *Escherichia coli* (13mm), *Klebsiella pneumoniae* and *Streptococcus pyogenes* (12mm) *Pseudomonas aeruginosa* (9mm). Under Soxhlet ethanolic extraction sensitivity was recorded in *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Escherichia coli* and *Streptococcus pyogenes* as 22mm, 21mm, 18mm, 15mm and 12mm respectively (Table 1).

Phytochemical analysis:

Phytochemical such as Tannin, Flavonoids, Alkaloids, saponin and Terpenoids were reported from *Catharanthus roseus*. Alkaloids and flavonoids found to be present in all the four extracts whereas Terpenoids present in crude ethanol. Tannin shows their presence in Crude methanol and ethanol. Saponin present in soxhlet ethanolic and methanol. Piovani and Fillipini also reported the presence of various alkaloids, viz. Vincristine, Vinblastine, Yohimbine is an alkaloid and another flavonoids hirsutidin in *Catharanthus roseus* (Table 2).

Thin layer chromatography:

Crude ethanol (R_f 0.9) and soxhlet ethanol (0.77) shows brown colour spots whereas yellow spots were identified in crude methanol extracts. Olive green (R_f 0.15) spot was observed in soxhlet methanol extract (Table 3).

Table 1

Antibacterial activity of *Catharanthus roseus* against isolated wound Pathogens.

Sr. No	Organisms	Crude methanolic extract (ul)	Concentration of Extracts/Zone of inhibition in mm															
			50	75	100	25	50	75	100	25	50	75	100	25	50	75	100	
1.	Escherichia coli	250	50	75	100	25	50	75	100	25	50	75	100	25	50	75	100	
2.	Klebsiella pneumoniae	13	15	19	22	16	19	22	25	6	8	11	14	10	11	13	16	
3.	Pseudomonas aeruginosa	13	16	19	21	17	19	24	30	-	6	8	10	10	16	19	23	
4.	Staphylococcus	13	18	22	26	20	21	24	26	6	8	10	18	9	15	18	22	
5.	Streptococcus pyogens	9	13	15	17	10	11	13	16	6	8	11	13	-	-	7	13	

Table 2
Photochemical screening of *Catharanthus roseus* extracts

Sr. No	Phytochemicals	Crude methanolic	Crude ethanol	Soxhlet methanolic	Soxhlet ethanol extract
1.	Allcaloids	+	+	+	+
2.	Flavanoids	+	+	+	+
3.	Steroids	-	-	-	-
4.	Glyeosides	-	-	-	-
5.	Terpanoids	-	+	-	-
6.	Tannins	+	+	-	-
7.	Quinine	-	-	-	-
8.	Coumarin	-	-	-	-
9.	Starch	-	-	-	-
10.	Saponins	-	+	-	+
11.	Phenols	-	-	-	-

Table 3
Rf value of various extracts of *Catharanthus roseus* Leaves.

Sr. No	Extracts	Observation of sports	Rf value
1.	Crude Methanol	Yellow	0.4
2.	Crude Ethanol	Brown	0.9
3.	Soxhlet	Olive green	0.15
4.	Soxhlet Ethanol	Brown	0.77

REFERENCE

1. Anushia C, Sampathkumar P and Ram Kumar L: Antibacterial and antioxidant activity of *Cassia auriculata*. Global Journal of Pharmacology. 2009; 3(3): 127-130.

2. Fischhof PK, Moslinger-Gehmayr R, Hermann WM, Friedmann A and Russmann DL: Therapeutic efficacy of Vincamine in dementia. *Neuropsychobiology* 1996, 34(1): 29-35.
3. Harborne JB: *Phytochemical methods*. Harsted Press. New York 1973.
4. Henry and John B: *Clinical Diagnosis and Management by Laboratory Methods*, 20th ed. Philadelphia, WB Saunders Company. *J. Agri. Food. Chem.* 2001; 49(2): 1030-1034.
5. Hindmarch I, Fuchs HH and Erzigkeit H: Efficacy and tolerance of vinpocetine in ambulant patients suffering from mild to moderate organic psychosyndromes. *Int Clin Psychopharmacology* 1991; 6(1): S1-43.
6. Kelmanson JE, Jager AK and Staden JV: Zulu medicinal plants with antibacterial activity. *Journal of Pharmacology*. 2000; 69: 241-246.
7. Koneman, William EW, Janda M, Stephen D, Alien, Paul C, Schreeken B. Cashington C and Winn, JR: *Introduction to diagnostic microbiology in Laboratory and clinical diagnosis of infectious diseases*. J.B. Lippincott Company, Philadelphia 1998; 1-19.
8. Kumar B, Vijaya KM, Govinda RR and Pushpangadan P: Ethno pharmacological approaches to wound healing-exploring medicinal plants of India. *J. Ethno pharmacology* 2007; 114: 103-113.
9. Natarajan V, Venugopal PV and Menon T: Effect of azadirachta (neem) on the growth pattern of dermatophytes. *Indian J Med Microbiol* 2003; 21: 98-101.
10. Perez C, Paul M and Bazerque P. An antibiotic assay by the agar-well diffusion method. *Acta Biol. Med. Exp.* 1990; 15: 113-115.
11. Piovan A and Fillipini R: Anthocyanin in *Catharanthus roseus* *in vivo* and *in vitro*: A review. *Phytochem Rev* 2007; 6: 235-242.
12. Russell AD: Antibiotic and biocide resistance in bacteria: Introduction. *J. Appl Microbial. Symp. Supply* 2002; 2: 176-181.

This academic article was published by The International Institute for Science, Technology and Education (IISTE). The IISTE is a pioneer in the Open Access Publishing service based in the U.S. and Europe. The aim of the institute is Accelerating Global Knowledge Sharing.

More information about the publisher can be found in the IISTE's homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

The IISTE is currently hosting more than 30 peer-reviewed academic journals and collaborating with academic institutions around the world. There's no deadline for submission. **Prospective authors of IISTE journals can find the submission instruction on the following page:** <http://www.iiste.org/journals/> The IISTE editorial team promises to review and publish all the qualified submissions in a **fast** manner. All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Printed version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Recent conferences: <http://www.iiste.org/conference/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

