

Milk Yield and Composition of Red Sokoto Goats Fed Varying Levels of Baobab (*Adansonia digitata*) Pulp and Seed Meal

Okunlola D.O^{1*}, Olorunnisomo O.A², Aderinola O.A³, Nuga H.A¹ and Balogun N.O¹

1, Department of Animal Nutrition and Biotechnology, Ladoko Akintola University of Technology, P.M.B. 4000, Ogbomosho, Oyo State, Nigeria.

2, Department of Animal Science, University of Ibadan, Ibadan, Oyo State, Nigeria.

3, Department of Animal Production and Health, Ladoko Akintola University of Technology, P.M.B. 4000, Ogbomosho, Oyo State, Nigeria.

Abstract

The milk yield and mineral composition of Red Sokoto goats fed varying levels of baobab (*Adansonia digitata*) pulp and seed meal was investigated over a period of 84 days. Sixteen (16) lactating Red Sokoto goats averaging 19.75 ± 1.26 kg body weight were randomly allotted to four dietary treatments with four animals per replicate in a completely randomized design to determine the quantity and quality of milk produced. Treatment 1 was control with no baobab pulp and seed supplement, Treatments 2, 3 and 4 had 10%, 20% and 30% baobab pulp and seed inclusion respectively. The milk yield and mineral composition were significantly affected ($P < 0.05$). Milk yield progressively increased for all treatments to week four where the peak was recorded for the yield with 20% baobab pulp and seed inclusion levels having a clear significant value of 720.20g, followed by 30%, 10% above the control with varying values of 700.75g, 690.80 and 560.33g respectively. The milk yield recorded a decreased values of 475.00g, 455.67, 454.75 and 360.50 for 20%, 10%, 30% and 0% inclusion levels at the end of 12th week. There were significant difference ($P < 0.05$) in the values of Total solids, Fat and lactose. For milk composition; 30% inclusion level recorded the highest values for Total solids, fat and lactose at figures of 17.38%, 5.78% and 6.64% respectively. The fat content of the milk in the study were at the increasing range of 3.0 %, 4.05%, 4.60% and 5.78% respectively for inclusion levels of 0 – 30% baobab pulp and seed in the experimental diets. Milk produced by the experimental animals on 20% and 30% levels of inclusion recorded the overall best results in term of mineral composition. The results of the study had shown that inclusion of baobab at 20 % in the diet of Red Sokoto goats gave the highest milk yield . Inclusion at 30% level also gave better results in term of mineral composition.

Keywords: baobab pulp and seed, red sokoto goat, milk yield and mineral composition.

INTRODUCTION

The rising demand for milk and its products in the tropics has made it imperative to find means for increasing local milk production. The traditional dairy subsector is dominated by indigenous cattle breeds whose potential for milk production is largely underexploited (Olorunnisomo *et al.*, 2013). Even though cattle produces best milk in terms of quantity to meet the nutritional need of ever increasing human population; milk production by goats also cannot be overlooked.

Goat production plays a very vital role in the livelihood of rural populations in Nigeria as it contributes significantly to improvement of family nutrition and health. Also, sales of animals and their products help to stabilize household income. Goat husbandry, therefore, is considered as a form of food security and source of independent income for rural households and subsistent farmers. . The demand for animal products including milk have informed the need to raise indigenous goats even outside their natural habitat/environment; this practice helps in performance evaluation of animals in locations outside their domain (Ahamefule *et al.*, 2012).

The nutritional role of milk and milk products in human diet especially in developed countries has been well documented (Ibeawuchi *et al.* 2000). Milk from ruminants is a good alternative to augment animal protein intake. The West African Dwarf and the Red Sokoto goats are indigenous goat breeds found in southern and northern Nigeria, respectively. Records of their dairy performance in their natural environment exist (Ahamefule and Ibeawuchi, 2005; Akpa *et al.* 2002). However, their potentials for milk and meat production outside their natural ecology are continuously under survey (Akpa, 1999; Akpa *et al.* 2003; Ahamefule and Ibeawuchi, 2005). It is possible that these small ruminant breeds can do well, or even better, outside their natural habitat if well managed. Good management practices include adequate feeding of materials of great potentials for better performance. Therefore, feeding regime using non-conventional feedstuff such as baobab (*Adansonia digitata*) fruit is a pointer to accomplish this due to its beneficial health properties as antioxidant, analgesic, anti-inflammatory (Ramadan *et al.*, 1994), and good source of micronutrients and energy (Lockett *et al.*, 2000). Folktales among fulani herdsmen has it that Baobab fruit pulp enhances milk yield and milk letdown among Zebu cows.

1.1 MATERIAL AND METHODS

The study was conducted at the Small Ruminant Unit of Ladoko Akintola University of Technology Teaching and Research Farm. Sixteen lactating Red Sokoto goats (Does) were used for the study. Milk collection lasted for 84 days (12 weeks). The animals were allotted to four treatment comprising of four (4) animal per treatment in a Completely Randomized Experimental Design.

1.1.1 Preparation of experimental diets

The baobab fruits were gathered from a rural community in Ogbomoso area of Oyo State, Nigeria. The fruit pulp and seeds were removed and sun-dried for a week to reduce the anti nutrients. Dried baobab pulp and seeds was later milled and included at varying levels of 10%, 20% and 30% respectively with other feed component to prepare a concentrate tagged baobab pulp and seed meal (BPSM).

1.1.2 Chemical evaluation

Proximate analysis

Proximate composition of the whole baobab fruit, pulp, seeds, pulp and seed mixture as well as experimental diets were carried out according to the procedure of AOAC (1990). The crude protein was determined by the Kjeldahl method as described by AOAC (1990). Crude fiber determination was carried out using trichloroacetic acid (TCA) method. The ash and crude fat content were obtained by charring in furnace and extraction with ether

1.1.3

Experimental Animals – Sixteen (16) lactating Red Sokoto goats fed varying levels of baobab (*Adansonia digitata*) pulp and seed meal were used for the study in a 84 days milk collection.

Experimental Design – Completely Randomized Design

In a completely randomized design with four replicates of lactating doe per treatment, animals were randomly distributed to 4 treatments.

1.1.4 Milk Collection

Collection of milk from each lactating doe commenced 7 days after parturition to allow for colostrums intake by the kids. Milking procedure of Olorunnisomo et.al., 2012 in a study on Sokoto gudali cows was adopted in milk collection in the study. Milking was done by hand twice daily at 7am and 4pm in the presence of the kids for 84 days. Milk let-down was stimulated by the suckling the kid for 5 minutes after which the doe was milked. Weight of the kids was taken before and immediately after suckling to estimate milk intake. Kids were separated from does after milking and part of the milk collected was fed back to the kids using a nipple-bottle. Milk yield was calculated as the summation of milk off-take and intake by the kid



Kid – A week after parturition



A doe after milking



Kid feeding from bottle

1.1.5 RESULT AND DISCUSSION

Table 4 presents the average milk yield (g) over 90 days lactation in Red Sokoto goats fed baobab pulp and seed meal. The result is significantly ($P < 0.05$) different among the treatments for the twelve weeks of collection. Milk yield progressively increased for all treatments from week one to week four where the peak was recorded for the yield with 20% baobab pulp and seed inclusion levels having a clear significant value of 720.20g, followed by 30%, 10% above the control with varying values of 700.75g, 690.80 and 560.33g respectively. This is in agreement with the findings of Ahamefule *et al.*, 2012, Ahamefule and Ibeawuchi (2005) and Akpa *et al.* (2003) where peak production was recorded in West African Dwarf and Red Sokoto goats at the 4th week of lactation. The differences in milk yield may be attributed to feed intake and varying inclusion levels of baobab pulp and seed mixture in the experimental diets. Meanwhile, even though a decrease in yield was observed generally across the treatment at the fifth week of the study; animals on 20% inclusion level recorded the highest values in milk yield. The milk yield recorded a decreased values of 475.00g, 455.67, 454.75 and 360.50 for 20%, 10%, 30% and 0% inclusion levels at the end of 12th week. This could be attributed to gradual reduction in milk synthesis by the animals with reference to gradual offset of lactation.

The milk composition of Red Sokoto goats fed varying levels of Baobab (*Adansonia digitata*) Pulp and Seed Meal was presented in table 5. There were significant difference ($P < 0.05$) in the values of Total solids, Fat and lactose. For milk composition; 30% inclusion level recorded the highest values for Total solids, fat and lactose at figures of 17.38%, 5.78% and 6.64% respectively. It has been documented that the concentration of lactose in milk cannot be easily altered by nutrition. Ahamefule *et al.*, 2012. Ahamefule *et al.* (2003) recorded lactose concentrations of 4.46, 4.62 and 4.60% in early, mid and late lactation stages in West African Dwarf goat which affirm the relative consistency of lactose in milk. Lactose is a disaccharide synthesized in the udder. It is composed of a molecule of galactose joined to a molecule of glucose. Ash values were not influenced ($P > 0.05$) by graded level of baobab pulp and seed in the experimental diets. The values recorded were 0.80%, 0.84% 0.90% and 0.90% respectively for treatment 1 to 4. These values were within the range of 0.90 and 0.86% recorded by Ahamefule *et al.* (2004) and Ibeawuchi *et al.* (2003) for West African Dwarf and Red Sokoto goats, respectively.

The fat content of the milk in the study were at the increasing range of 3.0 %, 4.05%, 4.60% and 5.78% respectively for inclusion levels of 0 – 30% baobab pulp and seed in the experimental diets.. These values were in agreement with 5.04 % and 4.94% recorded by Zarhraddeen *et al.*, 2007. Alawa and Oji, 2008 also put fat content of Red sokoto goats in an experiment at 5.7%, Mba *et al.*, 1975, put it at 5.32% . Sankey, 1991 recorded 4.30 while Akinsoyinu *et al.*, 1981 and Ehoche and Buvanendran, 1983 reported 4.60% and 4.75% milk fat content respectively for the Red sokoto goats.

Table 6 presents the mineral composition of milk produced by Red Sokoto goats fed varying levels of

baobab pulp and seed meal. The results were significantly different ($P < 0.05$) among the minerals examined except for 30% inclusion level of the test ingredient. Generally, the mineral composition of goat milk is between 0.7-0.8%. Among macro-minerals, potassium accounts for the greatest percentage in milk, followed by calcium, phosphorus, sodium and magnesium (Ahamefule, 2012). Milk produced by animals on 20% and 30% levels of inclusion recorded the overall best results in terms of mineral composition. 30% inclusion of baobab pulp and seed recorded 0.25% for calcium, 0.33% for potassium. However, there was no significant difference ($P > 0.05$) in the value of magnesium as they both recorded 0.13. For sodium content however, 10 and 20% inclusion levels are not significantly different as the both values were put at 0.14%. Even though there was variation in phosphorus levels as recorded in the study; its variation in value had no significant difference ($P > 0.05$). Alawa and Oji (2008) reported 0.14 and 0.13%, potassium to phosphorus level respectively, for Red Sokoto goat which compares favorably with the results obtained in this study even though potassium level in this study was higher. Ahamefule, 2012 reported 0.15%, 0.09%, 0.11%, 0.11% and 0.13% for Calcium, Potassium, Magnesium, Sodium and Phosphorus respectively in a study on Milk yield and composition of Red Sokoto goats. The significant variation of mineral composition recorded in this study was attributed to high mineral composition of baobab fruit. Baobab fruit especially the seed has been reported to contain high level of minerals. Nkafamiya et al. (2007) reported phosphorus, calcium and potassium as the major mineral elements present in the seed.

CONCLUSION

The results of the study had shown that inclusion of baobab at 20% in the diet of Red Sokoto goats gave the highest milk yield. Inclusion at 30% level also gave better results in terms of mineral composition especially the macro minerals such as Calcium, Potassium, Magnesium, Sodium and Phosphorus which are essential nutrients needed by man and animal for a good health condition. Calcium for example is vital for the formation of strong bones and teeth and is also important in the maintenance of regular heartbeat and the transmission of nerve impulses. It is needed for muscle growth and contraction and for the prevention of muscle cramps. Also, Potassium is important for a healthy nervous system and a regular heart rhythm. It aids in proper muscle contraction, and works with sodium to control the body's water balance. The roles of Magnesium in human and animal diet cannot be over-emphasized. Magnesium is vital to enzyme activity. It assists in potassium uptake. A deficiency interferes with the transmission of nerve and muscle impulses, causing irritability and nervousness. It plays a role in the formation of bone and in carbohydrate and mineral metabolism. So also is Sodium which is necessary for maintenance of proper water balance and blood pH. It is also needed for stomach, nerve, and muscle function. Importance of Phosphorus cannot be left out as its presence in human and animal diet play a major function in the existence of these species. Phosphorus is needed for bone and tooth formation, cell growth and contraction of the heart muscle. It also assists in the utilization of vitamins and the conversion of food to energy. Baobab pulp and seed meal has been established in this study to contain all these essential nutrients and as such recommended for inclusion in ruminant diet.

In addition, baobab fruit plays a significant role in the diet of man meeting their daily vitamin C, Iron, and protein requirement. Its antioxidant capacity among other numerous health benefits has been documented. Therefore, considering the potentials of baobab fruit to supply these vital nutrients both to human and animals, its inclusion in ruminant diet no doubt will bring about the production of healthy animals and their products for human consumption.

The availability of baobab fruit in the dry season during which fresh forages are scarcely found; even where available; the nutritive value used to be low. Feeding this naturally rich in nutrient material; baobab fruit would thereby improve the performance of Red Sokoto goats and as such promote good health quality and their products for human consumption.

References

- Ahamefule F.O., Ibeawuchi J.A. and Ejiofor C.A. (2003). A comparative study of the constituents of cattle, sheep and goat milk in a hot humid environment. *Disc. Innov.* **15**(1/2), 64-68.
- Ahamefule F.O., Odilinye O and Nwachukwu E.N. (2012) *Milk Yield and Composition of Red Sokoto and West African Dwarf Does Raised Intensively in a Hot Humid Environment. Iranian Journal of Applied Animal Science* 2(2), 143-149
- Ahamefule F.O. and Ibeawuchi J.A. (2005). Milk yield and composition of West African Dwarf (WAD) does fed pigeon pea-cassava peel based diets. *J. Anim. Vet. Adv.* **4**(12), 991-999.
- Akinsoyinu A.O., Tewe O.O., Ngere L.O. and Mba A.U. (1981). Milk composition and yield of Red Sokoto (Maradi) goats. *Trop. Anim. Prod.* **6**, 186-187.
- Akpa G.N. (1999). Evaluation of dairy characteristics and lactation curves of small holder Red Sokoto (Maradi) goats. Ph. D. Thesis. Ahmadu Bello University, Zaria, Nigeria.
- Alawa J.P. and Oji U.I. (2008). Effect of pendulous udder enlargement on yield of Red Sokoto goats. *J. Anim. Vet. Adv.* **7**, 870-87.

- Akpa G.N., Osuhor C.U., Olugbemi T.S. and Nwani P.I. (2003). Milk flow rate and milking frequency in Red Sokoto goats. *Pakistan J. Nutr.* **2(3)**, 192-195.
- Ibeawuchi J.A. Chindo L. and Ahamefule F.O. (2000). Milk production characteristics of imported Holstein-Friesian cattle maintained in a tropical high altitude environment. *J. Sust. Agric. Emt.* **2**, 14-19.
- AOAC. (1990). Official Methods of Analysis. Vol.I. 15th Ed., Association of Official Analytical Chemists, Arlington, Virginia.
- Ramadan, F.M, Harraz, S.A, El-Maigy. (1994). Anti-inflammatory, anagelsic and anti-pyretic effects of the pulp of *Adansonia digitata*. *Fitoterapia*, 65(5):418-422.
- Ibeawuchi J.A., Ahamefule F.O. and Ringim I.A. (2003). The influence of lactation stage on the milk constituents of Sahelian goats. *Nigerian J. Anim. Prod.* **30(2)**, 259-264.
- Lockett, C.T, Calvert, C.C, Grivetti, L.E. (2000). Energy and micronutrient composition of dietary and medicinal wild plants consumed during drought. Study of rural Fulani, Northeastern Nigeria. *Int J food Sci Nutr*,51,195-208
- Mba A.U., Boyo S.S. and Oyenuga V.A. (1975). Studies on the milk composition of West African Dwarf, Red Sokoto and Saanen goats at different stages of lactation. *J. Dairy Res.* **42**, 217-226.
- Olorunnisomo O.A (2013). Milk production in Sokoto Gudali cows fed legume or elephant grass ensiled with cassava peel. *J. of Livestock Research for rural development* 25(6) 2013
- Zahraddeen, I.S., R. Butswat and S.T. Mbat, 2007. Evaluation of some factors affecting milk composition of indigenous goats in Nigeria. *Livest. Res. Rural Dev.* Vol. 19.

Table 1
Chemical composition of Baobab fruit (*Adansonia digitata*)

Parameters (%)	Whole fruit	Pulp only	Pulp and seed	Seed
Dry matter	90.39	90.39	89.88	89.92
Crude protein	7.53	3.50	13.38	17.33
Crude fiber	23.00	8.00	13.00	16.00
Ether extract	15.00	11.00	18.00	22.00
Ash	6.00	6.00	7.00	7.80
NFE	48.47	71.50	48.62	36.87
NDF	71.00	48.00	60.60	65.75
ADF	41.00	16.50	26.00	30.50
ADL	26.59	10.50	15.00	18.50

NDF- Neutral Detergent Fiber, ADF- Acid Detergent Fiber, ADL- Acid Detergent Lignin

Table 2 : Gross composition of the experimental diets

Ingredient (%)	Bb - 0	Bb - 10	Bb - 20	Bb - 30
Baobab pulp and seed	0.00	10.00	20.00	30.00
Wheat offal	63.00	53.00	43.00	33.00
Cassava peels	20.00	20.00	20.00	20.00
PKC	15.00	15.00	15.00	15.00
Premix	0.50	0.50	0.50	0.50
DCP	0.50	0.50	0.50	0.50
Salt	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00

Bb- 0% - Baobab pulp and seed, Bb- 20% - Baobab pulp and seed, Bb- 30% - Baobab pulp and seed, Bb- 10% - Baobab pulp and seed, PKC–Palm kernel cake, DCP – Di-calcium phosphate.

Table 3 : Chemical composition of experimental diets

Parameters (%)	Bb – 0	Bb – 10	Bb – 20	Bb – 30
Dry matter	82.57	83.07	83.98	81.92
Crude protein	14.50	14.10	13.70	13.30
Crude fiber	11.68	12.08	12.48	12.83
Ether extract	15.50	16.00	16.40	16.83
Ash	9.56	8.65	7.97	7.29
NFE	48.76	49.17	49.45	49.74
NDF	40.50	46.02	52.50	56.85
ADF	32.00	35.75	40.80	45.05
ADL	10.75	12.00	12.65	12.80
M.E (kcal/kg)	2013.70	2172.20	2220.70	2320.20

NFE- Nitrogen free extract, NDF- Neutral Detergent Fiber, ADF- Acid Detergent Fiber, ADL- Acid Detergent Lignin, Bb- 0% - Baobab pulp and seed, Bb- 20% - Baobab pulp and seed
 Bb- 30% - Baobab pulp and seed, Bb- 10% - Baobab pulp and seed

Table 4 Average weekly milk yield (g) over 84 days lactation period in Red Sokoto Goats fed varying levels of baobab (*Adansonia digitata*) pulp and seed meal

Week	Bb – 0	Bb - 10	Bb – 20	Bb - 30	±SEM
1	525.50	605.00	630.00	615.25	23.05
2	540.00	618.00	655.00	640.50	22.75
3	550.00	640.50	695.85	672.80	31.63
4	560.33	690.80	720.50	700.75	35.08
5	525.85	675.75	700.00	666.50	35.00
6	560.00	640.00	675.50	615.00	34.35
7	500.00	625.50	690.00	600.00	28.31
8	445.50	600.00	655.75	572.50	34.06
9	412.50	525.50	602.90	550.00	33.95
10	380.00	510.00	575.25	550.00	30.98
11	360.50	470.00	500.00	490.50	23.04
12	360.50	455.67	475.00	454.75	20.75

Table 5 : Milk composition of Red Sokoto Goats fed varying levels of Baobab (*Adansonia digitata*) Pulp and Seed Meal

Components (%)	Bb – 0	Bb - 10	Bb – 20	Bb – 30	SEM
Total solids	11.05 ^d	13.33 ^c	15.25 ^b	17.38 ^a	1.05
Protein	3.50 ^b	3.98 ^b	4.28 ^a	4.06 ^a	0.17
Fat	3.0 ^d	4.05 ^c	4.60 ^b	5.78 ^a	0.10
Ash	0.80	0.84	0.90	0.90	0.05
Lactose	3.75 ^d	4.46 ^c	5.47 ^b	6.64 ^a	0.15

^{abcd} means within the same row with different superscripts are significantly different ($P < 0.05$).

*Lactose was calculated as total solids – (protein + fat + ash)

Table 6: Mineral composition of Milk produced by Red Sokoto Goats fed varying levels of Baobab (*Adansonia digitata*) Pulp and Seed Meal

Minerals (%)	Bb - 0	Bb - 10	Bb – 20	Bb – 30	SEM
Calcium	0.15 ^d	0.18 ^c	0.20 ^b	0.25 ^a	0.001
Potassium	0.09 ^c	0.20 ^b	0.30 ^a	0.33 ^a	0.005
Magnesium	0.08 ^b	0.11 ^a	0.13 ^a	0.13 ^a	0.002
Sodium	0.10 ^c	0.14 ^b	0.14 ^b	0.16 ^a	0.02
Phosphorus	0.12 ^b	0.15 ^a	0.17 ^a	0.20 ^a	0.01

^{abcd} means within the same row with different superscripts are significantly different ($P < 0.05$).

Bb- 0% - Baobab pulp and seed

Bb- 20% - Baobab pulp and seed

Bb- 30% - Baobab pulp and seed

Bb- 10% - Baobab pulp and seed

The IISTE is a pioneer in the Open-Access hosting service and academic event management. The aim of the firm is Accelerating Global Knowledge Sharing.

More information about the firm can be found on the homepage:

<http://www.iiste.org>

CALL FOR JOURNAL PAPERS

There are more than 30 peer-reviewed academic journals hosted under the hosting platform.

Prospective authors of journals can find the submission instruction on the following page: <http://www.iiste.org/journals/> All the journals articles are available online to the readers all over the world without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself. Paper version of the journals is also available upon request of readers and authors.

MORE RESOURCES

Book publication information: <http://www.iiste.org/book/>

Academic conference: <http://www.iiste.org/conference/upcoming-conferences-call-for-paper/>

IISTE Knowledge Sharing Partners

EBSCO, Index Copernicus, Ulrich's Periodicals Directory, JournalTOCS, PKP Open Archives Harvester, Bielefeld Academic Search Engine, Elektronische Zeitschriftenbibliothek EZB, Open J-Gate, OCLC WorldCat, Universe Digital Library, NewJour, Google Scholar

