



On Jordan (σ,τ) - Higher Reverse Derivations of Gamma-Rings

Fawaz Raad Jarullah

Department of Mathematics ,college of Education , Al-Mustansirya University , Iraq

Abstract:

Let M be a Γ -ring and σ^n, τ^n be two higher endomorphisms of a Γ -ring M , for all $n \in N$ in the present paper we show that under certain conditions of M , every Jordan (σ,τ) -higher reverse derivation of a Γ -Ring M is a (σ,τ) -higher reverse derivation

Mathematics Subject Classification(2000) : 16W25, 16N80 , 16U80

Key Words: derivation , reverse derivation , higher reverse derivation , Jordan higher reverse derivation

1- Introduction:

Let M and Γ be two additive abelian groups, suppose that there is a mapping from $M \times \Gamma \times M \rightarrow M$ (the image of (a, α, b) being denoted by $a\alpha b$, $a, b \in M$ and $\alpha \in \Gamma$). Satisfying for all $a, b, c \in M$ and $\alpha, \beta \in \Gamma$:

$$(i) \quad (a + b)\alpha c = a\alpha c + b\alpha c$$

$$a(\alpha + \beta)c = a\alpha c + a\beta c$$

$$a\alpha(b + c) = a\alpha b + a\alpha c$$

$$(ii) \quad (a\alpha b)\beta c = a\alpha(b\beta c)$$

Then M is called a Γ -ring. This definition is due to Barnes [1].

Let M be a Γ -ring then M is called 2-torsion free if $2a = 0$ implies $a = 0$, for every $a \in M$, this definition is due to [3].

Let M be a Γ -ring and $d: M \rightarrow M$ be an additive mapping (that is $d(a + b) = d(a) + d(b)$) of a Γ -ring M into itself then d is called a derivation on M if : $d(a\alpha b) = d(a)\alpha b + a\alpha d(b)$, for all $a, b \in M$ and $\alpha \in \Gamma$ and d is called a Jordan derivation on M if $d(a\alpha a) = d(a)\alpha a + a\alpha d(a)$, for all $a \in M$ and $\alpha \in \Gamma$, [2].

Let M be a Γ -ring and σ, τ be tow endomorphisms of M . such that $d: M \rightarrow M$ be an additive mapping. Then d is called (σ, τ) -derivation of M if:

$$d(a\alpha b) = d(a)\alpha \tau(b) + \sigma(a)\alpha d(b), \text{ for all } a, b \in M, \alpha \in \Gamma.$$

And d is called a Jordan (σ, τ) -derivation of M if:

$$d(a\alpha a) = d(a)\alpha\tau(a) + \sigma(a)\alpha d(a), \text{ for all } a \in M, \alpha \in \Gamma, [5].$$

Let M be a Γ -ring and $d: M \rightarrow M$ be an additive mapping of a Γ -ring M into itself then d is called reverse derivation on M if

$$d(aab) = d(b)\alpha a + b\alpha d(a), \text{ for all } a, b \in M \text{ and } \alpha \in \Gamma.$$

Let M be a Γ -ring and $d: M \rightarrow M$ be an additive mapping of a Γ -ring M into itself then d is called a Jordan reverse derivation on M if

$$d(a\alpha a) = d(a)\alpha a + a\alpha d(a), \text{ for all } a \in M \text{ and } \alpha \in \Gamma, [4].$$

Let M be a Γ -ring and $D = (d_i)_{i \in N}$ be a family of additive mappings of M , such that $d_0 = id_M$ then D is called a higher reverse derivation on M if for every $a, b \in M, \alpha \in \Gamma$ and $n \in N$

$$d_n(aab) = \sum_{i+j=n} d_i(b)\alpha d_j(a)$$

And D is called a Jordan higher reverse derivation on M if for every $a \in M, \alpha \in \Gamma$ and $n \in N$.

$$d_n(a\alpha a) = \sum_{i+j=n} d_i(a)\alpha d_j(a), [6].$$

Now, the main purpose of this paper is that every Jordan (σ, τ) -higher reverse derivation of a 2-torsion free Γ -ring M into itself, such that $aab\beta a = a\beta baa$, for all $a, b \in M$ and $\alpha, \beta \in \Gamma$ is a Jordan triple (σ, τ) -higher reverse derivation .

2- Jordan (σ, τ) -Higher Reverse Derivations on Γ -Ring :

Definition (2.1):

Let $D = (d_i)_{i \in N}$ be a family of additive mappings of a Γ -ring M into itself, such that $d_0 = id_M$ and σ, τ be two endomorphisms of M . D is called (σ, τ) -higher reverse derivation if

$$d_n(aab) = \sum_{i+j=n} d_i(\sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(a)), \text{ for all } a, b \in M, \alpha \in \Gamma \text{ and } n \in N.$$

Example (2.2):

Let R be a ring and $d = (d_i)_{i \in N}$ be a (σ, τ) -higher reverse derivation on R . Let $M = M_{1 \times 2}(R)$ and $\Gamma = \left\{ \begin{pmatrix} n \\ 0 \end{pmatrix}, n \in N \right\}$. Then M is a Γ -ring. We define $D = (D_i)_{i \in N}$ be a family of

additive mappings of M such that $D_n((a \quad b)) = ((d_n(a) \quad d_n(b))$

Let σ_1^n, τ_1^n be two endomorphisms of M , such that $\sigma_1^n((a \quad b)) = ((\sigma(a) \quad \sigma(b)),$

$\tau_1^n((a \quad b)) = ((\tau(a) \quad \tau(b)))$

Then D is a (σ, τ) -higher reverse derivation.

Definition (2.3):

Let $D = (d_i)_{i \in N}$ be a family of additive mappings of a Γ -ring M into itself, such that $d_0 = id_M$ and σ, τ be two endomorphisms of M . D is called Jordan (σ, τ) -higher reverse derivation if

$$d_n(a \alpha a) = \sum_{i+j=n} d_i(\sigma^{n-i}(a)) \text{ad}_j(\tau^{n-j}(a))$$

for all $a \in M$, $\alpha \in \Gamma$ and $n \in N$.

Definition (2.4):

Let $D = (d_i)_{i \in N}$ be a family of additive mappings of a Γ -ring M into itself, such that $d_0 = id_M$ and σ, τ be two endomorphisms of M . D is called Jordan triple (σ, τ) -higher reverse derivation if

$$d_n(a \alpha b \beta a) = d_n(a) \beta a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a))$$

for all $a, b \in M$, $\alpha, \beta \in \Gamma$ and $n \in N$.

Lemma (2.5):

Let $D = (d_i)_{i \in N}$ be a Jordan triple (σ, τ) -higher reverse derivations on a Γ -ring M into itself. Then for all $a, b, c \in M$, $\alpha, \beta \in \Gamma$ and $n \in N$

- (i) $d_n(a \alpha b + b \alpha a) = \sum_{i+j=n} d_i(\sigma^{n-i}(b)) \text{ad}_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a)) \text{ad}_j(\tau^{n-j}(b))$
- (ii) $d_n(a \alpha b \beta a + a \beta b \alpha a) = d_n(a) \beta a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a)) +$
 $d_n(a) \alpha a \beta b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \text{ad}_j(\sigma^k \tau^i(b)) \beta d_k(\tau^{n-k}(a))$

(iii) If M is a 2-torsion free Γ -ring.

- $d_n(a \alpha b \alpha a) = d_n(a) \alpha a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \text{ad}_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a))$
- (iv) $d_n(a \alpha b \beta c + c \alpha b \beta a) = d_n(c) \beta a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c)) \beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a)) +$
 $d_n(a) \beta c \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(c))$

(v) In particular, if M is a 2-torsion free commutative Γ -ring

- $d_n(a \alpha b \beta c) = d_n(c) \beta a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c)) \beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a))$
- (vi) $d_n(a \alpha b \alpha c + c \alpha b \alpha a) = d_n(c) \alpha a \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c)) \alpha d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a)) +$
 $d_n(a) \alpha c \alpha b + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a)) \alpha d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a))$

Proof:

(i) Replacing $a + b$ for a in the Definition (2.3), we get:

$$\begin{aligned}
 d_n((a+b)\alpha(a+b)) &= \sum_{i+j=n} d_i(\sigma^{n-i}(a+b))\alpha d_j(\tau^{n-j}(a+b)) \\
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(a) + \sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(a) + \tau^{n-j}(b)) \\
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(b)) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(b)) \\
 &\quad \dots(1)
 \end{aligned}$$

On the other hand:

$$\begin{aligned}
 d_n((a+b)\alpha(a+b)) &= d_n(a\alpha a + a\alpha b + b\alpha a + b\alpha b) \\
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(b)) + \dots(2) \\
 &\quad d_n(a\alpha b + b\alpha a)
 \end{aligned}$$

Comparing (1) and (2), we get:

$$d_n(a\alpha b + b\alpha a) = \sum_{i+j=n} d_i(\sigma^{n-i}(b))\alpha d_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(b))$$

(ii) Replace $a\beta b + b\beta a$ for b in (i), we get:

$$\begin{aligned}
 &d_n(a\alpha(a\beta b + b\beta a) + (a\beta b + b\beta a)\alpha a) \\
 &= d_n(a\alpha(a\beta b) + a\alpha(b\beta a) + (a\beta b)\alpha a + (b\beta a)\alpha a) \\
 &= d_n((a\alpha a)\beta b + (a\alpha b)\beta a + (a\beta b)\alpha a + (b\beta a)\alpha a) \\
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))\beta d_j(\tau^{n-j}(a\alpha a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\beta d_j(\tau^{n-j}(a\alpha b)) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(a\beta a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(b\beta a))
 \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))\beta(\sum_{r+s=j} d_r(\sigma^{j-r}\tau^{n-j}(a))ad_s(\tau^{j-s}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\beta(\sum_{e+f=j} d_e(\sigma^{j-e}\tau^{n-j}(b))ad_f(\tau^{j-f}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha(\sum_{p+q=j} d_p(\sigma^{j-p}\tau^{n-j}(b))\beta d_q(\tau^{j-q}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha(\sum_{x+y=j} d_x(\sigma^{j-x}\tau^{n-j}(a))\beta d_y(\tau^{j-y}\tau^{n-j}(b))) \\
 &= \sum_{i+r+s=n} d_i(\sigma^{n-i}(b))\beta d_r(\sigma^s\tau^i(a))ad_s(\tau^{n-s}(a)) + \\
 &\quad \sum_{i+e+f=n} d_i(\sigma^{n-i}(a))\beta d_e(\sigma^f\tau^i(b))ad_f(\tau^{n-f}(a)) + \\
 &\quad \sum_{i+p+q=n} d_i(\sigma^{n-i}(a))ad_p(\sigma^q\tau^i(b))\beta d_q(\tau^{n-q}(a)) + \\
 &\quad \sum_{i+x+y=n} d_i(\sigma^{n-i}(a))ad_x(\sigma^y\tau^i(a))\beta d_y(\tau^{n-y}(b)) \\
 &= d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k\tau^i(a))ad_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\beta a\alpha b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))ad_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha a\beta b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))ad_j(\sigma^k\tau^i(b))\beta d_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha b\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))ad_j(\sigma^k\tau^i(a))\beta d_k(\tau^{n-k}(b))
 \end{aligned} \tag{1}$$

On the other hand:

$$\begin{aligned}
 &d_n(a\alpha(a\beta b + b\beta a) + (a\beta b + b\beta a)\alpha a) \\
 &= d_n(a\alpha a\beta b + a\alpha b\beta a + a\beta b\alpha a + b\beta a\alpha a) \\
 &= d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k\tau^i(a))ad_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha b\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))ad_j(\sigma^k\tau^i(a))\beta d_k(\tau^{n-k}(b)) + d_n(a\alpha b\beta a + a\beta b\alpha a)
 \end{aligned} \tag{2}$$

Comparing (1) and (2), we get:

$$\begin{aligned}
 &d_n(a\alpha b\beta a + a\beta b\alpha a) = \\
 &= d_n(a)\beta a\alpha b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))ad_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha a\beta b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))ad_j(\sigma^k\tau^i(b))\beta d_k(\tau^{n-k}(a))
 \end{aligned}$$

(iii) Replace α for β in (ii), we get:

$$d_n(aab\alpha a + aab\alpha a) = 2d_n(aab\alpha a)$$

Since M is a 2-torsion free Γ -ring

$$= d_n(a)\alpha a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a))\alpha d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a))$$

(iv) Replace $a+c$ for a in Definition (2.4), we get:

$$\begin{aligned} d_n((a+c)ab\beta(a+c)) &= d_n(a+c)\beta(a+c)ab + \\ &\quad \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a+c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-j}(a+c)) \\ &= d_n(a)\beta a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-j}(a)) + \\ &\quad d_n(c)\beta a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-j}(a)) + \\ &\quad d_n(a)\beta c ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-j}(c)) + \\ &\quad d_n(c)\beta c ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-j}(c)) \end{aligned} \dots(1)$$

On the other hand

$$\begin{aligned} d_n((a+c)ab\beta(a+c)) &= d_n(aab\beta a + aab\beta c + cab\beta a + cab\beta c) \\ &= d_n(a)\beta a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a)) + \\ &\quad d_n(c)\beta c ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(c)) + d_n(aab\beta c + cab\beta a) \end{aligned} \dots(2)$$

Compare (1) and (2), we get:

$$\begin{aligned} d_n(aab\beta c + cab\beta a) &= d_n(c)\beta a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a)) + \\ &\quad d_n(a)\beta c ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(c)) \end{aligned}$$

(v) By (iv) and since M is a 2-torsion free commutative Γ -ring, we get:

$$\begin{aligned} d_n(aab\beta c + aab\beta c) &= 2d_n(aab\beta c) \\ &= d_n(c)\beta a ab + \sum_{i+j+k=n}^{i < n} d_i(\sigma^{n-i}(c))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a)) \end{aligned}$$

(vi) Replace α for β in (iv), we get:

$$\begin{aligned} d_n(aabac + cabaa) &= d_n(c)aaa + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(c))ad_j(\sigma^k\tau^i(b))ad_k(\tau^{n-k}(a)) + \\ &\quad d_n(a)acab + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))ad_j(\sigma^k\tau^i(b))ad_k(\tau^{n-k}(c)) \end{aligned}$$

Definition (2.6):

Let $D = (d_i)_{i \in N}$ be a Jordan (σ, τ) -higher reverse derivation of a Γ -ring M into itself, then for all $a, b \in M$, $\alpha \in \Gamma$ and $n \in N$, we define

$$\phi_n = d_n(aab) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a))$$

Lemma (2.7):

Let $D = (d_i)_{i \in N}$ be a Jordan (σ, τ) -higher reverse derivation of a Γ -ring M into itself, then for all $a, b, c \in M$, $\alpha, \beta \in \Gamma$ and $n \in N$:

- (i) $\phi_n(a,b)_\alpha = -\phi_n(b,a)_\alpha$
- (ii) $\phi_n(a+b,c)_\alpha = \phi_n(a,c)_\alpha + \phi_n(b,c)_\alpha$
- (iii) $\phi_n(a,b+c)_\alpha = \phi_n(a,b)_\alpha + \phi_n(a,c)_\alpha$
- (iv) $\phi_n(a,b)_{\alpha+\beta} = \phi_n(a,b)_\alpha + \phi_n(a,b)_\beta$

Proof:

(i) By Lemma (2.5) (i), we get:

$$\begin{aligned} d_n(aab + bab) &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))ad_j(\tau^{n-j}(b)) \\ d_n(aab) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a)) &= -(d_n(baa) - \sum_{i+j=n} d_i(\sigma^{n-i}(a))ad_j(\tau^{n-j}(b))) \\ \phi_n(a,b)_\alpha &= -\phi_n(b,a)_\alpha \\ (ii) \quad \phi_n(a+b,c)_\alpha &= d_n((a+b)ac) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(a+b)) \\ &= d_n(aac + bac) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(a)) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(b)) \\ &= d_n(aac) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(a)) + d_n(bac) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(b)) \\ &= \phi_n(a,c)_\alpha + \phi_n(b,c)_\alpha \\ (iii) \quad \phi_n(a,b+c)_\alpha &= d_n(aa(b+c)) - \sum_{i+j=n} d_i(\sigma^{n-i}(b+c))ad_j(\tau^{n-j}(a)) \\ &= d_n(aab + aac) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a)) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(a)) \\ &= d_n(aab) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a)) + d_n(aac) - \sum_{i+j=n} d_i(\sigma^{n-i}(c))ad_j(\tau^{n-j}(a)) \\ &= \phi_n(a,b)_\alpha + \phi_n(a,c)_\alpha \\ (iv) \quad \phi_n(a,b)_{\alpha+\beta} &= d_n(a(\alpha+\beta)b) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))(\alpha+\beta)d_j(\tau^{n-j}(a)) \\ &= d_n(aab) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))ad_j(\tau^{n-j}(a)) + d_n(a\beta b) - \sum_{i+j=n} d_i(\sigma^{n-i}(b)\beta)d_j(\tau^{n-j}(a)) \\ &= \phi_n(a,b)_\alpha + \phi_n(a,b)_\beta \end{aligned}$$

Remark (2.8):

Note that $D = (d_i)_{i \in N}$ is a (σ, τ) -higher reverse derivation of a Γ -ring M into itself if and only if $\phi_n = 0$, for all $n \in N$.

3- The Main Result :

Theorem (3.1):

Let $D = (d_i)_{i \in N}$ be a Jordan (σ, τ) -higher reverse derivation of a Γ -ring M into itself, then $\phi_n = 0$, for all $n \in N$.

Proof:

By Lemma (2.5) (i), we get

$$d_n(aab + baa) = \sum_{i+j=n} d_i(\sigma^{n-i}(b))\text{ad}_j(\tau^{n-j}(a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\text{ad}_j(\tau^{n-j}(b)) \quad \dots(1)$$

On the other hand

$$d_n(aab + baa) = d_n(aab) + d_n(baa) = d_n(aab) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\text{ad}_j(\tau^{n-j}(b)) \quad \dots(2)$$

Compare (1) and (2), we get:

$$\begin{aligned} d_n(aab) &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))\text{ad}_j(\tau^{n-j}(a)) \\ d_n(aab) - \sum_{i+j=n} d_i(\sigma^{n-i}(b))\text{ad}_j(\tau^{n-j}(a)) &= 0 \end{aligned}$$

By Definition (2.4), we get:

$\phi_n = 0$, for all $n \in N$.

Corollary (3.2):

Every Jordan (σ, τ) -higher reverse derivation of a Γ -ring M is a (σ, τ) -higher reverse derivation of M

Proof:

By Theorem (3.1), we get $\phi_n = 0$, for all $n \in N$ and by Remark (2.8) we get the require result.

Proposition (3.3):

Every Jordan (σ, τ) -higher reverse derivation of a 2-torsion free Γ -ring M into itself, such that $aab\beta a = a\beta baa$, for all $a, b \in M$ and $\alpha, \beta \in \Gamma$ is a Jordan triple (σ, τ) -higher reverse derivation .

Proof:

Let $D = (d_i)_{i \in N}$ be a Jordan (σ, τ) -higher reverse derivation of a Γ -ring M into itself.

Replace $a\beta b + b\beta a$ for b in Lemma (2.5) (i), we get:

$$\begin{aligned} d_n(a\alpha(a\beta b + b\beta a) + (a\beta b + b\beta a)\alpha a) \\ = d_n(a\alpha(a\beta b) + a\alpha(b\beta a) + (a\beta b)\alpha a + (b\beta a)\alpha a) \\ = d_n((a\alpha a)\beta b + (a\alpha b)\beta a + (a\beta b)\alpha a + (b\beta a)\alpha a) \end{aligned}$$

$$\begin{aligned}
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))\beta d_j(\tau^{n-j}(a\alpha a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\beta d_j(\tau^{n-j}(a\alpha b)) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(a\beta a)) + \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha d_j(\tau^{n-j}(b\beta a)) \\
 &= \sum_{i+j=n} d_i(\sigma^{n-i}(b))\beta (\sum_{r+s=j} d_r(\sigma^{j-r}\tau^{n-j}(a))\alpha d_s(\tau^{j-s}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\beta (\sum_{e+f=j} d_e(\sigma^{j-e}\tau^{n-j}(b))\alpha d_f(\tau^{j-f}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha (\sum_{p+q=j} d_p(\sigma^{j-p}\tau^{n-j}(b))\beta d_q(\tau^{j-q}\tau^{n-j}(a))) + \\
 &\quad \sum_{i+j=n} d_i(\sigma^{n-i}(a))\alpha (\sum_{x+y=j} d_x(\sigma^{j-x}\tau^{n-j}(a))\beta d_y(\tau^{j-y}\tau^{n-j}(b))) \\
 &= \sum_{i+r+s=n} d_i(\sigma^{n-i}(b))\beta d_r(\sigma^s\tau^i(a))\alpha d_s(\tau^{n-s}(a)) + \\
 &\quad \sum_{i+e+f=n} d_i(\sigma^{n-i}(a))\beta d_e(\sigma^f\tau^i(b))\alpha d_f(\tau^{n-f}(a)) + \\
 &\quad \sum_{i+p+q=n} d_i(\sigma^{n-i}(a))\alpha d_p(\sigma^q\tau^i(b))\beta d_q(\tau^{n-q}(a)) + \\
 &\quad \sum_{i+x+y=n} d_i(\sigma^{n-i}(a))\alpha d_x(\sigma^y\tau^i(a))\beta d_y(\tau^{n-y}(b)) \\
 &= d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k\tau^i(a))\alpha d_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\beta a\alpha b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha a\beta b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\alpha d_j(\sigma^k\tau^i(b))\beta d_k(\tau^{n-k}(a)) + \\
 &\quad d_n(a)\alpha b\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\alpha d_j(\sigma^k\tau^i(a))\beta d_k(\tau^{n-k}(b)) \\
 &= d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k\tau^i(a))\alpha d_k(\tau^{n-k}(a)) + \\
 &\quad 2(d_n(a)\beta a\alpha b + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k\tau^i(b))\alpha d_k(\tau^{n-k}(a))) + \\
 &\quad d_n(a)\alpha b\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\alpha d_j(\sigma^k\tau^i(a))\beta d_k(\tau^{n-k}(b))
 \end{aligned}$$

Since M is a 2-torsion free Γ -ring , then

$$= d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k\tau^i(a))\alpha d_k(\tau^{n-k}(a)) +$$



$$\begin{aligned}
 & d_n(a)\beta a ab + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a)) + \\
 & d_n(a)ab\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\text{ad}_j(\sigma^k \tau^i(a))\beta d_k(\tau^{n-k}(b))
 \end{aligned} \dots(1)$$

On the other hand :

$$\begin{aligned}
 & d_n(a\alpha(a\beta b + b\beta a) + (a\beta b + b\beta a)\alpha a) \\
 & = d_n(a\alpha a\beta b + a\alpha b\beta a + a\beta b\alpha a + b\beta a\alpha a)
 \end{aligned}$$

Since $aab\beta a = a\beta ba\alpha a$, for all $a, b \in M$ and $\alpha, \beta \in \Gamma$

$$\begin{aligned}
 & = d_n(a\alpha a\beta b + a\alpha b\beta a + a\beta b\alpha a + b\beta a\alpha a) \\
 & = d_n(b)\beta a\alpha a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(b))\beta d_j(\sigma^k \tau^i(a)) \text{ad}_k(\tau^{n-k}(a)) + \\
 & d_n(a)ab\beta a + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\text{ad}_j(\sigma^k \tau^i(a))\beta d_k(\tau^{n-k}(b)) + 2d_n(aab\beta a) \dots(2)
 \end{aligned}$$

Compare (1), (2) and since M is a 2-torsion free Γ -ring , we have :

$$d_n(aab\beta a) = d_n(a)\beta a ab + \sum_{i+j+k=n}^{i<n} d_i(\sigma^{n-i}(a))\beta d_j(\sigma^k \tau^i(b)) \text{ad}_k(\tau^{n-k}(a)).$$

References:

- [1] W.E.Barnes, "On The Γ -Rings of Nobusawa", Pacific J.Math., Vol.18(3)(1966) , pp.411-422 .
- [2] Y. Ceven and M.A. Ozturk, "On Jordan Generalized Derivations in Gamma Rings", Hacettepe Journal of Mathematics and Statistics, Vol.33(2004), pp.11-14.
- [3] S .Chakraborty and A.C .Paul, "On Jordan K-Derivations of 2-Torsion Free Prime Γ_N -Rings", Journal of Mathematics, Vol.40(2008), pp.97-101.
- [4] K.K.Dey ,A.C.Paul and I.S.Rakhimov "Semiprime Gamma Rings With Orthogonal Reverse Derivations", International Journal Of Pure and Applied Mathematics , Vol.83(2)(2013) , p.p 233 – 245 .
- [5] A.M. Kamal, " (σ,τ) -Derivations on Prime Γ -Rings ", M.Sc.Thesis, Department of Mathematics, college of Education, Al-Mustansirya University, 2012.
- [6] M.R.Salih,"Reverse Derivations on Prime Γ -Rings ", M.Sc.Thesis, Department of Mathematics, college of Education, Al-Mustansirya University, 2014.